

## Some Results of Ruin Probability for the Classical Risk Process<sup>†</sup>

HE YUANJIANG

*Department of Statistics Science, Zhongshan University, Ganglion, 510275 PRChina*

LI XUCHENG

*Department of Statistics Science, Zhongshan University, Ganglion, 510275 PRChina*

JOHN ZHANG\*

*Department of Mathematics, Indiana University of Pennsylvania, Indiana PA, 15705, USA*

**Abstract.** The computation of ruin probability is an important problem in the collective risk theory. It has applications in the fields of insurance, actuarial science, and economics. Many mathematical models have been introduced to simulate business activities and ruin probability is studied based on these models. Two of these models are the classical risk model and the Cox model. In the classical model, the counting process is a Poisson process and in the Cox model, the counting process is a Cox process. Thorin (1973) studied the ruin probability based on the classical model with the assumption that random sequence followed the  $\Gamma$  distribution with density function

$$f(x) = \frac{x^{\frac{1}{\beta}-1}}{\beta^{\frac{1}{\beta}} \Gamma(1/\beta)} e^{-\frac{x}{\beta}}, x > 0, \text{ where } \beta > 1. \text{ This paper studies the ruin probability of}$$

the classical model where the random sequence follows the  $\Gamma$  distribution with density function  $f(x) = \frac{\alpha^n}{\Gamma(n)} x^{n-1} e^{-\alpha x}$ ,  $x > 0$ , where  $\alpha > 0$  and  $n \geq 2$  is a positive integer. An intermediate general result is given and a complete solution is provided for  $n = 2$ . Simulation studies for the case of  $n = 2$  is also provided.

**Keywords:** Collective risk theory; Gamma distribution; Ruin probability; Simulation

### 1. Introduction

To study the probability of ruin, many models have been proposed and investigated. These models include the classical and the Cox model. Although the applications of ruin probability are not restricted to the insurance type problem, the idea can be explained easily by an insurance business model. An insurance company's capital at time  $t$  can be expressed as

---

<sup>†</sup> Supported in part by National Natural Science Foundation of China and China Bridges International

\* Requests for reprints should be sent to John Zhang, Department of Mathematics, Indiana University of Pennsylvania 233 Stright Hall, Indiana PA, 15705, USA.

$R(t) = u + C(t) - O(t)$ , where  $R(t)$  is the company's capital at time  $t$ ,  $u$  is the initial capital,  $C(t)$  is the income function, and  $O(t)$  is the expense function. The income function  $C(t)$  can be expressed as a relatively simple function and often taken to be  $C(t) = ct$ , where  $c > 0$  is the income coefficient. The expense function  $O(t)$ , however, is a random process because of the nature of the insurance claims. One way to describe  $O(t)$  is to define:

$$O(t) = \sum_{i=1}^{N(t)} Z_i,$$

where  $Z_i$  is the size of the payment and  $N(t)$  is a counting process. The classical model assumes that  $N(t)$  is a Poisson process and the Cox model assumes that  $N(t)$  is a Cox process. Thorin (1973) studied the classical model where  $Z_i$  has  $\Gamma$  distribution with density function

$$f(x) = \frac{x^{\frac{1}{\beta}-1}}{\beta^{\frac{1}{\beta}} \Gamma(1/\beta)} e^{-\frac{x}{\beta}}, \quad x > 0,$$

where  $\beta > 1$ . In this paper, we study the classical model when  $Z_i$  follows the  $\Gamma$  distribution with density function

$$f(x) = \frac{\alpha^n}{\Gamma(n)} x^{n-1} e^{-\alpha x}, \quad x > 0,$$

where  $\alpha > 0$  and  $n \geq 2$  is a positive integer.

Thorin's case models situations where the probability distribution of ten random payments  $Z_i$  is monotonically strictly decrease. The case we study here can be applied to situations when the probability distribution of the random payments  $Z_i$  places higher mass to values near the center. It is meaningful to study this case because many data sets in practice have mound shape distributions.

The organization of the paper is as follows: In section 2, we give the main theorems and section 3 presents the simulation study. Section 4 contains the concluding remarks.

## 2. Main Results

We define a risk process as

$$X(t) = ct - \sum_{k=1}^{N(t)} Z_k,$$

where  $N(t)$  is a counting process,  $\{Z_k\}_1^\infty$  is a sequence of independent random variables with identical distributions. We further assume that the common distribution function is  $F$  with mean  $\mu$  and variance  $\sigma^2$ , and  $F(0) = 0$ . The counting process  $N(t)$  and  $\{Z_k\}_1^\infty$  are independent, and  $c$  is a positive constant.

$N(t)$  can be interpreted as the number of claims of an insurance company in the time interval  $(0, t]$ . At each jump point of  $N(t)$ , the insurance company has to pay out a stochastic amount of money. On the other hand, the company receives  $c$  units of revenue per unit time.

Let  $\Psi(u) = P\{u + X(t) < 0 \text{ for some } t > 0\}$ . Then for  $u \geq 0$ ,  $\Psi(u)$  can be interpreted as the ruin probability of an insurance company which has initial capital  $u$  when facing the risk process  $X(t)$ .  $\Phi(u) = 1 - \Psi(u)$  is the non-ruin probability. Notice that  $\Psi(u) = 1$  for  $u < 0$ .

From now on, we assume that  $N(t)$  is a Poisson process with standing  $\lambda > 0$ , and therefore,  $E(N(t)) = \lambda t$ . Let

$$\rho = \frac{c - \lambda\mu}{\lambda\mu}.$$

Then

$$EX(t) = 3Dct - E \sum_{i=3D1}^{N(t)} Z_i = 3Dct - EN(t)EZ_1 = 3Dct - \lambda\mu t = \rho\lambda\mu t.$$

We can see that

$$\rho = \frac{EX(t)}{\lambda\mu t},$$

and is called the safety loading. It is the ratio of the expected profit to the expected payments.  $\rho = 1$ , for example, indicates that the company's expected profit is of the same size as its expected payments. Larger value of  $\rho$  indicates a company is in a "save" state. Thus  $\rho$  can be considered as an index to measure the safety (non-ruin) of an insurance company.

We also assume that

$$\rho > 0.$$

This condition can be interpreted as the premiums received per unit time exceed the expected claim payments per unit time.

If  $Z_i$  has exponential distribution, then

$$\Psi(u) = \frac{1}{1 + \rho} e^{-\frac{\rho u}{\mu(1+\rho)}}$$

(for example, see Grandell(1991)). If  $Z_i$  has  $\Gamma$  distribution with density function

$$f(x) = \frac{x^{\frac{1}{\beta}-1}}{\beta^{\frac{1}{\beta}}\Gamma(1/\beta)} e^{-\frac{x}{\beta}}, \quad x > 0,$$

where  $\beta > 1$ , then the ruin probability is

$$\Psi(u) = \frac{\rho(1 - \beta R)e^{-Ru}}{1 + (1 + \rho)(R + R\beta - 1)} + \frac{\rho}{\pi\beta} \sin \frac{\pi}{\beta} \int_0^\infty \frac{x^{\frac{1}{\beta}} e^{-(x+1)u/\beta} dx}{\{x^{\frac{1}{\beta}} [1 + (1 + \rho)\frac{x+1}{\beta}] - \cos \frac{\pi}{\beta}\}^2 + \sin^2 \frac{\pi}{\beta}},$$

where  $R$  is the positive solution of the equation

$$\int_0^{+\infty} e^{rz} dF(z) - 1 = \frac{cr}{\lambda}$$

for  $r < 1/\beta$ , that is, the positive solution of the equation

$$(1 - \beta r)^{-\frac{1}{\beta}} - 1 = (1 + \rho)r$$

for  $r < 1/\beta$  (for example, see Thorin (1973), or Grandell (1991), p.14).

Now we consider the case that  $Z_i$  has  $\Gamma$  distribution with density function

$$f(x) = \frac{\alpha^n}{\Gamma(n)} x^{n-1} e^{-\alpha x}, \quad x > 0,$$

where  $\alpha > 0$  and  $n$  is a positive integer.

We state a well-known result as a lemma:

**Lemma 1** For a classical risk model, if  $N(t)$  has intensity  $\lambda$  and  $Z_i$  has distribution function  $F(z)$ , then

$$D\Phi(u) = \frac{\lambda}{c}\Phi(u) - \frac{\lambda}{c} \int_0^u \Phi(u-z) dF(z).$$

Here,  $D = \frac{d}{du}$  is the differential operator.

(Interested readers please see equation (3) in Grandell (1991), p.4.)

**Theorem 1** For a classical risk model, if  $N(t)$  has intensity  $\lambda$  and  $Z_i$  has  $\Gamma$  distribution with density function

$$f(x) = \frac{\alpha^n}{\Gamma(n)} x^{n-1} e^{-\alpha x}, \quad x > 0,$$

where  $\alpha > 0$  and  $n \geq 1$  is an integer, then the non-ruin probability  $\Phi(u)$  is a solution of the ordinary differential equation:

$$D(D + \alpha)^n \Phi(u) - \frac{\lambda}{c}(D + \alpha)^n \Phi(u) + \frac{\lambda}{c} \alpha^n \Phi(u) = 0. \quad (1)$$

**Proof.** From Lemma 1, we have

$$D\Phi(u) = \frac{\lambda}{c}\Phi(u) - \frac{\lambda}{c} \int_0^u \Phi(u-z) dF(z). \quad (2)$$

Substituting  $dF(z) = f(z)dz$ , we have

$$D\Phi(u) = \frac{\lambda}{c}\Phi(u) - \frac{\lambda}{c} \int_0^u \Phi(u-z) \frac{\alpha^n}{\Gamma(n)} z^{n-1} e^{-\alpha z} dz.$$

The change of variables  $z = u - w$  leads to

$$D\Phi(u) = \frac{\lambda}{c}\Phi(u) - \frac{\lambda}{c} \int_0^u \Phi(w) \frac{\alpha^n}{\Gamma(n)} (u-w)^{n-1} e^{-\alpha(u-w)} dw. \quad (3)$$

Let  $A(u; 0) = \Phi(u)$  and let

$$A(u; k) = \int_0^u \Phi(w) \frac{\alpha^k}{\Gamma(k)} (u-w)^{k-1} e^{-\alpha(u-w)} dw \quad (4)$$

for  $1 \leq k \leq n$ .

By (2),  $\Phi(w)$  is differentiable. Let  $h(w, \eta) = \Phi(w)\alpha e^{-\alpha(\eta-w)}$ . Then  $h(w, \eta)$  and  $\frac{\partial h(w, \eta)}{\partial \eta}$  are continuous. Let

$$g(\xi(u), \eta(u)) = \int_0^{\xi(u)} \Phi(w)\alpha e^{-\alpha(\eta(u)-w)} dw, \quad \xi(u) = u, \quad \eta(u) = u.$$

Then by Lang (1979), p.119, Theorem 5 of Chapter V, we can exchange the order of differentiation and integration, so we have

$$\begin{aligned} DA(u; 1) &= Dg(\xi, \eta) \\ &= \frac{\partial g}{\partial \xi} \frac{d\xi}{du} + \frac{\partial g}{\partial \eta} \frac{d\eta}{du} \\ &= \alpha \Phi(\xi) e^{-\alpha(\eta-\xi)} \\ &\quad - \alpha \int_0^\xi \Phi(w) \alpha e^{-\alpha(\eta-w)} dw \\ &= \alpha \Phi(u) - \alpha \int_0^u \Phi(w) \alpha e^{-\alpha(u-w)} dw \\ &= \alpha A(u; 0) - \alpha A(u; 1). \end{aligned} \quad (5)$$

And by using similar methods, for  $2 \leq k \leq n$ , we have

$$\begin{aligned} & DA(u; k) \\ &= (k-1) \int_0^u \Phi(w) \frac{\alpha^k}{\Gamma(k)} (u-w)^{k-2} e^{-\alpha(u-w)} dw \\ &\quad - \alpha \int_0^u \Phi(w) \frac{\alpha^k}{\Gamma(k)} (u-w)^{k-1} e^{-\alpha(u-w)} dw \\ &= \alpha A(u; k-1) - \alpha A(u; k). \end{aligned} \quad (6)$$

So for  $1 \leq k \leq n$ , we have

$$\frac{D+\alpha}{\alpha} A(u; k) = A(u; k-1).$$

Thus

$$\left( \frac{D+\alpha}{\alpha} \right)^n A(u; n) = A(u; 0) = \Phi(u). \quad (7)$$

From (3) and (4), we have

$$D\Phi(u) = \frac{\lambda}{c} \Phi(u) - \frac{\lambda}{c} A(u; n). \quad (8)$$

From (5) and (6), we have

$$\begin{aligned} D \left( \frac{D+\alpha}{\alpha} \right)^n \Phi(u) &= \left( \frac{D+\alpha}{\alpha} \right)^n D\Phi(u) \\ &= \frac{\lambda}{c} \left( \frac{D+\alpha}{\alpha} \right)^n \Phi(u) - \frac{\lambda}{c} \left( \frac{D+\alpha}{\alpha} \right)^n A(u; n) \\ &= \frac{\lambda}{c} \left( \frac{D+\alpha}{\alpha} \right)^n \Phi(u) - \frac{\lambda}{c} \Phi(u). \end{aligned} \quad (9)$$

So

$$D(D+\alpha)^n \Phi(u) - \frac{\lambda}{c} (D+\alpha)^n \Phi(u) + \frac{\lambda}{c} \alpha^n \Phi(u) = 0.$$

□

**Remark.** If  $n = 1$ , equation (1) becomes

$$D(D+\alpha)\Phi(u) - \frac{\lambda}{c}(D+\alpha)\Phi(u) + \frac{\lambda\alpha}{c}\Phi(u) = 0.$$

After simplification, we have

$$D^2\Phi(u) + \left(\alpha - \frac{\lambda}{c}\right)D\Phi(u) = 0,$$

or

$$D^2\Phi(u) = -\frac{\alpha\rho}{1+\rho}D\Phi(u).$$

The above equation is solved in Grandell (1991), page 6.  $\square$

The following theorem gives a complete solution of differential equation (1) for  $n = 2$ .

**Theorem 2** *For a classical risk model, if  $N(t)$  has intensity  $\lambda$  and  $Z_i$  has  $\Gamma$  distribution with density function*

$$f(x) = \frac{\alpha^2}{\Gamma(2)}xe^{-\alpha x}, \quad x > 0,$$

then the non-ruin probability  $\Phi(u)$  is :

$$\Phi(u) = 1 + \frac{\nu_2(\nu_1 + \alpha)^2}{(\nu_1 - \nu_2)\alpha^2}e^{\nu_1 u} + \frac{\nu_1(\nu_2 + \alpha)^2}{(\nu_2 - \nu_1)\alpha^2}e^{\nu_2 u},$$

where

$$\nu_1 = \frac{\lambda - 2c\alpha + \sqrt{\lambda^2 + 4c\alpha\lambda}}{2c},$$

$$\nu_2 = \frac{\lambda - 2c\alpha - \sqrt{\lambda^2 + 4c\alpha\lambda}}{2c}.$$

**Proof.** By Theorem 1,  $\Phi(u)$  is a solution of the ordinary differential equation:

$$D(D + \alpha)^2\Phi(u) - \frac{\lambda}{c}(D + \alpha)^2\Phi(u) + \frac{\lambda}{c}\alpha^2\Phi(u) = 0. \quad (10)$$

The characteristic equation

$$x(x + \alpha)^2 - \frac{\lambda}{c}(x + \alpha)^2 + \frac{\lambda}{c}\alpha^2 = 0 \quad (11)$$

can be rewritten as

$$x[x^2 + (2\alpha - \frac{\lambda}{c})x + (\alpha^2 - \frac{2\alpha\lambda}{c})] = 0, \quad (12)$$

and has solutions

$$\nu_0 = 0,$$

$$\nu_1 = \frac{\lambda - 2c\alpha + \sqrt{\lambda^2 + 4c\alpha\lambda}}{2c},$$

$$\nu_2 = \frac{\lambda - 2c\alpha - \sqrt{\lambda^2 + 4c\alpha\lambda}}{2c}.$$

And (7) has general solution

$$\Phi(u) = c_0 + c_1 e^{\nu_1 u} + c_2 e^{\nu_2 u}. \quad (13)$$

Since

$$\rho = \frac{c - \lambda\mu}{\lambda\mu} > 0, \quad \mu = \frac{2}{\alpha},$$

we have

$$\frac{c\alpha - 2\lambda}{2\lambda} > 0.$$

So

$$\alpha > \frac{2\lambda}{c},$$

and hence

$$2\alpha > \alpha > \frac{2\lambda}{c} > \frac{\lambda}{c}.$$

Thus

$$\alpha^2 - \frac{2\alpha\lambda}{c} > 0 \quad \text{and} \quad 2\alpha - \frac{\lambda}{c} > 0.$$

Since  $\nu_1$  and  $\nu_2$  are the solutions of equation (9), we have

$$\nu_1 < 0 \quad \text{and} \quad \nu_2 < 0.$$

Thus, let  $u \rightarrow \infty$ , we have:

$$c_0 = \Phi(\infty) = 1. \quad (14)$$

From (3), (10) and (11), we have

$$\begin{aligned} & c_1 \nu_1 e^{\nu_1 u} + c_2 \nu_2 e^{\nu_2 u} \\ = & \frac{\lambda}{c} (1 + c_1 e^{\nu_1 u} + c_2 e^{\nu_2 u}) - \frac{\lambda}{c} \int_0^u (1 + c_1 e^{\nu_1 w} + c_2 e^{\nu_2 w}) \frac{\alpha^2}{\Gamma(2)} (u-w) e^{-\alpha(u-w)} dw. \end{aligned}$$

After simplification, we have

$$\begin{aligned} & [\nu_1 - \frac{\lambda}{c} + \frac{\lambda\alpha^2}{c(\nu_1 + \alpha)^2}] c_1 e^{\nu_1 u} + [\nu_2 - \frac{\lambda}{c} + \frac{\lambda\alpha^2}{c(\nu_2 + \alpha)^2}] c_2 e^{\nu_2 u} \\ = & \frac{\lambda}{c} e^{-\alpha u} \left\{ (\alpha u + 1) + \alpha^2 \left\{ \frac{c_1}{(\nu_1 + \alpha)^2} [(\nu_1 + \alpha)u + 1] + \right. \right. \\ & \left. \left. \frac{c_2}{(\nu_2 + \alpha)^2} [(\nu_2 + \alpha)u + 1] \right\} \right\}. \quad (15) \end{aligned}$$



From (8), we have

$$\nu_1 - \frac{\lambda}{c} + \frac{\lambda\alpha^2}{c(\nu_1 + \alpha)^2} = 0$$

and

$$\nu_2 - \frac{\lambda}{c} + \frac{\lambda\alpha^2}{c(\nu_2 + \alpha)^2} = 0.$$

So

$$\alpha u + 1 + \alpha^2 \left\{ \frac{c_1}{(\nu_1 + \alpha)^2} [(\nu_1 + \alpha)u + 1] + \frac{c_2}{(\nu_2 + \alpha)^2} [(\nu_2 + \alpha)u + 1] \right\} = 0,$$

that is,

$$\left( \alpha + \frac{\alpha^2 c_1}{\nu_1 + \alpha} + \frac{\alpha^2 c_2}{\nu_2 + \alpha} \right) u + \frac{\alpha^2 c_1}{(\nu_1 + \alpha)^2} + \frac{\alpha^2 c_2}{(\nu_2 + \alpha)^2} + 1 = 0.$$

Therefore we have

$$\alpha + \frac{\alpha^2 c_1}{\nu_1 + \alpha} + \frac{\alpha^2 c_2}{\nu_2 + \alpha} = 0$$

and

$$\frac{\alpha^2 c_1}{(\nu_1 + \alpha)^2} + \frac{\alpha^2 c_2}{(\nu_2 + \alpha)^2} + 1 = 0.$$

The solution is

$$c_1 = \frac{\nu_2(\nu_1 + \alpha)^2}{(\nu_1 - \nu_2)\alpha^2}, \quad c_2 = \frac{\nu_1(\nu_2 + \alpha)^2}{(\nu_2 - \nu_1)\alpha^2}.$$

Thus we have

$$\Phi(u) = 1 + \frac{\nu_2(\nu_1 + \alpha)^2}{(\nu_1 - \nu_2)\alpha^2} e^{\nu_1 u} + \frac{\nu_1(\nu_2 + \alpha)^2}{(\nu_2 - \nu_1)\alpha^2} e^{\nu_2 u}.$$

□

Theorem 2 has an interesting equivalent version. It is stated as follows:

**Theorem 3** *For a classical risk model, if  $N(t)$  has intensity  $\lambda = 1$  and  $Z_i$  has  $\Gamma$  distribution with density function*

$$f(x) = \frac{1}{\Gamma(2)} x e^{-x}, \quad x > 0,$$

*then the non-ruin probability  $\Phi(u)$  is:*

$$\Phi(u) = 1 + \frac{\nu_2(\nu_1 + 1)^2}{\nu_1 - \nu_2} e^{\nu_1 u} + \frac{\nu_1(\nu_2 + 1)^2}{\nu_2 - \nu_1} e^{\nu_2 u}, \quad (16)$$

where

$$\nu_1 = \frac{1 - 2c + \sqrt{1 + 4c}}{2c},$$

$$\nu_2 = \frac{1 - 2c - \sqrt{1 + 4c}}{2c}.$$

□

It is obvious that Theorem 2 implies Theorem 3 (by letting  $\lambda = 1, \alpha = 1$ ). Now we show that Theorem 3 implies Theorem 2 as follows:

Suppose that Theorem 3 holds. For a classical risk model, let  $N(t)$  be a Poisson process with intensity  $\lambda$ , and assume all  $Z_i$ 's have  $\Gamma$  distribution with identical density function

$$f(z) = \frac{\alpha^2}{\Gamma(2)} z e^{-\alpha z}, \quad z > 0.$$

Let

$$\tilde{N}(t) = N(t/\lambda),$$

and

$$\tilde{Z}_i = \alpha Z_i,$$

for all  $i$ . Then  $\tilde{N}(t)$  is a Poisson process with intensity 1, all  $\tilde{Z}_i$ s have identical density function

$$\tilde{f}(\tilde{z}) = \frac{1}{\Gamma(2)} \tilde{z} e^{-\tilde{z}}, \quad \tilde{z} > 0.$$

Let

$$R(t) = u + ct - \sum_{i=1}^{N(t)} Z_i,$$

and

$$\tilde{R}(t) = \tilde{u} + \tilde{c}t - \sum_{i=1}^{\tilde{N}(t)} \tilde{Z}_i,$$

where

$$\tilde{u} = \alpha u, \quad \tilde{c} = \frac{\alpha c}{\lambda}.$$

Then

$$\tilde{R}(\lambda t) = \alpha u + \frac{c\alpha}{\lambda} \cdot \lambda t - \alpha \sum_{i=1}^{\tilde{N}(\lambda t)} Z_i = \alpha u + \alpha ct - \alpha \sum_{i=1}^{N(t)} Z_i = \alpha R(t). \quad (17)$$

Let

$$\left. \begin{aligned} \Phi(u) &= 1 - P\{R(t) < 0 \text{ for some } t\} \\ \tilde{\Phi}(\tilde{u}) &= 1 - P\{\tilde{R}(t) < 0 \text{ for some } t\} \end{aligned} \right\}.$$

Then by (13) and Theorem 3 we have

$$\Phi(u) = \tilde{\Phi}(\tilde{u}) = 1 + \frac{\nu_2(\nu_1 + 1)^2}{\nu_1 - \nu_2} e^{\nu_1 \tilde{u}} + \frac{\nu_1(\nu_2 + 1)^2}{\nu_2 - \nu_1} e^{\nu_2 \tilde{u}}, \quad (18)$$

where

$$\begin{aligned} \nu_1 &= \frac{1 - 2\tilde{c} + \sqrt{1 + 4\tilde{c}}}{2\tilde{c}}, \\ \nu_2 &= \frac{1 - 2\tilde{c} - \sqrt{1 + 4\tilde{c}}}{2\tilde{c}}. \end{aligned}$$

It is easy to see that Theorem 2 comes from (14) if we replace  $\tilde{u}$  and  $\tilde{c}$  by  $\alpha u$  and  $\alpha c/\lambda$ , respectively.  $\square$

### 3. Simulation Study

For the risk process in Theorem 3, we conducted a simulation study to demonstrate the relationships between the non-ruin probability, the initial capital and the revenue coefficient  $c$ . The simulation is carried out using SAS version 6.12 on a PC with CPU PII-300. In example 1, we simulate the risk process 1000000 times, while in example 2, for each case (observation 1-18) we simulate the risk process 10000 times. The following is part of the program and the results:

```
Example 1: data new;
d =10; c =2.4; u =5; r =0; nr =0; w =u;
v1 =(1-2*c+sqrt(1+4*c))/(2*c);
v2 =(1-2*c-sqrt(1+4*c))/(2*c);
p =1+(v2*(v1+1)**2*exp(v1*w))/(v1-v2)+(v1*(v2+1)**2*exp(v2*w))/(v2-
v1);
do i =1 to 1000000;
w =u;
```

```

k: w =w+c*ranexp(1)-rangam(d,2);
if w >=0 and w <=500 then goto k;
if w > 500 then nr =nr+1;
if w < 0 then r =r+1;
end;
f =nr/(r+nr);
e =f-p;
s =1000*e/sqrt(p*(1-p));
proc print;
var c u nr f p e s;
run;

```

OBS	C	U	NR	F	P	E	S
1	2.4	5	516550	0.51655	0.51681	-.00026197	-0.52424

NOTE: The DATA statement used 4 hours 14 minutes 33.72seconds.  
 Example 2:

OBS	C	U	NR	F	P	E	S
1	2.1	3	1350	0.1350	0.12984	0.0051617	1.53565
2	2.1	5	1803	0.1803	0.18360	-.0032976	-0.85176
3	2.1	10	3027	0.3027	0.30402	-.0013219	-0.28738
4	2.1	50	8043	0.8043	0.80586	-.0015609	-0.39464
5	2.1	100	9588	0.9588	0.96064	-.0018441	-0.94842
6	2.1	200	9985	0.9985	0.99838	0.0001173	0.29203
7	2.2	3	2302	0.2302	0.23523	-.0050277	-1.18538
8	2.2	5	3197	0.3197	0.32330	-.0036049	-0.77071
9	2.2	10	5021	0.5021	0.50181	0.0002863	0.05727
10	2.2	50	9565	0.9565	0.95701	-.0005116	-0.25223
11	2.2	100	9979	0.9979	0.99799	-.0000895	-0.19984
12	2.2	200	10000	1.0000	1.00000	0.0000044	0.20970
13	2.4	3	3911	0.3911	0.39403	-.0029332	-0.60029
14	2.4	5	5223	0.5223	0.51681	0.0054880	1.09823
15	2.4	10	7321	0.7321	0.72589	0.0062069	1.39148
16	2.4	50	9969	0.9969	0.99706	-.0001606	-0.29658
17	2.4	100	10000	1.0000	0.99999	0.0000101	0.31849
18	2.4	200	10000	1.0000	1.00000	0.0000000	0.00110

Here, C and U are the c and u in Theorem 3, respectively. (Recall that u is the initial capital of an insurance company and the company receives c units of revenue per unit time.) P is the non-ruin probability  $\Phi(u)$  in

Theorem 3 calculated by using equation (12); NR is the number of the non-ruin risk processes; F is the relative frequency of non-ruin, and E is the difference of F and P. E can be interpreted as the error of simulation results (assuming our theory is correct.) The small E values in our study confirm our theoretical results from a different perspective.

For the estimation of the ruin probability  $\Psi(u)$ , we can use the Lundberg inequality

$$\Psi(u) \leq e^{-Ru}, \quad (19)$$

where  $R$  is the Lundberg exponent, that is,  $R$  is the positive solution of the equation

$$\int_0^{+\infty} e^{rz} dF(z) - 1 = \frac{cr}{\lambda}. \quad (20)$$

(For example, see Grandell, p.11.) For the risk process in Theorem 3, equation (16) becomes

$$\int_0^{+\infty} e^{rz} z e^{-z} dz - 1 = cr.$$

If  $r \geq 1$ , then

$$\int_0^{+\infty} e^{rz} z e^{-z} dz = +\infty.$$

If  $r < 1$ , then we have

$$\frac{1}{(1-r)^2} - 1 = cr,$$

or

$$r[cr^2 - (2c-1)r + (c-2)] = 0. \quad (21)$$

Since  $\rho > 0$ , it follows that  $c > 2$ . Hence

$$R = \frac{2c-1 - \sqrt{(2c-1)^2 - 4c(c-2)}}{2c} = \frac{2c-1 - \sqrt{4c+1}}{2c}$$

is the unique solution of equation (17) satisfying  $0 < R < 1$ . For  $c = 2.1, 2.2$  and  $2.4$ , we have  $R = 0.03191, 0.06125$  and  $0.11338$  respectively.  $\Psi(u)$  is strictly decreasing in  $R$  for fixed  $u > 0$ , it is suffice to examine  $R = 0.03191$ . So if  $u > 500$ , then by (15), the ruin probability

$$\Psi(u) \leq e^{-0.03191 \times 500} = 1.1771 \times 10^{-7}$$

is sufficiently small. So if  $\inf\{t : R(t) > 500\} < \inf\{t : R(t) < 0\}$ , we classify the company as "non-ruin". Otherwise, if  $\inf\{t : R(t) < 0\} < \inf\{t : R(t) > 500\}$ , we consider the company as "ruin". In other words, in

our example, "non-ruin" means that the time for the total capital exceeding 500 comes prior to the time the total capital being less than 0. This assumption is reasonable because once the total capital exceeds 500, the chance of "ruin" is slim.

We consider the hypothesis test problem  $H_0 : \Phi(u) = P$ . If  $H_0$  is true, then  $S = E\sqrt{R + NR}/\sqrt{P(1 - P)}$  has approximately a standard normally distribution,  $P(-1.96 \leq S \leq 1.96) = 0.95$ . (Note that  $\sqrt{(R + NR)} = \sqrt{(10^6)} = 1000$  in Example 1, while in Example 2,  $\sqrt{(R + NR)} = \sqrt{(10^4)} = 100$ .) The simulation results show that the relative frequency of ruin (i.e.,  $F$ ) is very close to the probability of ruin computed by Theorem 3 (i.e.,  $P$ ), and all  $S$  values fall in the interval  $[-1.96, 1.96]$ . These results confirm that Theorem 3 is correct.

#### 4. Concluding Remarks

The computation of a ruin probability is in many cases difficult. This paper gives a way of computing the ruin probability in a special case. Although the result of Theorem 1 is for gamma distributions with  $n$  being positive integer, we are currently only able to give a complete solution for the case of  $n = 2$ . We have obtained solutions for the characteristic equations corresponding to equation (1) in Theorem 1 for  $n = 3$  and 4. The computation of the ruin probabilities for these cases is ongoing.

#### References

1. Grandell, J., (1991) Aspects of Risk Theory. Springer-Verlag, New York.
2. S. Lang, (1979) Calculus of several variables, second edition. Addison-Wesley Publishing Company, Massachusetts.
3. Thorin, O., (1973) The ruin problem in case the tail of the claim distribution is completely monotone. Skand. Aktuar Tidskr., 100-119.

## Special Issue on Singular Boundary Value Problems for Ordinary Differential Equations

### Call for Papers

The purpose of this special issue is to study singular boundary value problems arising in differential equations and dynamical systems. Survey articles dealing with interactions between different fields, applications, and approaches of boundary value problems and singular problems are welcome.

This Special Issue will focus on any type of singularities that appear in the study of boundary value problems. It includes:

- Theory and methods
- Mathematical Models
- Engineering applications
- Biological applications
- Medical Applications
- Finance applications
- Numerical and simulation applications

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/bvp/guidelines.html>. Authors should follow the Boundary Value Problems manuscript format described at the journal site <http://www.hindawi.com/journals/bvp/>. Articles published in this Special Issue shall be subject to a reduced Article Processing Charge of €200 per article. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	May 1, 2009
First Round of Reviews	August 1, 2009
Publication Date	November 1, 2009

### Lead Guest Editor

**Juan J. Nieto**, Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de

Compostela, Santiago de Compostela 15782, Spain;  
[juanjose.nieto.roig@usc.es](mailto:juanjose.nieto.roig@usc.es)

### Guest Editor

**Donal O'Regan**, Department of Mathematics, National University of Ireland, Galway, Ireland;  
[donal.oregan@nuigalway.ie](mailto:donal.oregan@nuigalway.ie)