

RELIABILITY FOR SOME BIVARIATE GAMMA DISTRIBUTIONS

SARALEES NADARAJAH

Received 23 July 2004 and in revised form 23 November 2004

In the area of stress-strength models, there has been a large amount of work as regards estimation of the reliability $R = \Pr(X < Y)$. The algebraic form for $R = \Pr(X < Y)$ has been worked out for the vast majority of the well-known distributions when X and Y are independent random variables belonging to the same univariate family. In this paper, we consider forms of R when (X, Y) follows a bivariate distribution with dependence between X and Y . In particular, we derive explicit expressions for R when the joint distribution is bivariate gamma. The calculations involve the use of special functions.

1. Introduction

Bivariate gamma distributions arise as tractable “lifetime” models in many areas, including life testing and telecommunications. In the context of reliability, the stress-strength model describes the life of a component which has a random strength Y and is subjected to a random stress X . The component fails at the instant that the stress applied to it exceeds the strength and the component will function satisfactorily whenever $Y > X$. Thus, $R = \Pr(X < Y)$ is a measure of the component reliability. It has many applications especially in engineering concepts such as structures, deterioration of rocket motors, static fatigue of ceramic components, fatigue failure of aircraft structures, and the aging of concrete pressure vessels. Some examples are as follows.

- (i) If X represents the maximum chamber pressure generated by ignition of a solid propellant and Y represents the strength of the rocket chamber, then R is the probability of successful firing of the engine.
- (ii) If X represents the diameter of a shaft and Y represents the diameter of a bearing that is to be mounted on the shaft, then R is the probability that the bearing fits without interference.
- (iii) Let Y and X be the remission times of two chemicals when they are administered to two kinds of mechanical systems. Inferences about R present a comparison of the effectiveness of the two chemicals.
- (iv) If X and Y are future observations on the stability of an engineering design, then R would be the predictive probability that X is less than Y . Similarly, if X and

Y represent lifetimes of two electronic devices, then R is the probability that one fails before the other.

- (v) If Y represents the distance of a pyrotechnic igniter from its adjacent pellet and X represents its *ignition distance*, then R is the probability that the igniter succeeds to bridge the gap in the pyrotechnic chain.
- (vi) A certain unit—be it a receptor in a human eye or ear or any other organ (including sexual)—operates only if it is stimulated by the source of random magnitude Y and the stimulus exceeds a lower threshold X specific for that unit. In this case, R is the probability that the unit functions.
- (vii) In military warfare, R could be interpreted as the probability that a given round of ammunition will penetrate its target.

Because of these applications, the calculation and the estimation of $R = \Pr(X < Y)$ are important for the class of bivariate gamma distributions. The calculation of R has been extensively investigated in the literature when X and Y are independent random variables belonging to the same univariate family of distributions. The algebraic form for R has been worked out for the majority of the well-known distributions, including normal, uniform, exponential, gamma, beta, extreme value, Weibull, Laplace, logistic, and the Pareto distributions (Nadarajah [21, 22, 23, 24, 25]; Nadarajah and Kotz [27]). However, there is relative little work when X and Y are dependent random variables. A complete literature search shows that the cases of bivariate beta (Nadarajah [26]), bivariate exponential (Awad et al. [2], Klein and Basu [18], Jana [13], Jana and Roy [15], Jana [14], Jeevanand [17], Hanagal [10, 12]), bivariate normal (Mukherjee and Saran [20], Nandi and Aich [28, 29], Gupta and Subramanian [8]), and bivariate Pareto (Hanagal [11], Jeevanand [16]) have been considered. The cases of multivariate exponential (Cramer and Kamps [5]) and multivariate normal (Singh [33], R. D. Gupta and R. C. Gupta [9]) have also been considered. We have been able to trace no other work for dependent variables.

The aim of this paper is to calculate R when X and Y are dependent random variables from six flexible families of bivariate gamma distributions. We will assume throughout this paper that (X, Y) has a bivariate gamma distribution with joint probability density function (pdf) f and joint survivor function \bar{F} . One can write

$$R = \int_0^{\infty} \int_x^{\infty} f(x, y) dy dx. \quad (1.1)$$

Our calculations of R make use of a number of special functions. They are the complementary incomplete gamma function defined by

$$\Gamma(a, x) = \int_x^{\infty} t^{a-1} \exp(-t) dt, \quad (1.2)$$

the confluent hypergeometric function (${}_1F_1$) defined by

$${}_1F_1(a; b; x) = \sum_{k=0}^{\infty} \frac{(a)_k}{(b)_k} \frac{x^k}{k!}, \quad (1.3)$$

the Gauss hypergeometric function (${}_2F_1$) defined by

$${}_2F_1(a, b; c; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k} \frac{x^k}{k!}, \tag{1.4}$$

the generalized hypergeometric function (${}_2F_2$) defined by

$${}_2F_2(a, b; c, d; x) = \sum_{k=0}^{\infty} \frac{(a)_k (b)_k}{(c)_k (d)_k} \frac{x^k}{k!}, \tag{1.5}$$

the Kummer function defined by

$$\Psi(a, b, x) = \frac{1}{\Gamma(a)} \int_0^{\infty} t^{a-1} (1+t)^{b-a-1} \exp(-zt) dt, \tag{1.6}$$

and the modified Bessel function of the first kind defined by

$$I_m(z) = \sum_{k=0}^{\infty} \frac{z^{2k+m}}{2^{2k+m} k! \Gamma(k+m+1)}, \tag{1.7}$$

where $(e)_k = e(e+1) \cdots (e+k-1)$ denotes the ascending factorial. The properties of these special functions being used can be found in Prudnikov et al. [30, 31, 32] and Gradshteyn and Ryzhik [7].

The paper is organized as follows. Sections 2 to 7 calculate expressions for R for the six distributions (Section 2 for McKay’s bivariate gamma distribution, Section 3 for Dus-sauchoy and Berland’s bivariate gamma distribution, Section 4 for Cherian’s bivariate gamma distribution, Section 5 for Arnold and Strauss’ bivariate gamma distribution, Section 6 for Becker and Roux’s bivariate gamma distribution, and Section 7 for Smith and Adelfang’s bivariate gamma distribution.) Conclusions and suggestions for future work are described in Section 8.

2. McKay’s bivariate gamma distribution

McKay’s [19] bivariate gamma distribution has the joint pdf given by

$$f(x, y) = \frac{a^{p+q}}{\Gamma(p)\Gamma(q)} x^{p-1} (y-x)^{q-1} \exp(-ay) \tag{2.1}$$

for $y > x > 0, a > 0, p > 0,$ and $q > 0$. For this distribution, it is easily seen that R is given by

$$\begin{aligned} R &= \frac{a^{p+q}}{\Gamma(p)\Gamma(q)} \int_0^{\infty} x^{p-1} \int_x^{\infty} (y-x)^{q-1} \exp(-ay) dy dx \\ &= \frac{a^{p+q}}{\Gamma(p)\Gamma(q)} \int_0^{\infty} x^{p-1} \Gamma(q) dx \\ &= a^{p+q}. \end{aligned} \tag{2.2}$$

3. Dussauchoy and Berland's bivariate gamma distribution

Dussauchoy and Berland's [6] bivariate gamma distribution has the joint pdf specified by

$$f(x, y) = \frac{\beta a_2^q}{\Gamma(p)\Gamma(q-p)} (\beta x)^{p-1} (y - \beta x)^{q-p-1} \exp(-a_2 x) \exp\left\{-\frac{a_2}{\beta}(y - \beta x)\right\} \times {}_1F_1\left(p, q-p; \left(\frac{a_1}{\beta} - a_2\right)(y - \beta x)\right) \tag{3.1}$$

for $x > 0, y > 0, 0 \leq \beta \leq 1, 0 < a_2 \leq a_1/\beta$, and $0 < p < q$. Using the definition of the confluent hypergeometric function given in Section 1, one can express the corresponding form of R as

$$\begin{aligned} R &= \frac{\beta a_2^q}{\Gamma(p)\Gamma(q-p)} \int_0^\infty (\beta x)^{p-1} \exp(-a_2 x) \\ &\quad \times \int_x^\infty (y - \beta x)^{q-p-1} \exp\left\{-\frac{a_2}{\beta}(y - \beta x)\right\} \\ &\quad \times \sum_{k=0}^\infty \frac{(p)_k}{k!(q-p)_k} \left(\frac{a_1}{\beta} - a_2\right)^k (y - \beta x)^k dy dx \\ &= \frac{\beta a_2^q}{\Gamma(p)\Gamma(q-p)} \int_0^\infty (\beta x)^{p-1} \exp(-a_2 x) \sum_{k=0}^\infty \frac{(p)_k}{k!(q-p)_k} \left(\frac{a_1}{\beta} - a_2\right)^k \\ &\quad \times \int_x^\infty (y - \beta x)^{k+q-p-1} \exp\left\{-\frac{a_2}{\beta}(y - \beta x)\right\} dy dx \\ &= \frac{\beta^q a_2^p}{\Gamma(p)\Gamma(q-p)} \int_0^\infty x^{p-1} \exp(-a_2 x) \sum_{k=0}^\infty \frac{(p)_k}{k!(q-p)_k} \left(\frac{a_1}{a_2} - \beta\right)^k \\ &\quad \times \int_{a_2(1-\beta)x/\beta}^\infty w^{k+q-p-1} \exp(-w) dw dx \\ &= \frac{\beta^q a_2^p}{\Gamma(p)\Gamma(q-p)} \int_0^\infty x^{p-1} \exp(-a_2 x) \sum_{k=0}^\infty \frac{(p)_k}{k!(q-p)_k} \left(\frac{a_1}{a_2} - \beta\right)^k \\ &\quad \times \Gamma\left(k+q-p, \frac{a_2(1-\beta)x}{\beta}\right) dx \\ &= \frac{\beta^q a_2^p}{\Gamma(p)\Gamma(q-p)} \sum_{k=0}^\infty \frac{(p)_k}{k!(q-p)_k} \left(\frac{a_1}{a_2} - \beta\right)^k I(k), \end{aligned} \tag{3.2}$$

where the transformation $w = a_2(y - \beta x)/\beta$ has been applied and $I(k)$ denotes the integral

$$I(k) = \int_0^\infty x^{p-1} \exp(-a_2 x) \Gamma\left(k+q-p, \frac{a_2(1-\beta)x}{\beta}\right) dx. \tag{3.3}$$

Application of Lemma A.1 in the appendix shows that $I(k)$ can be expressed in terms of the Gauss hypergeometric function as

$$I(k) = \frac{\beta^p(1-\beta)^{k+q}}{pa_2^p} \Gamma(k+q) {}_2F_1(1, k+q; p+1; \beta). \quad (3.4)$$

If $q-p$ is an integer, then by using the property

$$\Gamma(n, z) = (n-1)! \exp(-z) \sum_{m=0}^{n-1} \frac{z^k}{k!}, \quad (3.5)$$

one can obtain the elementary expression

$$\begin{aligned} I(k) &= \int_0^\infty x^{p-1} \exp(-a_2x) (k+q-p-1)! \sum_{m=0}^{k+q-p-1} \frac{1}{m!} \left\{ \frac{a_2(1-\beta)x}{\beta} \right\}^m dx \\ &= (k+q-p-1)! \sum_{m=0}^{k+q-p-1} \frac{1}{m!} \left\{ \frac{a_2(1-\beta)}{\beta} \right\}^m \int_0^\infty x^{m+p-1} \exp(-a_2x) dx \\ &= (k+q-p-1)! \left(\frac{a_2}{\beta} \right)^p \sum_{m=0}^{k+q-p-1} \frac{\Gamma(p+m)(1-\beta)^m}{m!}. \end{aligned} \quad (3.6)$$

Expressions for R can be obtained by substituting (3.4) and (3.6) into (3.2).

4. Cherian's bivariate gamma distribution

Cherian's [4] bivariate gamma distribution has the joint pdf specified by

$$f(x, y) = K \exp(-x) \exp(-y) \int_0^{\min(x,y)} (x-z)^{\theta_1-1} (y-z)^{\theta_2-1} z^{\theta_3-1} \exp(z) dz \quad (4.1)$$

for $x > 0$, $y > 0$, $\theta_1 > 0$, $\theta_2 > 0$, and $\theta_3 > 0$, where K denotes the normalizing constant given by

$$\frac{1}{K} = \Gamma(\theta_1) \Gamma(\theta_2) \Gamma(\theta_3). \quad (4.2)$$

For this distribution, the form of R can be expressed as

$$\begin{aligned} R &= K \int_0^\infty \int_x^\infty \int_0^{\min(x,y)} (x-z)^{\theta_1-1} (y-z)^{\theta_2-1} z^{\theta_3-1} \exp(z-x-y) dz dy dx \\ &= K \int_0^\infty \int_0^x (x-z)^{\theta_1-1} \exp(-x) z^{\theta_3-1} \exp(z) \left\{ \int_x^\infty (y-z)^{\theta_2-1} \exp(-y) dy \right\} dz dx \\ &= K \int_0^\infty \int_0^x (x-z)^{\theta_1-1} \exp(-x) z^{\theta_3-1} \exp(z) \left\{ \int_{x-z}^\infty w^{\theta_2-1} \exp(-w) dw \right\} dz dx \end{aligned}$$

$$\begin{aligned}
 &= K \int_0^\infty \int_0^x (x-z)^{\theta_1-1} \exp(-x) z^{\theta_3-1} \Gamma(\theta_2, x-z) dz dx \\
 &= K \int_0^\infty \int_0^x (x-z)^{\theta_1-1} \exp(-x) z^{\theta_3-1} \Gamma(\theta_2, x-z) dz dx \\
 &= K \int_0^\infty x^{\theta_1+\theta_3-1} \exp(-x) \int_0^1 t^{\theta_1-1} (1-t)^{\theta_3-1} \Gamma(\theta_2, xt) dt dx,
 \end{aligned} \tag{4.3}$$

where the transformations $w = y - z$ and $t = (x - z)/x$ have been applied. Application of Lemma A.2 shows that the inner integral in (4.3) can be calculated as

$$\begin{aligned}
 &\int_0^1 t^{\theta_1-1} (1-t)^{\theta_3-1} \Gamma(\theta_2, xt) dt \\
 &= \Gamma(\theta_2) B(\theta_1, \theta_3) + \theta_2^{-1} B(\theta_3, \theta_1 + \theta_2) x^{\theta_2} {}_2F_2(\theta_2, \theta_1 + \theta_2; \theta_2 + 1; \theta_1 + \theta_2 + \theta_3; -x),
 \end{aligned} \tag{4.4}$$

and thus R can be rewritten as

$$R = \frac{1}{\Gamma(\theta_1)\Gamma(\theta_3)} + \theta_2^{-1} B(\theta_3, \theta_1 + \theta_2) I, \tag{4.5}$$

where I denotes the integral

$$I = \int_0^\infty x^{\theta_1+\theta_2+\theta_3-1} \exp(-x) {}_2F_2(\theta_2, \theta_1 + \theta_2; \theta_2 + 1; \theta_1 + \theta_2 + \theta_3; -x) dx. \tag{4.6}$$

This integral can be calculated by an application of Lemma A.3 to yield

$$I = \Gamma(\theta_1 + \theta_2 + \theta_3) {}_2F_1(\theta_2, \theta_1 + \theta_2; \theta_2 + 1; -1), \tag{4.7}$$

and hence it follows from (4.5) that the form of R for Cherian's bivariate gamma distribution is given by

$$R = 1 + \theta_2^{-1} \Gamma(\theta_1 + \theta_2) \Gamma(\theta_3) {}_2F_1(\theta_2, \theta_1 + \theta_2; \theta_2 + 1; -1). \tag{4.8}$$

5. Arnold and Strauss' bivariate gamma distribution

Arnold and Strauss's [1] bivariate gamma distribution has the joint pdf specified by

$$f(x, y) = K x^{\alpha-1} y^{\beta-1} \exp\{- (ax + by + cxy)\} \tag{5.1}$$

for $x > 0, y > 0, \alpha > 0, \beta > 0, a > 0, b > 0,$ and $c > 0,$ where $K = K(a, b, c, \alpha, \beta)$ denotes the normalizing constant given by

$$\frac{1}{K} = b^{\alpha-\beta} c^{-\alpha} \Gamma(\alpha) \Gamma(\beta) \Psi\left(\alpha, \alpha - \beta + 1, \frac{ab}{c}\right). \tag{5.2}$$

For this distribution, the form of R can be derived as follows:

$$\begin{aligned}
 R &= K \int_0^\infty x^{\alpha-1} \exp(-ax) \int_x^\infty y^{\beta-1} \exp\{-(b+cx)y\} dy dx \\
 &= K \int_0^\infty \frac{x^{\alpha-1} \exp(-ax)}{(b+cx)^\beta} \int_{(b+cx)x}^\infty z^{\beta-1} \exp(-z) dz dx \\
 &= K \int_0^\infty \frac{x^{\alpha-1} \exp(-ax) \Gamma(\beta, (b+cx)x)}{(b+cx)^\beta} dx,
 \end{aligned} \tag{5.3}$$

where the transformation $z = (b+cx)y$ has been applied. The integral in (5.3) cannot be simplified further into its general form. However, if β is an integer, then using (3.5) one can rewrite (5.3) as

$$\begin{aligned}
 R &= K(\beta-1)! \int_0^\infty \frac{x^{\alpha-1} \exp(-ax)}{(b+cx)^\beta} \exp\{-(b+cx)x\} \sum_{l=0}^{\beta-1} \frac{(b+cx)^l x^l}{l!} dx \\
 &= K(\beta-1)! \sum_{l=0}^{\beta-1} \frac{I(l)}{l!},
 \end{aligned} \tag{5.4}$$

where $I(l)$ denotes the integral

$$I(l) = \int_0^\infty (b+cx)^{l-\beta} x^{l+\alpha-1} \exp\{-(ax+bx+cx^2)\} dx. \tag{5.5}$$

Again, this integral cannot be simplified further unless α is also an integer. If this is assumed then—by transforming $y = b+cx$ —one can express

$$\begin{aligned}
 I(l) &= c^{-(l+\alpha)} \exp\left\{\frac{(a+b)^2}{4c}\right\} \int_b^\infty y^{l-\beta} (y-b)^{l+\alpha-1} \exp\left\{-\frac{1}{c}\left(y+\frac{a-b}{2}\right)^2\right\} dy \\
 &= c^{-(l+\alpha)} \exp\left\{\frac{(a+b)^2}{4c}\right\} \int_b^\infty y^{l-\beta} \left[\sum_{m=0}^{l+\alpha-1} \binom{l+\alpha-1}{m} (-b)^{l+\alpha-1-m} y^m \right] \\
 &\quad \times \exp\left\{-\frac{1}{c}\left(y+\frac{a-b}{2}\right)^2\right\} dy \\
 &= c^{-(l+\alpha)} \exp\left\{\frac{(a+b)^2}{4c}\right\} \sum_{m=0}^{l+\alpha-1} \binom{l+\alpha-1}{m} (-b)^{l+\alpha-1-m} J(m, l),
 \end{aligned} \tag{5.6}$$

where

$$J(m, l) = \int_b^\infty y^{m-\beta+l} \exp\left\{-\frac{1}{c}\left(y+\frac{a-b}{2}\right)^2\right\} dy. \tag{5.7}$$

The integral $J(m, l)$ can be easily calculated. Three cases need to be considered.

(1) If $m = \beta - l$, then

$$J(m, l) = \sqrt{\pi c} \left\{ 1 - \Phi \left(\frac{a+b}{\sqrt{2c}} \right) \right\}, \tag{5.8}$$

where $\Phi(\cdot)$ denotes the cdf of the standard normal distribution.

(2) If $m > \beta - l$, then—by transforming $z = \{y + (a - b)/2\}^2$ —one can write

$$\begin{aligned} J(m, l) &= \frac{1}{2} \int_{(a+b)^2/4}^{\infty} \frac{1}{\sqrt{z}} \left(\sqrt{z} + \frac{b-a}{2} \right)^{m-\beta+l} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \int_{(a+b)^2/4}^{\infty} \frac{1}{\sqrt{z}} \left\{ \sum_{n=0}^{m-\beta+l} \binom{m-\beta+l}{n} \left(\frac{b-a}{2} \right)^{m-\beta+l-n} z^{n/2} \right\} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \sum_{n=0}^{m-\beta+l} \binom{m-\beta+l}{n} \left(\frac{b-a}{2} \right)^{m-\beta+l-n} \int_{(a+b)^2/4}^{\infty} z^{(n-1)/2} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \sum_{n=0}^{m-\beta+l} \binom{m-\beta+l}{n} \left(\frac{b-a}{2} \right)^{m-\beta+l-n} c^{(n+1)/2} \Gamma \left(\frac{n+1}{2}, \frac{(a+b)^2}{4c} \right), \end{aligned} \tag{5.9}$$

a finite sum of incomplete gamma functions.

(3) If $m < \beta - l$, then the same transformation as above yields

$$\begin{aligned} J(m, l) &= \frac{1}{2} \int_{(a+b)^2/4}^{\infty} z^{(m-\beta+l-1)/2} \left(1 + \frac{b-a}{2\sqrt{z}} \right)^{m-\beta+l} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \int_{(a+b)^2/4}^{\infty} z^{(m-\beta+l-1)/2} \left\{ \sum_{n=0}^{\infty} \binom{m-\beta+l}{n} \left(\frac{b-a}{2\sqrt{z}} \right)^n \right\} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \binom{m-\beta+l}{n} \left(\frac{b-a}{2} \right)^n \int_{(a+b)^2/4}^{\infty} z^{(m-\beta+l-n-1)/2} \exp \left(-\frac{z}{c} \right) dz \\ &= \frac{1}{2} \sum_{n=0}^{\infty} \binom{m-\beta+l}{n} \left(\frac{b-a}{2} \right)^n c^{(m-\beta+l-n+1)/2} \times \Gamma \left(\frac{m-\beta+l-n+1}{2}, \frac{(a+b)^2}{4c} \right), \end{aligned} \tag{5.10}$$

an infinite sum of incomplete gamma functions.

Combining (5.4)–(5.10), the reliability R can be expressed as an infinite sum of incomplete gamma functions for integer values of α and β and general a, b , and c . However, if $a = b$, then one can express R as a finite sum of incomplete gamma functions. This follows because if $a = b$, then $J(m, l)$ reduces to

$$\begin{aligned} J(m, l) &= 2^{-1} c^{1+(m-\beta)/2} \int_{a^2/c}^{\infty} z^{(m-\beta)/2} \exp(-z) dz \\ &= 2^{-1} c^{1+(m-\beta)/2} \Gamma \left(1 + \frac{m-\beta}{2}, \frac{a^2}{c} \right) \end{aligned} \tag{5.11}$$

(after substituting $z = y^2/c$), and combining (5.11) with (5.4)–(5.6), one obtains

$$R = \frac{K(\beta - 1)! \exp(a^2/c)}{2(-a)^{1-\alpha} c^{\alpha+\beta/2-1}} \sum_{l=0}^{\beta-1} \sum_{m=0}^{l+\alpha-1} \frac{(-a)^{l-m}}{l!c^{l-m/2}} \binom{l+\alpha-1}{m} \Gamma\left(1 + \frac{m-\beta}{2}, \frac{a^2}{c}\right), \tag{5.12}$$

a finite sum of incomplete gamma functions. If also $\alpha = \beta$, then it is easy to see that $R = 1/2$.

6. Becker and Roux’s bivariate gamma distribution

Becker and Roux’s [3] bivariate gamma distribution has the joint pdf given by

$$f(x, y) = \begin{cases} \frac{\lambda_2}{\Gamma(h)\Gamma(l)\alpha_1^h\alpha_2^l} x^{h-1} \{\lambda_2(y-x) + x\}^{l-1} \exp\left\{-\left(\frac{1}{\alpha_1} + \frac{1-\lambda_2}{\alpha_2}\right)x - \frac{\lambda_2}{\alpha_2}y\right\} & \text{if } x < y, \\ \frac{\lambda_1}{\Gamma(h)\Gamma(l)\alpha_1^h\alpha_2^l} y^{l-1} \{\lambda_1(x-y) + y\}^{h-1} \exp\left\{-\left(\frac{1}{\alpha_1} + \frac{1-\lambda_1}{\alpha_1}\right)y - \frac{\lambda_1}{\alpha_1}x\right\} & \text{if } x > y, \end{cases} \tag{6.1}$$

for $x > 0, y > 0, h > 0, l > 0, \lambda_1 > 0, \lambda_2 > 0, \alpha_1 > 0$, and $\alpha_2 > 0$. For this distribution, the form of R can be expressed as

$$\begin{aligned} R &= \frac{\lambda_2}{\Gamma(h)\Gamma(l)\alpha_1^h\alpha_2^l} \int_0^\infty x^{h-1} \exp\left\{-\left(\frac{1}{\alpha_1} + \frac{1-\lambda_2}{\alpha_2}\right)x\right\} \\ &\quad \times \int_x^\infty \{\lambda_2(y-x) + x\}^{l-1} \exp\left\{-\frac{\lambda_2}{\alpha_2}y\right\} dy dx \\ &= \frac{1}{\Gamma(h)\Gamma(l)\alpha_1^h\alpha_2^l} \int_0^\infty x^{h-1} \exp\left(-\frac{x}{\alpha_1}\right) \int_x^\infty z^{l-1} \exp\left(-\frac{z}{\alpha_2}\right) dz dx \\ &= \frac{1}{\Gamma(h)\Gamma(l)\alpha_1^h} \int_0^\infty x^{h-1} \exp\left(-\frac{x}{\alpha_1}\right) \Gamma\left(l, \frac{x}{\alpha_2}\right) dx, \end{aligned} \tag{6.2}$$

where the transformation $z = \lambda_2(y-x) + x$ has been applied. By application of Lemma A.1 in the appendix, the integral in (6.2) can be calculated as

$$\int_0^\infty x^{h-1} \exp\left(-\frac{x}{\alpha_1}\right) \Gamma\left(l, \frac{x}{\alpha_2}\right) dx = \frac{\alpha_1^{h+l}\alpha_2^h}{h(\alpha_1 + \alpha_2)^{h+l}} {}_2F_1\left(1, h+l; h+1; \frac{\alpha_2}{\alpha_1 + \alpha_2}\right). \tag{6.3}$$

Thus, the form of R can be expressed in terms of the Gauss hypergeometric function as

$$R = \frac{\alpha_1^l\alpha_2^h}{hB(h, l)(\alpha_1 + \alpha_2)^{h+l}} {}_2F_1\left(1, h+l; h+1; \frac{\alpha_2}{\alpha_1 + \alpha_2}\right). \tag{6.4}$$

If l is an integer, then the hypergeometric term reduces to a finite sum, yielding

$$R = \frac{\alpha_2^h}{hB(h, l)(\alpha_1 + \alpha_2)^h} \sum_{k=0}^{l-1} \frac{(h)_k(1-l)_k}{(h+1)_kk!} \left(\frac{\alpha_2}{\alpha_1 + \alpha_2}\right)^k. \tag{6.5}$$

7. Smith and Adelfang’s bivariate gamma distribution

Smith and Adelfang’s [34] bivariate gamma distribution has the joint pdf specified by

$$f(x, y) = Kx^{\gamma_1-1}y^{\gamma_2-1} \exp\left(-\frac{x+y}{1-\eta}\right) \sum_{k=0}^{\infty} a_k I_{\gamma_2+k-1}\left(\frac{2\sqrt{\eta xy}}{1-\eta}\right) \tag{7.1}$$

for $x > 0, y > 0, \gamma_1 > 0, \gamma_2 > 0$, and $0 < \eta < 1$, where K denotes the normalizing constant given by

$$\frac{1}{K} = (1-\eta)^{\gamma_1} \Gamma(\gamma_1) \Gamma(\gamma_2 - \gamma_1). \tag{7.2}$$

Using the definition of $I_m(\cdot)$ given in Section 1, the corresponding form of R can be expressed as

$$\begin{aligned} R &= K \sum_{k=0}^{\infty} a_k \sum_{l=0}^{\infty} \frac{1}{l! \Gamma(\gamma_2 + l + k)} \left(\frac{\sqrt{\eta}}{1-\eta}\right)^{k+2l+\gamma_2-1} \\ &\quad \times \int_0^{\infty} x^{\gamma_1+l+(\gamma_2+k)/2-1} \int_x^{\infty} y^{\gamma_2+l+(\gamma_2+k-1)/2-1} \exp\left(-\frac{x+y}{1-\eta}\right) dy dx \tag{7.3} \\ &= K \sum_{k=0}^{\infty} a_k \sum_{l=0}^{\infty} \frac{\eta^{(k+2l+\gamma_2-1)/2} (1-\eta)^{(\gamma_2-k-2l+1)/2}}{l! \Gamma(\gamma_2 + l + k)} J(k, l), \end{aligned}$$

where $J(k, l)$ denotes the integral

$$J(k, l) = \int_0^{\infty} x^{\gamma_1+(k+2l+\gamma_2-1)/2-1} \exp\left(-\frac{x}{1-\eta}\right) \Gamma\left(\frac{k+2l+3\gamma_2-1}{2}, \frac{x}{1-\eta}\right) dx. \tag{7.4}$$

Application of Lemma A.1 in the appendix shows that $J(k, l)$ can be expressed in terms of the Gauss hypergeometric function as

$$\begin{aligned} J(k, l) &= \frac{(1-\eta)^{\gamma_1+l+(\gamma_2+k-1)/2} \Gamma(\gamma_1 + 2\gamma_2 + k + 2l - 1)}{2^{\gamma_1+2\gamma_2+k+2l-2} (2\gamma_1 + \gamma_2 + k + 2l - 1)} \\ &\quad \times {}_2F_1\left(1, \gamma_1 + 2\gamma_2 + k + 2l - 1; \gamma_1 + l + \frac{\gamma_2 + k + 1}{2}; \frac{1}{2}\right). \tag{7.5} \end{aligned}$$

Substituting this into (7.3), one obtains the expression

$$\begin{aligned} R &= K (1-\eta)^{\gamma_1+\gamma_2} 2^{2-\gamma_1-2\gamma_2} \\ &\quad \times \sum_{k=0}^{\infty} a_k \sum_{l=0}^{\infty} \frac{\eta^{(k+2l+\gamma_2-1)/2} \Gamma(\gamma_1 + 2\gamma_2 + k + 2l - 1)}{l! 2^{k+2l} (2\gamma_1 + \gamma_2 + k + 2l - 1) \Gamma(\gamma_2 + l + k)} \\ &\quad \times {}_2F_1\left(1, \gamma_1 + 2\gamma_2 + k + 2l - 1; \gamma_1 + l + \frac{\gamma_2 + k + 1}{2}; \frac{1}{2}\right), \tag{7.6} \end{aligned}$$

a double infinite sum of terms involving the Gauss hypergeometric function.

8. Conclusions

We have calculated the forms of $R = \Pr(X < Y)$ for six flexible families of bivariate gamma distributions. It would be of interest to emulate this work for other continuous bivariate distributions, including bivariate Laplace distributions, bivariate logistic distributions, and bivariate extreme value distributions. It would also be of interest to extend this work for continuous multivariate distributions. We hope to address some of these issues in a future paper.

Appendix

Some technical lemmas required for the calculations above are noted below.

LEMMA A.1 (Gradshteyn and Ryzhik [7, equation (6.455.1)]). For $c > 0$, $p > 0$, and $\alpha + \nu > 0$,

$$\int_0^{\infty} x^{\alpha-1} \exp(-px) \Gamma(\nu, cx) dx = -\frac{c^{\nu} \Gamma(\alpha + \nu)}{\nu p^{\alpha+\nu}} {}_2F_1\left(\nu, \alpha + \nu; \nu + 1; -\frac{c}{p}\right) + \frac{\Gamma(\nu) \Gamma(\alpha)}{p^{\alpha}}. \quad (\text{A.1})$$

LEMMA A.2 (Prudnikov et al. [31, Volume 2, equation (2.10.2.2)]). For $\alpha > 0$ and $\beta > 0$,

$$\begin{aligned} \int_0^a x^{\alpha-1} (a-x)^{\beta-1} \Gamma(\nu, cx) dx \\ = a^{\alpha+\beta-1} \Gamma(\nu) B(\alpha, \beta) - \nu^{-1} a^{\alpha+\beta+\nu-1} c^{\nu} B(\beta, \alpha + \nu) {}_2F_2(\nu, \alpha + \nu; \nu + 1, \alpha + \beta + \nu; -ac). \end{aligned} \quad (\text{A.2})$$

LEMMA A.3 (Prudnikov et al. [32, Volume 3, equation (2.22.3.5)]). For $c > 0$ and $\sigma > w$,

$$\int_0^{\infty} x^{c-1} \exp(-\sigma x) {}_2F_2(a, b; d, c; wx) dx = \sigma^{-c} \Gamma(c) {}_2F_1\left(a, b; d; \frac{w}{\sigma}\right). \quad (\text{A.3})$$

Acknowledgments

The author is grateful to Professor Samuel Kotz (George Washington University, USA) and to Professor Marianna Pensky (University of Central Florida, USA) for introducing him to this area of research. The author would also like to thank the referee and the editors for carefully reading the paper and for their great help in improving the paper.

References

- [1] B. C. Arnold and D. Strauss, *Bivariate distributions with exponential conditionals*, J. Amer. Statist. Assoc. **83** (1988), no. 402, 522–527.
- [2] A. M. Awad, M. A. Hamdan, and M. M. Azzam, *Some inference results on $\Pr(X < Y)$ in the bivariate exponential model*, Comm. Statist. Theory Methods **10** (1981), no. 24, 2515–2525.
- [3] P. J. Becker and J. J. J. Roux, *A bivariate extension of the gamma distribution*, South African Statist. J. **15** (1981), no. 1, 1–12.
- [4] K. C. Cherian, *A bi-variate correlated gamma-type distribution function*, J. Indian Math. Soc. (N.S.) **5** (1941), 133–144.
- [5] E. Cramer and U. Kamps, *The UMVUE of $P(X < Y)$ based on type-II censored samples from Weinman multivariate exponential distributions*, Metrika **46** (1997), no. 2, 93–121.

- [6] A. Dussauchoy and R. Berland, *Lois gamma à deux dimensions*, C. R. Acad. Sci. Paris Sér. A-B **274** (1972), A1946–A1949 (French).
- [7] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*, Academic Press, California, 2000.
- [8] R. C. Gupta and S. Subramanian, *Estimation of reliability in a bivariate normal distribution with equal coefficients of variation*, Comm. Statist. Simulation Comput. **27** (1998), no. 3, 675–698.
- [9] R. D. Gupta and R. C. Gupta, *Estimation of $\Pr(a'x > b'y)$ in the multivariate normal case*, Statistics **21** (1990), no. 1, 91–97.
- [10] D. D. Hanagal, *Testing reliability in a bivariate exponential stress-strength model*, J. Indian Statist. Assoc. **33** (1995), 41–45.
- [11] ———, *Note on estimation of reliability under bivariate Pareto stress-strength model*, Statist. Papers **38** (1997), no. 4, 453–459.
- [12] ———, *Estimation of reliability of a component subjected to bivariate exponential stress*, Statist. Papers **40** (1999), no. 2, 211–220.
- [13] P. K. Jana, *Estimation for $P[Y < X]$ in the bivariate exponential case due to Marshall and Olkin*, J. Indian Statist. Assoc. **32** (1994), no. 1, 35–37.
- [14] ———, *Estimation of $P[Y < X]$ under a bivariate exponential stress-strength model*, Frontiers in Probability and Statistics (Calcutta, 1994/1995) (S. P. Mukherjee, S. K. Basu, and Sinha B. K., eds.), Narosa, New Delhi, 1998, pp. 207–214.
- [15] P. K. Jana and D. Roy, *Estimation of reliability under stress-strength model in a bivariate exponential set-up*, Calcutta Statist. Assoc. Bull. **44** (1994), no. 175–176, 175–181.
- [16] E. S. Jeevanand, *Bayes estimation of $P(X_2 < X_1)$ for a bivariate Pareto distribution*, The Statistician **46** (1997), no. 1, 93–99.
- [17] ———, *Bayes estimation of reliability under stress-strength model for the Marshall–Olkin bivariate exponential distribution*, IAPQR Trans. **23** (1998), no. 2, 133–136.
- [18] J. P. Klein and A. P. Basu, *Estimating reliability for bivariate exponential distributions*, Sankhyā Ser. B **47** (1985), no. 3, 346–353.
- [19] A. T. McKay, *Sampling from batches*, Journal of the Royal Statistical Society—Supplement **1** (1934), 207–216.
- [20] S. P. Mukherjee and L. K. Saran, *Estimation of failure probability from a bivariate normal stress-strength distribution*, Microelectronics Reliability **25** (1985), 692–702.
- [21] S. Nadarajah, *Reliability for beta models*, Serdica Math. J. **28** (2002), no. 3, 267–282.
- [22] ———, *Reliability for extreme value distributions*, Math. Comput. Modelling **37** (2003), no. 9–10, 915–922.
- [23] ———, *Reliability for lifetime distributions*, Math. Comput. Modelling **37** (2003), no. 7–8, 683–688.
- [24] ———, *Reliability for logistic distributions*, Engineering Simulation **26** (2004), 81–98.
- [25] ———, *Reliability for Laplace*, Math. Probl. Eng. **2004** (2004), no. 2, 169–183.
- [26] ———, *Reliability for some bivariate beta distributions*, Math. Probl. Eng. **2005** (2005), no. 1, 101–111.
- [27] S. Nadarajah and S. Kotz, *Reliability for Pareto models*, Metron **61** (2003), no. 2, 191–204.
- [28] S. B. Nandi and A. B. Aich, *A note on confidence bounds for $P(X > Y)$ in bivariate normal samples*, Sankhyā Ser. B **56** (1994), no. 2, 129–136.
- [29] ———, *A note on testing hypothesis regarding $P(X > Y)$ in bivariate normal samples*, IAPQR Trans. **21** (1996), no. 2, 149–153.
- [30] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Integrals and Series. Vol. 1*, Gordon & Breach Science Publishers, New York, 1986.
- [31] ———, *Integrals and Series. Vol. 2*, Gordon & Breach Science Publishers, New York, 1986.
- [32] ———, *Integrals and Series. Vol. 3*, Gordon & Breach Science Publishers, New York, 1986.

- [33] N. Singh, *MVUE of $\Pr(X < Y)$ for multivariate normal populations: an application to stress-strength models*, IEEE Trans. Rel. **30** (1981), 192–193.
- [34] O. E. Smith and S. I. Adelfang, *Gust model based on the bivariate gamma probability distribution*, Journal of Spacecraft and Rockets **18** (1981), 545–549.

Saralees Nadarajah: Department of Statistics, University of Nebraska, Lincoln, NE 68583, USA
E-mail address: snadaraj@unlserve.unl.edu

Special Issue on Decision Support for Intermodal Transport

Call for Papers

Intermodal transport refers to the movement of goods in a single loading unit which uses successive various modes of transport (road, rail, water) without handling the goods during mode transfers. Intermodal transport has become an important policy issue, mainly because it is considered to be one of the means to lower the congestion caused by single-mode road transport and to be more environmentally friendly than the single-mode road transport. Both considerations have been followed by an increase in attention toward intermodal freight transportation research.

Various intermodal freight transport decision problems are in demand of mathematical models of supporting them. As the intermodal transport system is more complex than a single-mode system, this fact offers interesting and challenging opportunities to modelers in applied mathematics. This special issue aims to fill in some gaps in the research agenda of decision-making in intermodal transport.

The mathematical models may be of the optimization type or of the evaluation type to gain an insight in intermodal operations. The mathematical models aim to support decisions on the strategic, tactical, and operational levels. The decision-makers belong to the various players in the intermodal transport world, namely, drayage operators, terminal operators, network operators, or intermodal operators.

Topics of relevance to this type of decision-making both in time horizon as in terms of operators are:

- Intermodal terminal design
- Infrastructure network configuration
- Location of terminals
- Cooperation between drayage companies
- Allocation of shippers/receivers to a terminal
- Pricing strategies
- Capacity levels of equipment and labour
- Operational routines and lay-out structure
- Redistribution of load units, railcars, barges, and so forth
- Scheduling of trips or jobs
- Allocation of capacity to jobs
- Loading orders
- Selection of routing and service

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/jamds/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/>, according to the following timetable:

Manuscript Due	June 1, 2009
First Round of Reviews	September 1, 2009
Publication Date	December 1, 2009

Lead Guest Editor

Gerrit K. Janssens, Transportation Research Institute (IMOB), Hasselt University, Agoralaan, Building D, 3590 Diepenbeek (Hasselt), Belgium; Gerrit.Janssens@uhasselt.be

Guest Editor

Cathy Macharis, Department of Mathematics, Operational Research, Statistics and Information for Systems (MOSI), Transport and Logistics Research Group, Management School, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium; Cathy.Macharis@vub.ac.be