

Research Article

Variational Problems with Moving Boundaries Using Decomposition Method

Reza Memarbashi

Received 12 December 2006; Revised 23 April 2007; Accepted 25 July 2007

Recommended by T. Zolezzi

The aim of this paper is to present a numerical method for solving variational problems with moving boundaries. We apply Adomian decomposition method on the Euler-Lagrange equation with boundary conditions that yield from transversality conditions and natural boundary conditions.

Copyright © 2007 Reza Memarbashi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In many problems arising in mathematics, physics, engineering, economics, and other sciences, it is necessary to minimize amounts of a certain functional. Because of the important role of this subject, considerable attention has been devoted to these kinds of problems. Such problems are called variational problems, see [1, 2].

The simplest form of a variational problem can be considered as

$$J[y] = \int_a^b f(x, y(x), y'(x)) dx, \quad (1.1)$$

where J is called a functional, and its minimum must be found. Functional J can be considered by two kinds of boundary conditions. The first kind, we refer to as fixed boundary problems, the admissible function $y(x)$ must satisfy the boundary conditions

$$y(a) = y_a, \quad y(b) = y_b. \quad (1.2)$$

The second kind, we call problems with moving boundaries and concern ourselves with them, is variational problems for which at least one of the boundary points of the admissible function is movable along a boundary curve.

2 Mathematical Problems in Engineering

The variational problem where the beginning point and the endpoint are fixed is often referred to as point-point problem, and the problems with variable boundaries as point-curve, curve-point, and curve-curve problems.

A necessary condition for the admissible solutions of such problems is to satisfy the Euler-Lagrange equation

$$\frac{\partial f}{\partial y} - \frac{d}{dx} \frac{\partial f}{\partial y'} = 0. \quad (1.3)$$

For the point-point problem, Euler-Lagrange equation must be considered by the boundary conditions (1.2). For the problems with variable boundaries, we have two cases.

Type 1: As the first case, we consider problems for which at least one of the boundary points move freely along a line parallel to the y -axis, in fact at this point $y(x)$ is not specified. In this case, admissible functions must fulfill the natural boundary conditions (2.3) and (2.4), (or only one of them).

Type 2: For the second case, we will let the beginning and endpoints (or only one of them) move freely on given curves $y = \psi(x)$, $y = \varphi(x)$. In this case, the admissible function $y(x)$ and points a and b (or one of them) must satisfy the necessary conditions (2.5) and (2.6) called transversality conditions.

The decomposition method proposed by G. Adomian is a power tool to investigate various models in applied science, engineering, and so forth, (see [3, 4]). The method provides the solution in a rapidly convergent series with components that can be computed recursively, see [5, 6] for convergence. Adomian decomposition is applied on various kinds of problems, for example, see [7–24] for some recent works. It produces an efficient solution with high accuracy and minimal calculation for finding the approximate solution of the linear and nonlinear problems. Dehghan and Tatari in [9] used Adomian decomposition to solve variational problems with fixed boundaries. In this work we solve variational problems with moving boundaries by this method. In fact, we apply Adomian decomposition method to solve the Euler-Lagrange equation considered by the boundary conditions which yield from natural boundary conditions and transversality conditions. In the last section, we use some examples to show the numerical treatment of the proposed method.

2. Statement of the problem

2.1. Boundary conditions. The first type of problem here is one for which at least at one of the endpoints, $y(x)$ is not specified. In this case, all admissible functions have the same domain of definition $[a, b]$ and satisfy the Euler-Lagrange equation in this interval. Furthermore, such function must satisfy conditions called natural boundary conditions prescribed in the following theorem.

THEOREM 2.1. *Suppose the function $y = y_0(x) \in C^1[a, b]$ yield a relative minimum of the functional (1.1) for which*

$$y(a) = y_a \text{ is given, } \quad y(b) \text{ is arbitrary (free right endpoint)} \quad (2.1)$$

or

$$y(a), y(b) \text{ are arbitrary (free endpoints).} \quad (2.2)$$

Then $y_0(x)$ satisfies, respectively, the following natural boundary conditions:

$$\frac{\partial f}{\partial y'}(b, y_0(b), y'_0(b)) = 0 \quad (\text{free right endpoint}) \quad (2.3)$$

or

$$\frac{\partial f}{\partial y'}(a, y_0(a), y'_0(a)) = 0 = \frac{\partial f}{\partial y'}(b, y_0(b), y'_0(b)) \quad (\text{free endpoints}). \quad (2.4)$$

We now consider a generalized form of the problem. We let endpoints move freely on given curves.

In this case, we seek for a function $y(x)$, which emanates at some $x = a$ from the curve $y = \psi(x)$ and terminates for some $x = b$ on the curve $y = \varphi(x)$ and minimizes the functional (1.1). In this problem also the points a, b are not known, they must satisfy the necessary conditions called transversality conditions, prescribed in the following theorem.

THEOREM 2.2. *If the function $y = y_0(x) \in C^1[a, b]$, which emanates at some $x = a$ from the curve $y = \psi(x) \in C^1(-\infty, \infty)$ and terminates for some $x = b$ on the curve $y = \varphi(x) \in C^1(-\infty, \infty)$, yields a relative minimum for functional (1.1), where $f \in C^1(R)$, R being a domain in the (x, y, y') space that contains all lineal elements of $y = y_0(x)$, then it is necessary that $y = y_0(x)$ must satisfy the Euler-Lagrange equation in the interval $[a, b]$ and that at the point of departure and the point of arrival, the transversality conditions*

$$\frac{\partial f}{\partial y'}(a, y_0(a), y'_0(a))(\psi'(a) - y'_0(a)) + f(a, y_0(a), y'_0(a)) = 0, \quad (2.5)$$

$$\frac{\partial f}{\partial y'}(b, y_0(b), y'_0(b))(\varphi'(b) - y'_0(b)) + f(b, y_0(b), y'_0(b)) = 0 \quad (2.6)$$

are satisfied. In the case that one of the points is fixed, then the transversality condition has to hold at the other point.

We see that the natural boundary conditions (2.3) and (2.4) may be viewed as a special case of the transversality conditions for $\psi'(a) = \infty$ and $\varphi'(b) = \infty$.

If the initial curve or the terminal curve (or both) is a horizontal line, that is, if $\psi'(x) = 0$ or $\varphi'(x) = 0$ (or both), then we obtain from (2.5) that

$$\frac{\partial f}{\partial y'}(a, y_0(a), y'_0(a))y'_0(a) - f(a, y_0(a), y'_0(a)) = 0, \quad (2.7)$$

and an analogous condition for the right endpoint holds.

One can consider transversality conditions for the problems with more than one unknown function.

4 Mathematical Problems in Engineering

For example, in the two-dimensional case we seek a vector function $\hat{y} = \hat{y}(x) = (y_1(x), y_2(x))$ such that it minimizes

$$J[y_1, y_2] = \int_a^b f(x, y_1(x), y_2(x), y_1'(x), y_2'(x)) dx, \quad (2.8)$$

in which $y_1(a) = y_{1,a}$, $y_2(a) = y_{2,a}$ and the endpoint lies on a two-dimensional surface that is given by $x = u(y_1, y_2)$. Here, the necessary condition for the extremum of the functional (2.8) is to satisfy the following system of second-order differential equations:

$$\frac{\partial f}{\partial y_i} - \frac{d}{dx} \frac{\partial f}{\partial y_i'} = 0, \quad i = 1, 2; \quad (2.9)$$

and the terminal point $x = b$ must satisfy the following transversality conditions:

$$\begin{aligned} \left(\frac{\partial u}{\partial y_1} f + \left(1 - \frac{\partial u}{\partial y_1} y_1^{0'} - \frac{\partial u}{\partial y_2} y_2^{0'} \right) \frac{\partial f}{\partial y_1'} \right) \Big|_{x=b} &= 0, \\ \left(\frac{\partial u}{\partial y_2} f + \left(1 - \frac{\partial u}{\partial y_1} y_1^{0'} - \frac{\partial u}{\partial y_2} y_2^{0'} \right) \frac{\partial f}{\partial y_2'} \right) \Big|_{x=b} &= 0, \end{aligned} \quad (2.10)$$

in which $\hat{y} = \hat{y}_0(x) = (y_1^0(x), y_2^0(x))$ is an admissible vector function. For further information on transversality conditions, specially for the proofs of Theorems 2.1 and 2.2 and condition (2.10), see [2].

2.2. Solution of problem using Adomian decomposition method. The Euler-Lagrange (1.3) can be considered in an operator form

$$L(y) - N(y) = g \quad (2.11)$$

for $a \leq x \leq b$, where $L = d^2/dx^2$ is the second-order derivative operator, N is an operator (which may be nonlinear), and g is a given function. We consider the inverse operator in the following form:

$$L^{-1}(h(x)) = \int_a^x \int_a^{x_2} h(x_1) dx_1 dx_2. \quad (2.12)$$

Applying operator L^{-1} to both sides of (2.11), we have

$$y(x) - y(a) - y'(a)x + y'(a)a = L^{-1}N(y) + L^{-1}g. \quad (2.13)$$

So that,

$$y(x) = \alpha + Ax - Aa + L^{-1}g + L^{-1}N(y). \quad (2.14)$$

Now according to the decomposition procedure of Adomian, we construct the unknown function $y(x)$ by a sum of components defined by the following series:

$$y(x) = \sum_{n=0}^{\infty} y_n(x) \quad (2.15)$$

and consider the nonlinear expression $N(y)$ by the infinite series of the polynomials given by $N(y) = \sum_{n=0}^{\infty} N_n$ where components N_n have the following form:

$$N_n(y_0, y_1, \dots, y_n) = \frac{1}{n!} \frac{d^n}{d\lambda^n} \left[N \left[\sum_{k=0}^{\infty} \lambda^k y_k \right] \right]_{\lambda=0}, \quad n \geq 0. \quad (2.16)$$

Notice that if N be a linear operator then we have $N_n = y_n$. Now we have the following recursive relations:

$$\begin{aligned} y_0(x) &= \alpha + Ax + L^{-1}g(x), \\ y_{n+1}(x) &= L^{-1}N_n, \quad n \geq 0. \end{aligned} \quad (2.17)$$

Based on the Adomian decomposition method, we constructed the solution $y(x)$ as

$$y = \lim_{n \rightarrow \infty} \phi_n, \quad (2.18)$$

where the $n + 1$ term approximation of the solution is defined in the following form:

$$\phi_n = \sum_{k=0}^n y_k(x), \quad n > 0. \quad (2.19)$$

The solution here is given in a series form that converges rapidly in physical problems.

Applying the decomposition procedure of Adomian, we find the series solution of $y(x)$ follows with constants α, A which are unknown. For the determination of these constants, we have two cases.

Case (1): For the variational problems of type 1, we impose the boundary conditions that yield from natural boundary conditions (2.3) and (2.4) to the obtained approximation of the solution (2.19) which results equations in α, A . Remark that for the point-curve and curve-point problems of this type only constant A is undetermined, also we have one equation in A . However, by solving these equations we find α, A .

Case (2): Consider variational problems of type 2, for example, a point-curve problem. In such problem, the point a and constant α are known but the point b and the constant A are unknown. In this case, we have two equations, one equation yields from transversality condition at $x = b$ and another equation yields from the intersection of $y(x)$ and the terminal curve $y = \varphi(x)$ at $x = b$. By solving this system of equations, which is usually nonlinear, we find both b and A and the solution of the Euler-Lagrange equation follows immediately.

Note that for a curve-curve problem of this type, we have four unknown a, b, α, A . In this case, we have four equations. Two equations yield from transversality conditions at $x = a, b$ and two equations obtained from the intersection of $y(x)$ by $y = \psi(x)$ and $y = \varphi(x)$ at $x = a$ and $x = b$, respectively. Solution of this system of equations determines these unknown.

3. Numerical tests

To give an obvious overview of the Adomian decomposition method on the variational problems with moving boundaries, we present some examples.

3.1. Example 1 (Ramsey growth model). Consider the following variational problem:

$$J[y] = \int_0^T a(by(t) - y'(t) - c^*)^2 dt, \quad (3.1)$$

in which $a, b > 0$, $C^* > 0$ and $y(t)$ is the amount of a capital at time t (see [1] for further information on this economic model).

Here, the capital stock $y(0)$ at the initial time $t = 0$ of the planing period is assumed to be known: $y(0) = y_0$; on the other hand, the planner will not want to prescribe how large the capital will be at time $t = T$. We have therefore a variational problem with free right endpoint.

For this numerical example, we let $a = b = c^* = 1$, $T = 1$, and $y_0 = 2$ which has the analytical solution $y(t) = 1 + e^t$.

The corresponding Euler-Lagrange equation is

$$y''(t) - y(t) + 1 = 0. \quad (3.2)$$

Now we consider the natural boundary condition (2.3) at $t = 1$:

$$\frac{\partial f}{\partial y'}(1, y(1), y'(1)) = -2(y(t) - y'(t) - 1) \Big|_{t=1} = 0. \quad (3.3)$$

Therefore, we have the following boundary conditions for (3.2):

$$y(0) = 2, \quad y(1) - y'(1) - 1 = 0. \quad (3.4)$$

Using the operator form of (3.2), we have

$$\begin{aligned} Ly &= y - 1, \\ y(x) &= 2 + Ax - L^{-1}(1) + L^{-1}(y(x)). \end{aligned} \quad (3.5)$$

Applying the Adomian decomposition, we have

$$\sum_{n=0}^{\infty} y_n(x) = 2 + Ax - \frac{x^2}{2} + L^{-1} \left(\sum_{n=0}^{\infty} y_n(x) \right). \quad (3.6)$$

Therefore, we use the recursive relations

$$\begin{aligned} y_0(x) &= 2 + Ax - \frac{x^2}{2}, \\ y_{n+1}(x) &= L^{-1}(y_n(x)), \quad n \geq 0. \end{aligned} \quad (3.7)$$




FIGURE 3.1. Plot of $y(x)$ and $\phi_3(x)$ in example 1.

Now we use $\phi_n(x)$ as the approximation of $y(x)$, for example, for $n = 3$ we have

$$\begin{aligned} \phi_3(x) = 2 + Ax + \frac{1}{2}x^2 + \frac{1}{6}Ax^3 + \frac{1}{24}x^4 + \frac{1}{120}Ax^5 \\ + \frac{1}{720}x^6 + \frac{1}{5040}Ax^7 - \frac{1}{40320}x^8. \end{aligned} \tag{3.8}$$

For the determination of A , we solve the equation $\phi_3(1) - \phi_3'(1) - 1 = 0$. Figure 3.1 shows $y(x)$ and $\phi_3(x)$ and Table 3.1 shows the error of ϕ_3 .

3.2. Example 2 (Fermat’s principle). We consider in this example the problem of a light ray emanating from a fixed point P_a , propagating in a plane through a medium with $n(x, y) > 0$ as an index of refraction, and terminating at some point $x = b$ on the curve $y = \varphi(x)$.

By Fermat’s principle, the light ray will take the path that minimizes the total traveling time from P_a to $x = b$ on $y = (x)$, that is,

$$J[y] = \int_a^b n(x, y(x))\sqrt{1 + y'^2(x)}dx. \tag{3.9}$$

Observe that the Brachistocherone problem, the minimal surfaces of revolution and the shortest path problem are special cases of Fermat’s principle for $n(x, y) = 1/\sqrt{y}$, $n(x, y) = y$, and $n(x, y) = 1$, respectively.

TABLE 3.1. Error ($e(x) = |y(x) - \phi_3(x)|$) in Example 1.

x	0	0.2	0.4	0.6	0.8	1
$e(x)$	0	.203617e-3	.415371e-3	.643003e-3	.889442e-3	.1135848e-2

The transversality condition in this case is of the form

$$\left(n(x, y)\sqrt{1 + y'^2} + \frac{n(x, y)y'}{\sqrt{1 + y'^2}}(\phi' - y') \right) \Big|_{x=b} = 0 \tag{3.10}$$

or

$$\left(\frac{n(x, y)(1 + \phi' y')}{\sqrt{1 + y'^2}} \right) \Big|_{x=b} = 0. \tag{3.11}$$

Finally, we have $y'(b) = -1/\phi'(b)$, that is, the transversality condition is in this case reduced to the orthogonality condition.

For this numerical example, we let $n(x, y) = 1 + y$, $P_a = (0, 0)$, and $y = \varphi(x) = x + 1$ which has the analytical solution $y(x) = -1 + d \cosh(x/d)$, in which d, b must satisfy transversality condition at $x = b$. The corresponding Euler-Lagrange equation for this problem is

$$y''(x) = \frac{1 + y'^2(x)}{1 + y(x)}. \tag{3.12}$$

By using (2.16), we obtain the following Adomian polynomials for $N(y) = (1 + y'^2)/(1 + y)$:

$$\begin{aligned} N_0(y_0) &= \frac{1 + y_0'^2}{1 + y_0}, & N_1(y_0, y_1) &= \frac{2y_1'y_0'(1 + y_0) - y_1(1 + y_0'^2)}{(1 + y_0)^2}, \\ N_2(y_0, y_1, y_2) &= \frac{B}{2(1 + y_0)^4}, \end{aligned} \tag{3.13}$$

in which

$$\begin{aligned} B &= (1 + y_0)^2(2y_1'^2(1 + y_0) + 4y_0'y_2'(1 + y_0) + 2y_0'y_1'y_1 - 2y_2(1 + y_0'^2) - 2y_1y_1'y_0') \\ &\quad - 2y_1(1 + y_0)(2y_0'y_1'(1 + y_0) - y_1(1 + y_0'^2)). \end{aligned} \tag{3.14}$$

Now we have the following recursive relations:

$$y_0(x) = Ax, \quad y_{n+1}(x) = L^{-1}(N_n(y_0, \dots, y_n)), \quad n \geq 0. \tag{3.15}$$

In this example, we use 2-term approximation of the solution, $\phi_2(x)$. For the determination of A and b , we solve the system of two equations $\phi_2(b) = b + 1$ and $\phi_2'(b) = -1$.

Figure 3.2 shows $y(x)$ and $\phi_2(x)$ and Table 3.2 shows the error of ϕ_2 .




FIGURE 3.2. Plot of $y(x)$ and $\phi_2(x)$ in example 2.

TABLE 3.2. Error ($e(x) = |y(x) - \phi_2(x)|$) in Example 2.

x	-0.3	0	0.3	0.6	1.25
$e(x)$.68239187e-3	.38763200e-1	.72982030e-1	.9805628e-1	.6617189e-2

4. Conclusion

In this work, the Adomian decomposition method was successfully used to solve variational problems with moving boundaries. We solve Euler-Lagrange equation with the boundary conditions that yield from natural boundary conditions and transversality conditions, using Adomian method. It is important to note that this method does not require any discretization or linearization and is not affected by computation roundoff errors.

References

- [1] U. Brechtken-Manderschied, *Introduction to the Calculus of Variations*, Translated from the German by P. G. Engstrom, Chapman and Hall Mathematics Series, Chapman & Hall, London, UK, 1991.
- [2] H. Sagan, *Introduction to the Calculus of Variations*, Dover, New York, NY, USA, 1992.
- [3] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, vol. 60 of *Fundamental Theories of Physics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1994.
- [4] G. Adomian, "A review of the decomposition method in applied mathematics," *Journal of Mathematical Analysis and Applications*, vol. 135, no. 2, pp. 501–544, 1988.

10 Mathematical Problems in Engineering

- [5] Y. Cherruault, "Convergence of Adomian's method," *Mathematical and Computer Modelling*, vol. 14, pp. 83–86, 1990.
- [6] Y. Cherruault, G. Saccomandi, and B. Some, "New results for convergence of Adomian's method applied to integral equations," *Mathematical and Computer Modelling*, vol. 16, no. 2, pp. 85–93, 1992.
- [7] M. Dehghan, "The use of adomian decomposition method for solving the one-dimensional parabolic equation with non-local boundary specifications," *International Journal of Computer Mathematics*, vol. 81, no. 1, pp. 25–34, 2004.
- [8] M. Dehghan, "Application of the Adomian decomposition method for two-dimensional parabolic equation subject to nonstandard boundary specifications," *Applied Mathematics and Computation*, vol. 157, no. 2, pp. 549–560, 2004.
- [9] M. Dehghan and M. Tatari, "The use of Adomian decomposition method for solving problems in calculus of variations," *Mathematical Problems in Engineering*, vol. 2006, Article ID 65379, 12 pages, 2006.
- [10] A. M. A. El-Sayed and M. Gaber, "The Adomian decomposition method for solving partial differential equations of fractal order in finite domains," *Physics Letters A*, vol. 359, no. 3, pp. 175–182, 2006.
- [11] I. Hashim, M. S. M. Noorani, and M. R. Said Al-Hadidi, "Solving the generalized Burgers-Huxley equation using the Adomian decomposition method," *Mathematical and Computer Modelling*, vol. 43, no. 11-12, pp. 1404–1411, 2006.
- [12] D. Kaya and S. M. El-Sayed, "Numerical soliton-like solutions of the potential Kadomtsev-Petviashvili equation by the decomposition method," *Physics Letters A*, vol. 320, no. 2-3, pp. 192–199, 2003.
- [13] D. Lesnic, "The decomposition method for Cauchy advection-diffusion problems," *Computers & Mathematics with Applications*, vol. 49, no. 4, pp. 525–537, 2005.
- [14] D. Lesnic, "The decomposition method for linear, one-dimensional, time-dependent partial differential equations," *International Journal of Mathematics and Mathematical Sciences*, vol. 2006, Article ID 42389, 29 pages, 2006.
- [15] R. Memarbashi, "Numerical solution of the Laplace equation in annulus by Adomian decomposition method," to appear in *Chaos, Solitons and Fractals (2006)*, doi:10.1016/j.chaos.2006.06.016.
- [16] S. Momani and Z. Odibat, "Analytical approach to linear fractional partial differential equations arising in fluid mechanics," *Physics Letters A*, vol. 355, no. 4-5, pp. 271–279, 2006.
- [17] S. Saha Ray and R. K. Bera, "Analytical solution of a fractional diffusion equation by Adomian decomposition method," *Applied Mathematics and Computation*, vol. 174, no. 1, pp. 329–336, 2006.
- [18] S. Saha Ray and R. K. Bera, "Approximate solution of a nonlinear fractional diffusion equation by Adomian decomposition method," *Applied Mathematics and Computation*, vol. 167, pp. 561–571, 2005.
- [19] M. Tatari and M. Dehghan, "The use of the Adomian decomposition method for solving multi-point boundary value problems," *Physica Scripta*, vol. 73, no. 6, pp. 672–676, 2006.
- [20] M. Tatari, M. Dehghan, and M. Razzaghi, "Application of the Adomian decomposition method for the Fokker-Planck equation," *Mathematical and Computer Modelling*, vol. 45, no. 5-6, pp. 639–650, 2007.
- [21] A.-M. Wazwaz, "A reliable modification of Adomian decomposition method," *Applied Mathematics and Computation*, vol. 102, no. 1, pp. 77–86, 1999.
- [22] A.-M. Wazwaz and A. Gorguis, "An analytic study of Fisher's equation by using Adomian decomposition method," *Applied Mathematics and Computation*, vol. 154, no. 3, pp. 609–620, 2004.

- [23] A.-M. Wazwaz, "The modified decomposition method and Padé approximants for a boundary layer equation in unbounded domain," *Applied Mathematics and Computation*, vol. 177, no. 2, pp. 737–744, 2006.
- [24] A.-M. Wazwaz, "A comparison between the variational iteration method and Adomian decomposition method," *Journal of Computational and Applied Mathematics*, vol. 207, no. 1, pp. 129–136, 2007.

Reza Memarbashi: Department of Mathematics, Faculty of Sciences, Semnan University,
P.O. Box 35195-363, Semnan, Iran
Email address: rmemar@semnan.ac.ir

Special Issue on Space Dynamics

Call for Papers

Space dynamics is a very general title that can accommodate a long list of activities. This kind of research started with the study of the motion of the stars and the planets back to the origin of astronomy, and nowadays it has a large list of topics. It is possible to make a division in two main categories: astronomy and astrodynamics. By astronomy, we can relate topics that deal with the motion of the planets, natural satellites, comets, and so forth. Many important topics of research nowadays are related to those subjects. By astrodynamics, we mean topics related to spaceflight dynamics.

It means topics where a satellite, a rocket, or any kind of man-made object is travelling in space governed by the gravitational forces of celestial bodies and/or forces generated by propulsion systems that are available in those objects. Many topics are related to orbit determination, propagation, and orbital maneuvers related to those spacecrafts. Several other topics that are related to this subject are numerical methods, nonlinear dynamics, chaos, and control.

The main objective of this Special Issue is to publish topics that are under study in one of those lines. The idea is to get the most recent researches and published them in a very short time, so we can give a step in order to help scientists and engineers that work in this field to be aware of actual research. All the published papers have to be peer reviewed, but in a fast and accurate way so that the topics are not outdated by the large speed that the information flows nowadays.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	July 1, 2009
First Round of Reviews	October 1, 2009
Publication Date	January 1, 2010

Lead Guest Editor

Antonio F. Bertachini A. Prado, Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, 12227-010 São Paulo, Brazil; prado@dem.inpe.br

Guest Editors

Maria Cecilia Zanardi, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; cecilia@feg.unesp.br

Tadashi Yokoyama, Universidade Estadual Paulista (UNESP), Rio Claro, 13506-900 São Paulo, Brazil; tadashi@rc.unesp.br

Silvia Maria Giuliatti Winter, São Paulo State University (UNESP), Guaratinguetá, 12516-410 São Paulo, Brazil; silvia@feg.unesp.br