

ON EXISTENCE OF KNESER SOLUTIONS OF A CERTAIN CLASS
OF n -TH ORDER NONLINEAR DIFFERENTIAL EQUATIONS

OLEG PALUMBÍNY, Trnava

(Received August 7, 1996)

Abstract. The paper deals with existence of Kneser solutions of n -th order nonlinear differential equations with quasi-derivatives.

Keywords: nonlinear differential equation, quasi-derivative, monotone solution, Kneser solution

MSC 1991: 34C10, 34D05

1. INTRODUCTION

The aim of our paper is to give some conditions for existence of Kneser solutions of the differential equation

$$(L) \quad L(y) \equiv 0,$$

where

$$\begin{aligned} L(y) &\equiv L_n y + \sum_{k=1}^{n-1} P_k(t) L_k y + P_0(t) f(y), \\ L_0 y(t) &= y(t), \\ L_1 y(t) &= p_1(t) (L_0 y(t))' = p_1(t) \frac{dy(t)}{dt}, \\ L_k y(t) &= p_k(t) (L_{k-1} y(t))' \quad \text{for } k = 2, 3, \dots, n-1, \\ L_n y(t) &= (L_{n-1} y(t))', \end{aligned}$$

n is an arbitrary positive integer, $n \geq 2$, $P_k(t)$, $k = 0, 1, \dots, n-1$, $p_i(t)$, $i = 1, 2, \dots, n-1$ are real-valued continuous functions on the interval $I_a = [a, \infty)$, $-\infty < a < \infty$; $f(t)$ is a real-valued function continuous on $E_1 = (-\infty, \infty)$.

If $n = 1$, then $L(y) \equiv L_1 y + P_0(t)f(y) = y' + P_0(t)f(y)$, $P_0(t)$ and $f(t)$ are real-valued continuous functions on I_a and on E_1 , respectively.

It is assumed throughout that

- (A) $P_k(t) \leq 0$, $p_i(t) > 0$ for all $t \in I_a$, $k = 0, 1, \dots, n-1$, $i = 1, 2, \dots, n-1$; $f(0) \neq 0$, $f(t) \geq 0$ for all $t \in E_1$; $P_0(t)$ is not identically zero in any subinterval of I_a ; n is an arbitrary positive integer, $n \geq 2$. If $n = 1$, then $P_0(t) \leq 0$ and $f(t) \geq 0$ for all $t \in I_a$ and E_1 , respectively.

The problems of existence of monotone or Kneser solutions for third order ordinary differential equations with quasi-derivatives were studied in several papers ([5], [7], [8], [10]). The equation (L), where $p_i(t) \equiv 1$, $i = 1, 2, 3$ ($n = 4$) was studied, for example, in ([6], [9], [12]). Equations of the fourth order with quasi-derivatives were also studied, for instance, in ([1], [3], [13]).

Existence of monotone solutions for n -th order equations with quasi-derivatives was studied in [4].

In our paper, Theorem 1 and Theorem 2 give sufficient conditions for existence of a Kneser solution of (L) on $[a, \infty)$ for n an even number or for an odd one, respectively.

Now we explain the concept of a Kneser solution, and other useful ones:

Definition 1. A nontrivial solution $y(t)$ of a differential equation of the n -th order is called a Kneser solution on $I_a = [a, \infty)$ iff $(y(t) > 0, (-1)^k L_k y(t) \geq 0)$ or $(y(t) < 0, (-1)^k L_k y(t) \leq 0)$ for all $t \in I_a$, $k = 1, 2, \dots, n-1$.

Definition 2. Let J be an arbitrary type of an interval with endpoints t_1, t_2 , where $-\infty \leq t_1 < t_2 \leq \infty$. The interval J is called the maximum interval of existence of $u: J \rightarrow E_1^n$, where $u(t)$ is a solution of the differential system $u' = F(t, u)$ iff $u(t)$ can be continued neither to the right nor to the left of J .

Definition 3. Let $y' = U(t, y)$ be a scalar differential equation. Then $y_0(t)$ is called the maximum solution of the Cauchy problem

$$(*) \quad y' = U(t, y), \quad y(t_0) = y_0$$

iff $y_0(t)$ is a solution of (*) on the maximum interval of existence and if $y(t)$ is another solution of (*), then $y(t) \leq y_0(t)$ for all t belonging to the common interval of existence of $y(t)$ and $y_0(t)$.

We give some preliminary results.

Lemma 1. Let $A(t, s)$ be a nonpositive and continuous function for $a \leq t \leq s \leq t_0$. If $g(t)$, $\psi(t)$ are continuous functions in the interval $[a, t_0]$ and

$$\psi(t) \geq g(t) + \int_{t_0}^t A(t, s)\psi(s) ds \quad \text{for } t \in [a, t_0],$$

then every solution $y(t)$ of the integral equation

$$y(t) = g(t) + \int_{t_0}^t A(t, s)y(s) ds$$

satisfies the inequality $y(t) \leq \psi(t)$ in $[a, t_0]$.

Proof. See [6], page 331. □

Lemma 2. (Wintner) Let $U(t, u)$ be a continuous function on a domain $t_0 \leq t \leq t_0 + \alpha$, $\alpha > 0$, $u \geq 0$, let $u(t)$ be a maximum solution of the Cauchy problem $u' = U(t, u)$, $u(t_0) = u_0 \geq 0$ ($u' = U(t, u)$ is a scalar differential equation) existing on $[t_0, t_0 + \alpha]$; for example, let $U(t, u) = \psi(u)$, where $\psi(u)$ is a continuous and positive function for $u \geq 0$ such that

$$\int^{\infty} \frac{du}{\psi(u)} = \infty.$$

Let us assume $f(t, y)$ to be continuous on $t_0 \leq t \leq t_0 + \alpha$, $y \in E_1^n$, y arbitrary, and to satisfy the condition

$$|f(t, y)| \leq U(t, |y|).$$

Then the maximum interval of existence of a solution of the Cauchy problem

$$y' = f(t, y), \quad y(t_0) = y_0,$$

where $|y_0| \leq u_0$, is $[t_0, t_0 + \alpha]$.

Proof. See [2], Theorem III.5.1. □

Lemma 3. Let (A) hold, and let there exist real nonnegative constants a_1, a_2 such that $f(t) \leq a_1|t| + a_2$ for all $t \in E_1$. Let initial values $L_k y(a) = b_k$ be given for $k = 0, 1, \dots, n-1$. Then there exists a solution $y(t)$ of (L) on $[a, \infty)$, which fulfils these initial conditions.

Proof. See [4], Lemma 3. □

2. RESULTS

Lemma 4. *Let us assume $g(t, z)$ to be continuous on $t_0 - \alpha \leq t \leq t_0$, α a positive constant, $z \in E_1^n$, z is arbitrary and satisfies a condition*

$$|g(t, z)| \leq \psi(|z|),$$

where $\psi(t)$ is a continuous and positive function for $t \geq 0$ such that

$$\int^{\infty} \frac{dt}{\psi(t)} = \infty.$$

Then the maximum interval of existence of a solution of the Cauchy problem

$$z' = g(t, z), \quad z(t_0) = z_0,$$

is $[t_0 - \alpha, t_0]$.

Proof. Let us consider the Cauchy problem

$$(u) \quad u' = \psi(u), \quad u(-t_0) = u_0 = |z_0|.$$

According to the assumptions, the problem (u) admits a single solution $u_0(t)$ on $[-t_0, \infty)$, where

$$u_0(t) = R_{-1}(t + t_0)$$

and $R: [u_0, \infty) \rightarrow [0, \infty)$, $R(u) = \int_{u_0}^u \frac{1}{\psi(t)} dt$, $R_{-1}(R(u)) = u$, $u \in [u_0, \infty)$. Let us consider the Cauchy problems

$$(U) \quad u' = U(t, u) = \psi(u), \quad u(-t_0) = u_0 = |z_0|, \quad (t, u) \in [-t_0, -t_0 + \alpha] \times [0, \infty),$$

$$(y) \quad y'(t) = g(-t, -y), \quad y(-t_0) = -z_0, \quad (t, y) \in [-t_0, -t_0 + \alpha] \times E_1^n,$$

$$(z) \quad z'(t) = g(t, z), \quad z(t_0) = z_0, \quad (t, z) \in [t_0 - \alpha, t_0] \times E_1^n.$$

Then $u_0(t) = R_{-1}(t + t_0)$ is the maximum solution of (U) on the maximum interval of existence $[-t_0, -t_0 + \alpha]$. According to Lemma 2 there exists a solution $y_0(t)$ of (y) on $[-t_0, -t_0 + \alpha]$. Then the Cauchy problem (z) admits the solution $z_0(t) = -y_0(-t)$ on $[t_0 - \alpha, t_0]$ because of

$$z'_0(t) = y'_0(-t) = g(t, -y_0(-t)) = g(t, z_0(t))$$

on $[t_0 - \alpha, t_0]$. So the maximum interval of existence of (z) is $[t_0 - \alpha, t_0]$. □

Lemma 5. *Let (A) hold, and let there exist nonnegative real constants a_1, a_2 such that $f(t) \leq a_1|t| + a_2$ for all $t \in E_1$. Let initial values $L_k y(t_0) = b_k$ be given for $k = 0, 1, \dots, n-1$, $t_0 > a$. Then there exists a solution $y(t)$ of (L) on $[a, \infty)$, which fulfils these initial conditions.*

Proof. According to Lemma 3 there exists a solution of (L) on $[t_0, \infty)$ such that the initial conditions hold. To prove our lemma we need to prove existence of a solution $y(t)$ of (L) on $[a, t_0]$ satisfying the given initial conditions. Consider now the following system (S), which corresponds to the equation (L):

$$(S) \quad \begin{aligned} u'_k(t) &= \frac{u_{k+1}(t)}{p_k(t)}, \quad k = 1, 2, \dots, n-1, \\ u'_n(t) &= - \sum_{k=1}^{n-1} P_k(t)u_{k+1}(t) - P_0(t)f(u_1(t)), \end{aligned}$$

where $u_k(t) = L_{k-1}y(t)$, $k = 1, 2, \dots, n$, $f_k = u_{k+1}/p_k$, $k = 1, \dots, n-1$, $f_n = - \sum P_k u_{k+1} - P_0 f(u_1)$, $F = (f_1, f_2, \dots, f_n)$, $u = (u_1, u_2, \dots, u_n)$, $u' = (u'_1, u'_2, \dots, u'_n)$, $|u| = \sum_{k=1}^n |u_k|$, $|F| = \sum_{k=1}^n |f_k|$, $(t, u) \in [a, t_0] \times E_1^n$. Then

$$\begin{aligned} |F(t, u)| &= \sum_{k=1}^{n-1} \left| \frac{u_{k+1}}{p_k} \right| + \left| - \sum_{k=1}^{n-1} P_k u_{k+1} - P_0 f(u_1) \right| \\ &\leq \sum_{k=1}^{n-1} \left(-P_k + \frac{1}{p_k} \right) |u_{k+1}| - P_0 (a_1 |u_1| + a_2) \leq K_1 |u| + K_2 = \psi(|u|), \end{aligned}$$

where K_1, K_2 are appropriate positive real constants. It is obvious that

$$\int_a^\infty \frac{ds}{\psi(s)} = \infty$$

for $s \in E_1$, $s > 0$. Lemma 4 yields existence of a solution of (S) on $[a, t_0]$. This fact implies existence of a solution $y(t)$ of the equation (L) on $[a, t_0]$ which satisfies the given initial conditions. The lemma is proved. \square

Lemma 6. *Let (A) hold, and let $y(t)$ be a solution of (L) on $[t_1, \infty)$, where $t_1 \geq a$. Let (B) hold, where $(s_0 = s)$*

$$(B) \quad \sum_{k=1}^{n-1} (-1)^{k-1} M_k(t, s) \leq 0, \quad N_n(t) \leq 0, \quad n \geq 2$$

and

$$\begin{aligned}
M_k(t, s) &= \int_t^s \frac{ds_1}{p_{n-2}(s_1)} \int_t^{s_1} \frac{ds_2}{p_{n-3}(s_2)} \cdots \int_t^{s_{k-2}} \frac{-P_{n-k}(s_{k-1})}{p_{n-1}(s)} ds_{k-1}, \\
M_1(t, s) &= -P_{n-1}(s), \quad N_n(t) = \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)G_k(s)) ds, \\
G_k(s) &= L_{n-k}y(t_2) + (-1)^1 L_{n-k+1}y(t_2) \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} + (-1)^2 L_{n-k+2}y(t_2) \\
&\quad \times \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} + \dots + (-1)^{k-2} L_{n-2}y(t_2) \\
&\quad \times \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} \cdots \int_{s_{k-3}}^{t_2} \frac{ds_{k-2}}{p_{n-2}(s_{k-2})}
\end{aligned}$$

for $k = 2, 3, \dots, n-1$, $G_1(s) = 0$.

- a) Let n be an even number and $t_2 \in (t_1, \infty)$ such that $(-1)^k L_k y(t_2) \geq 0$ for $k = 0, 1, \dots, n-1$. Then $(-1)^k L_k y(t) \geq 0$ for $t \in [t_1, t_2]$, $k = 0, 1, \dots, n-1$.
- b) Let n be an odd number and $t_2 \in (t_1, \infty)$ such that $(-1)^k L_k y(t_2) \leq 0$ for $k = 0, 1, \dots, n-1$. Then $(-1)^k L_k y(t) \leq 0$ for $t \in [t_1, t_2]$, $k = 0, 1, \dots, n-1$.

Proof. Let $n \geq 2$. Integration of the identity $L_n y = (L_{n-1} y)'$ over $[t_2, t]$, where $t_1 \leq t \leq t_2$ (n can be an even number as well as an odd one) yields

$$\begin{aligned}
&L_{n-1}y(t) \\
&= L_{n-1}y(t_2) - \int_{t_2}^t \sum_{k=1}^{n-1} P_k(s)L_k y(s) ds - \int_{t_2}^t P_0(s)f(y(s)) ds \\
&= L_{n-1}y(t_2) + \int_{t_2}^t (-P_0(s)f(y(s))) ds + \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)L_{n-k}y(s)) ds.
\end{aligned}$$

Let us denote the expression $L_{n-1}y(t_2) + \int_{t_2}^t (-P_0(s)f(y(s))) ds$ by $K_n(t)$. It is obvious that $K_n(t) \leq 0$ for all $t \in [t_1, t_2]$. We have

$$L_{n-1}y(t) = K_n(t) + \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)L_{n-k}y(s)) ds.$$

It can be proved that

$$\begin{aligned}
& L_{n-k}y(s) \\
&= L_{n-k}y(t_2) + L_{n-k+1}y(t_2) \int_{t_2}^s \frac{ds_1}{p_{n-k+1}(s_1)} \\
&+ L_{n-k+2}y(t_2) \int_{t_2}^s \frac{ds_1}{p_{n-k+1}(s_1)} \int_{t_2}^{s_1} \frac{ds_2}{p_{n-k+2}(s_2)} + \dots \\
&+ L_{n-2}y(t_2) \int_{t_2}^s \frac{ds_1}{p_{n-k+1}(s_1)} \int_{t_2}^{s_1} \frac{ds_2}{p_{n-k+2}(s_2)} \dots \int_{t_2}^{s_{k-3}} \frac{ds_{k-2}}{p_{n-2}(s_{k-2})} \\
&+ \int_{t_2}^s \frac{ds_1}{p_{n-k+1}(s_1)} \int_{t_2}^{s_1} \frac{ds_2}{p_{n-k+2}(s_2)} \int_{t_2}^{s_2} \frac{ds_3}{p_{n-k+3}(s_3)} \dots \int_{t_2}^{s_{k-2}} \frac{L_{n-1}y(s_{k-1}) ds_{k-1}}{p_{n-1}(s_{k-1})}
\end{aligned}$$

for $k = 2, 3, \dots, n-1$. By interchanging the upper and the lower bounds in the previous integrals, we have

$$\begin{aligned}
& L_{n-k}y(s) \\
&= L_{n-k}y(t_2) + (-1)^1 L_{n-k+1}y(t_2) \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \\
&+ (-1)^2 L_{n-k+2}y(t_2) \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} + \dots \\
&+ (-1)^{k-2} L_{n-2}y(t_2) \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} \dots \int_{s_{k-3}}^{t_2} \frac{ds_{k-2}}{p_{n-2}(s_{k-2})} \\
&+ (-1)^{k-1} \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} \dots \int_{s_{k-2}}^{t_2} \frac{L_{n-1}y(s_{k-1}) ds_{k-1}}{p_{n-1}(s_{k-1})}.
\end{aligned}$$

Denoting the last $(k-1)$ -dimensional integral by $I_k(s)$, the previous sum by $G_k(s)$, $I_1(s) = L_{n-1}y(s)$, $G_1(s) = 0$ for $k = 1, 2, \dots, n-1$ ($s_0 = s$) we obtain

$$L_{n-k}y(s) = G_k(s) + (-1)^{k-1} I_k(s).$$

Hence

$$\begin{aligned}
& L_{n-1}y(t) \\
&= K_n(t) + \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)[G_k(s) + (-1)^{k-1} I_k(s)]) ds \\
&= K_n(t) + \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)G_k(s)) ds + \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)(-1)^{k-1} I_k(s)) ds.
\end{aligned}$$

Denoting $K_n(t) + \int_t^{t_2} \sum_{k=1}^{n-1} (-P_{n-k}(s)G_k(s)) ds$ by $g_n(t)$ and denoting $\int_t^{t_2} (-P_{n-k}(s) \times (-1)^{k-1} I_k(s)) ds$ by $(-1)^{k-1} J_k(t)$ we have

$$L_{n-1}y(t) = g_n(t) + \sum_{k=1}^{n-1} (-1)^{k-1} J_k(t),$$

where $J_k(t)$ is the k -dimensional integral

$$J_k(t) = - \int_t^{t_2} (-P_{n-k}(s)) ds \int_s^{t_2} \frac{ds_1}{p_{n-k+1}(s_1)} \int_{s_1}^{t_2} \frac{ds_2}{p_{n-k+2}(s_2)} \cdots \\ \cdots \int_{s_{k-2}}^{t_2} \frac{L_{n-1}y(s_{k-1}) ds_{k-1}}{p_{n-1}(s_{k-1})}$$

for $k = 2, 3, \dots, n-1$ and $J_1(t) = - \int_t^{t_2} (-P_{n-1}(s)L_{n-1}y(s)) ds$.

By changing the notation of the variables we have

$$J_k(t) = - \int_t^{t_2} (-P_{n-k}(s_{k-1})) ds_{k-1} \int_{s_{k-1}}^{t_2} \frac{ds_{k-2}}{p_{n-k+1}(s_{k-2})} \int_{s_{k-2}}^{t_2} \frac{ds_{k-3}}{p_{n-k+2}(s_{k-3})} \cdots \\ \cdots \int_{s_1}^{t_2} \frac{L_{n-1}y(s) ds}{p_{n-1}(s)}.$$

$J_k(t)$ is a k -dimensional integral on a k -dimensional domain. This domain can be described as an elementary domain in the following way:

$$\begin{aligned} t &\leq s_{k-1} \leq t_2 \\ s_{k-1} &\leq s_{k-2} \leq t_2 \\ s_{k-2} &\leq s_{k-3} \leq t_2 \\ &\vdots \\ s_2 &\leq s_1 \leq t_2 \\ s_1 &\leq s \leq t_2, \end{aligned}$$

as well as like

$$\begin{aligned} t &\leq s \leq t_2 \\ t &\leq s_1 \leq s \\ t &\leq s_2 \leq s_1 \\ &\vdots \\ t &\leq s_{k-2} \leq s_{k-3} \\ t &\leq s_{k-1} \leq s_{k-2} \end{aligned}$$

for $k = 2, 3, \dots, n - 1$. Hence

$$J_k(t) = - \int_t^{t_2} L_{n-1}y(s) ds \int_t^s \frac{ds_1}{p_{n-2}(s_1)} \int_t^{s_1} \frac{ds_2}{p_{n-3}(s_2)} \cdots \int_t^{s_{k-2}} \frac{-P_{n-k}(s_{k-1})}{p_{n-1}(s)} ds_{k-1}.$$

The last integral can be rewritten into the form

$$J_k(t) = - \int_t^{t_2} M_k(t, s) L_{n-1}y(s) ds = \int_{t_2}^t M_k(t, s) L_{n-1}y(s) ds,$$

where

$$M_k(t, s) = \int_t^s \frac{ds_1}{p_{n-2}(s_1)} \int_t^{s_1} \frac{ds_2}{p_{n-3}(s_2)} \cdots \int_t^{s_{k-2}} \frac{-P_{n-k}(s_{k-1})}{p_{n-1}(s)} ds_{k-1}$$

for $k = 2, 3, \dots, n - 1$, $M_1(t, s) = -P_{n-1}(s)$. Hence

$$\begin{aligned} L_{n-1}y(t) &= g_n(t) + \sum_{k=1}^{n-1} (-1)^{k-1} J_k(t) = g_n(t) + \sum_{k=1}^{n-1} (-1)^{k-1} \int_{t_2}^t M_k(t, s) L_{n-1}y(s) ds \\ &= g_n(t) + \int_{t_2}^t \left[\sum_{k=1}^{n-1} (-1)^{k-1} M_k(t, s) \right] L_{n-1}y(s) ds = g_n(t) + \int_{t_2}^t A_n(t, s) L_{n-1}y(s) ds, \end{aligned}$$

where $A_n(t, s) = \sum_{k=1}^{n-1} (-1)^{k-1} M_k(t, s)$. We note that $s \leq t_2$, $s_i \leq t_2$, $t \leq s$, $t \leq s_i$ for $i = 1, 2, \dots, n - 3$. According to the assumptions of the lemma, we have $g_n(t) = K_n(t) + N_n(t)$ and $g_n(t) \leq 0$, $A_n(t, s) \leq 0$. According to Lemma 1 we have $L_{n-1}y(t) \leq 0$ for all $t \in [t_1, t_2]$. By virtue of

$$L_{n-2}y(t) = L_{n-2}y(t_2) + \int_{t_2}^t \frac{L_{n-1}y(s)}{p_{n-1}(s)} ds \geq L_{n-2}y(t_2) \geq 0,$$

we have $L_{n-2}y(t) \geq 0$ on $[t_1, t_2]$. By using of a similar procedure (n can be an even number or an odd one), we get for $n \geq 2$:

- a) $(-1)^k L_k y(t) \geq 0$ on $[t_1, t_2]$ for $k = 0, 1, \dots, n - 1$, for n an even number,
- b) $(-1)^k L_k y(t) \leq 0$ on $[t_1, t_2]$ for $k = 0, 1, \dots, n - 1$, for n an odd number.

If $n = 1$, then the assertion of the lemma is obvious. \square

Lemma 7. Consider a solution $y(t)$ of (L) on $[t_1, \infty)$, $t_1 \geq a$ such that (A) holds, let n be an even number and $t_2 \in (t_1, \infty)$ such that $(-1)^k L_k y(t_2) \geq 0$ for $k = 0, 1, \dots, n-1$. Let $P_k(t) \equiv 0$ on $[t_1, t_2]$ for all odd integers $k \in [1, n]$. Then (B) holds.

Proof. We have $G_k(s) \geq 0$ for all even numbers $k \in [1, n]$, and $G_k(s) \leq 0$ for all odd ones. If k is an odd number, then $n-k$ is an odd number too, and $P_{n-k}(t) \equiv 0$ on $[t_1, t_2]$. Therefore $N_n(t) = \int_{t_2}^t \sum_{k=1}^{n-1} (-P_{n-k}(s)G_k(s)) ds \leq 0$. Similarly, $M_k(t, s) = 0$ for all odd $k \leq n$. So $A_n(t, s) = \sum_{k=1}^{n-1} (-1)^{k-1} M_k(t, s) \leq 0$ because $M_k(t, s) \geq 0$ for all $k = 1, 2, \dots, n-1$. \square

Lemma 8. Consider a solution $y(t)$ of (L) on $[t_1, \infty)$, $t_1 \geq a$ such that (A) holds, let $n > 1$ be an odd number and $t_2 \in (t_1, \infty)$ such that $(-1)^k L_k y(t_2) \leq 0$ for $k = 0, 1, \dots, n-1$. Let $P_k(t) \equiv 0$ on $[t_1, t_2]$ for all even integers $k \in [1, n]$. Then (B) holds.

Proof. The proof is similar to the proof of the previous lemma, so it is omitted. \square

Lemma 9. Let $\{y_m(t)\}_{m=n_0}^\infty$ be a sequence of solutions of (L) on $[t_0, \infty)$, where $a < t_0 < n_0$, n is an even number, and $L_k y_m(m) = (-1)^k$ for all $m \geq n_0$, $k = 0, 1, \dots, n-1$. Let (A) hold, and let $P_k(t) \equiv 0$ on $[a, \infty)$ for all odd integer numbers $k \in [1, n]$. Let $-\infty < \int_{t_0}^\infty P_0(s) ds = P < 0$, $\int_{t_0}^\infty P_k(s) ds \geq -\frac{1}{2}$ for $k = 1, 2, \dots, n-1$, let P_k be nondecreasing functions for $k = 0, 1, \dots, n-1$, $\int_{t_0}^\infty 1/p_r(s) ds \leq \frac{1}{2}$ for $r = 1, 2, \dots, n-1$, and let K be a real positive constant such that $0 \leq f(t) \leq K$ for $t \in (-\infty, \infty)$. Then there exists a subsequence of $\{y_m(t)\}_{m=n_0}^\infty$ which converges to $\varphi_0(t)$. This function $\varphi_0(t)$ is a solution of (L) on $[t_0, \infty)$, and $(-1)^k L_k \varphi_0(t) \geq 0$ on $[t_0, \infty)$ for $k = 0, 1, \dots, n-1$.

Proof. Because $L_n y_m(t) \geq 0$ on $[t_0, m]$ for $m = n_0, n_0 + 1, \dots$ (this follows from Lemma 7 and Lemma 6, part a)), we have that $L_{n-1} y_m(t)$ is nondecreasing and negative on $[t_0, n_0]$ for $m > n_0$. If we prove that $L_{n-1} y_m(t_0)$ is bounded from below, it means $L_{n-1} y_m(t)$ is uniformly bounded on $[t_0, n_0]$. Using the expression (C) several times, where

$$(C) \quad L_k y_m(s) = L_k y_m(m) + \int_m^s \left(L_{k+1} \frac{y_m(s)}{p_{k+1}(s)} \right) ds \text{ for } k = 0, 1, \dots, n-2,$$

we obtain for $n > 3$, $2 \leq k < n - 1$ ($s_0 = s$):

$$\begin{aligned}
L_k y_m(s) &= L_k y_m(m) + L_{k+1} y_m(m) \int_m^s \frac{ds_1}{p_{k+1}(s_1)} \\
&+ L_{k+2} y_m(m) \int_m^s \frac{ds_1}{p_{k+1}(s_1)} \int_m^{s_1} \frac{ds_2}{p_{k+2}(s_2)} + \dots \\
&+ L_{n-2} y_m(m) \int_m^s \frac{ds_1}{p_{k+1}(s_1)} \int_m^{s_1} \frac{ds_2}{p_{k+2}(s_2)} \dots \int_m^{s_{n-k-3}} \frac{ds_{n-k-2}}{p_{n-2}(s_{n-k-2})} \\
&+ \int_m^s \frac{ds_1}{p_{k+1}(s_1)} \int_m^{s_1} \frac{ds_2}{p_{k+2}(s_2)} \dots \int_m^{s_{n-k-2}} \frac{L_{n-1} y_m(s_{n-k-1})}{p_{n-1}(s_{n-k-1})} ds_{n-k-1}.
\end{aligned}
\tag{D}$$

Integration of (L) over $[t_0, m]$ yields

$$\begin{aligned}
&L_{n-1} y_m(t_0) \\
&= L_{n-1} y_m(m) + \int_{t_0}^m P_0(s) f(y_m(s)) ds + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) L_{2k} y_m(s) ds \\
&= L_{n-1} y_m(m) + \int_{t_0}^m P_0(s) f(y_m(s)) ds + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) [B_{2k}(s) + C_{2k}(s)] ds,
\end{aligned}$$

where $C_k(s)$ is the last integral in (D) and $B_k(s)$ is the rest of the right-hand side of (D). Let us denote the expression $L_{n-1} y_m(m) + \int_{t_0}^m P_0(s) f(y_m(s)) ds$ by F_m . Then

$$\begin{aligned}
&L_{n-1} y_m(t_0) \\
&= F_m + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) B_{2k}(s) ds + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) C_{2k}(s) ds \\
&\geq F_m + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) B_{2k}(s) ds + L_{n-1} y_m(t_0) \\
&\quad \times \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) \left[\int_m^s \frac{ds_1}{p_{2k+1}(s_1)} \int_m^{s_1} \frac{ds_2}{p_{2k+2}(s_2)} \dots \int_m^{s_{n-2k-2}} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})} \right] ds \\
&\geq F_m + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) B_{2k}(s) ds + L_{n-1} y_m(t_0) \\
&\quad \times \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} \left[-P_{2k}(s) \left[\int_{t_0}^{\infty} \frac{ds_1}{p_{2k+1}(s_1)} \int_{t_0}^{\infty} \frac{ds_2}{p_{2k+2}(s_2)} \dots \int_{t_0}^{\infty} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})} \right] \right] ds.
\end{aligned}$$

(We have used the fact that the last integral has the dimension $n - 2k$, which is an even number, and $t_0 \leq s_i \leq m < \infty$ for $i = 1, 2, \dots, n - 2k - 2$, $t_0 \leq s \leq m < \infty$). An easy arrangement yields

$$L_{n-1}y_m(t_0) \left[1 + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} P_{2k}(s) ds \int_{t_0}^{\infty} \frac{ds_1}{p_{2k+1}(s_1)} \int_{t_0}^{\infty} \frac{ds_2}{p_{2k+2}(s_2)} \cdots \right. \\ \left. \cdots \int_{t_0}^{\infty} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})} \right] \geq F_m + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) B_{2k}(s) ds.$$

According to the assumptions, the expression in the parentheses above is a positive number because of $\sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} [-P_{2k}(s)] ds \cdots \int_{t_0}^{\infty} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})} \leq \sum_{k=1}^{\frac{n}{2}-1} (\frac{1}{2})^{n-2k} < 1$. Therefore

$$L_{n-1}y_m(t_0) \geq \frac{F_m + \sum_{k=1}^{\frac{n}{2}-1} \int_{k_0}^m P_{2k}(s) B_{2k}(s) ds}{1 + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} P_{2k}(s) ds \int_{t_0}^{\infty} \frac{ds_1}{p_{2k+1}(s_1)} \cdots \int_{t_0}^{\infty} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})}}.$$

We have

$$F_m = L_{n-1}y_m(m) + \int_{t_0}^m P_0(s) f(y_m(s)) ds \geq -1 + \int_{t_0}^{\infty} P_0(s) f(y_m(s)) ds \\ \geq -1 + K \int_{t_0}^{\infty} P_0(s) ds = -1 + KP,$$

$$B_{2k}(s) = L_{2k}y_m(m) + L_{2k+1}y_m(m) \int_m^s \frac{ds_1}{p_{2k+1}(s_1)} + \cdots + L_{n-2}y_m(m) \int_m^s \frac{ds_1}{p_{2k+1}(s_1)} \cdots \\ \cdots \int_m^{s_{n-2k-3}} \frac{ds_{n-2k-2}}{p_{n-2}(s_{n-2k-2})} = 1 + 1 \int_s^m \frac{ds_1}{p_{2k+1}(s_1)} + \cdots + 1 \int_s^m \frac{ds_1}{p_{2k+1}(s_1)} \cdots \\ \cdots \int_{s_{n-2k-3}}^m \frac{ds_{n-2k-2}}{p_{n-2k-2}(s_{n-2k-2})} \leq 1 + (n - 2k - 2) \frac{1}{2} \leq n$$

because of $s \leq m$, $s_i \leq m$ for $i = 1, 2, \dots, n - 2k - 3$. So we have

$$\sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) B_{2k}(s) ds \geq n \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^m P_{2k}(s) ds \\ \geq n \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} P_{2k}(s) ds \geq -n \left(\frac{n}{2} - 1 \right) \frac{1}{2}.$$

Hence

$$\begin{aligned} L_{n-1}y_m(t_0) &\geq \frac{-1 + KP - \frac{n}{2}(\frac{n}{2} - 1)}{1 + \sum_{k=1}^{\frac{n}{2}-1} \int_{t_0}^{\infty} P_{2k}(s) \, ds \int_{t_0}^{\infty} \frac{ds_1}{p_{2k+1}(s_1)} \cdots \int_{t_0}^{\infty} \frac{ds_{n-2k-1}}{p_{n-1}(s_{n-2k-1})}} \\ &= S_{n-1} \in (-\infty, 0) \end{aligned}$$

for $n > 3$. If $n = 2$, then $L_{n-1}y_m(t_0) = F_m \geq -1 + KP \in (-\infty, 0)$. It implies that $\{L_{n-1}y_m(t_0)\}_{m=n_0}^{\infty}$ is bounded from below for any fixed even number $n \geq 2$. So we have

$$\begin{aligned} 0 \leq L_{n-2}y_m(t_0) &= L_{n-2}y_m(m) + \int_{t_0}^m \frac{-L_{n-1}y_m(s)}{p_{n-1}(s)} \, ds \leq 1 - L_{n-1}y_m(t_0) \int_{t_0}^{\infty} \frac{ds}{p_{n-1}(s)} \\ &\leq 1 - S_{n-1} \int_{t_0}^{\infty} \frac{ds}{p_{n-1}(s)} = S_{n-2} \in (0, \infty), \\ 0 \geq L_{n-3}y_m(t_0) &= L_{n-3}y_m(m) + \int_{t_0}^m \frac{-L_{n-2}y_m(s)}{p_{n-2}(s)} \, ds \geq -1 - L_{n-2}y_m(t_0) \int_{t_0}^{\infty} \frac{ds}{p_{n-2}(s)} \\ &\geq -1 - S_{n-2} \int_{t_0}^{\infty} \frac{ds}{p_{n-2}(s)} = S_{n-3} \in (-\infty, 0). \end{aligned}$$

Similarly, it can be proved that $\{L_k y_m(t_0)\}_{m=n_0}^{\infty}$ is bounded for $k = 0, 1, \dots, n-1$. However,

$$\begin{aligned} 0 \leq L_n y_m(t) &= - \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t) L_{2k} y_m(t) - P_0(t) f(y_m(t)) \\ &\leq - \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t_0) L_{2k} y_m(t_0) - P_0(t_0) K \\ &\leq - \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t_0) S_{2k} - P_0(t_0) K = S_n \in (0, \infty), \end{aligned}$$

and this implies that $\{L_n y_m(t)\}_{m=n_0}^{\infty}$ is uniformly bounded on $[t_0, n_0]$ for $m \geq n_0$ and so $L_{n-1}y_m(t)$ are uniformly equicontinuous on $[t_0, n_0]$ for $m \geq n_0$. According to Arzelà-Ascoli theorem, there exists a subsequence $\{L_{n-1}y_{k_m}\}_{m=n_0}^{\infty}$ of $\{L_{n-1}y_m\}_{m=n_0}^{\infty}$ such that $\{L_{n-1}y_{k_m}\}_{m=n_0}^{\infty}$ converges uniformly on $[t_0, n_0]$ to, for example, a function $\varphi_{n-1}(t)$.

To ensure uniform convergence of $\{L_{n-2}y_{k_m}\}_{m=n_0}^\infty$ on $[t_0, n_0]$ to, for instance, a function $\varphi_{n-2}(t)$, it suffices to show convergence of $\{L_{n-2}y_{k_m}\}_{m=n_0}^\infty$ at an inner point of $[t_0, n_0]$. This follows from the fact that $L_{n-2}y_{k_m}(t_0 + \varepsilon) \leq L_{n-2}y_{k_m}(t_0) \leq S_{n-2}$ for $\varepsilon > 0$, $\varepsilon < n_0 - t_0$. Then there exists a convergent subsequence $\{L_{n-2}y_{k_{l_m}}(t_0 + \varepsilon)\}_{m=n_0}^\infty$ of $\{L_{n-2}y_{k_m}(t_0 + \varepsilon)\}_{m=n_0}^\infty$ and therefore $\{L_{n-2}y_{k_{l_m}}\}_{m=n_0}^\infty$ converges uniformly to $\varphi_{n-2}(t)$ on $[t_0, n_0]$. It is obvious that $L_{n-1}y_{k_{l_m}} \rightrightarrows \varphi_{n-1}$ on $[t_0, n_0]$, too. In a similar way we can prove uniform convergence of a subsequence $\{y_{r_m}\}_{m=n_0}^\infty$ of $\{y_m\}_{m=n_0}^\infty$ such that $L_k y_{r_m}(t) \rightrightarrows \varphi_k(t)$ on $[t_0, n_0]$ for $k = 0, 1, \dots, n$. Due to the fact that uniform convergence makes changing of the order of limit processes possible (a quasi-derivative is a certain kind of limit), we have

$$\begin{aligned} 0 &= \lim_{m \rightarrow \infty} L(y_{r_m}(t)) \\ &= \lim_{m \rightarrow \infty} L_n y_{r_m}(t) + \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t) \lim_{m \rightarrow \infty} L_{2k} y_{r_m}(t) + P_0(t) f(\lim_{m \rightarrow \infty} y_{r_m}(t)) \\ &= \varphi_n(t) + \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t) \varphi_{2k}(t) + P_0(t) f(\varphi_0(t)) \end{aligned}$$

for all $t \in [t_0, n_0]$.

But $\varphi_k(t) = \lim_{m \rightarrow \infty} L_k y_{r_m}(t) = L_k(\lim_{m \rightarrow \infty} y_{r_m}(t)) = L_k(\lim_{m \rightarrow \infty} L_0 y_{r_m}(t)) = L_k \varphi_0(t)$, so $\varphi_0(t)$ fulfils (L) on $[t_0, n_0]$. It is important that we are able to continue $\varphi_0(t)$ on $[t_0, n_0+1]$ in such a way that $\varphi_0(t)$ be a solution of (L) on $[t_0, n_0+1]$. Indeed, it suffices to repeat the whole previous part of the proof with the sequence y_{r_m} for $m \geq n_0 + 1$ instead of y_m for $m \geq n_0$. Now it is obvious that $\varphi_0(t)$ can be continued on $[t_0, n_0+v]$ (v is an arbitrary integer greater than 1) and therefore $\varphi_0(t)$ fulfils (L) on $[t_0, \infty)$. Now let us take an arbitrary point $t_1 \in [t_0, \infty)$. Then there exists $m_0 \in \{1, 2, \dots\}$, $t_1 < m_0$ and a subsequence $\{y_{s_m}\}_{m=n_0}^\infty$ of $\{y_m\}_{m=n_0}^\infty$ such that $L_k y_{s_m} \rightrightarrows L_k \varphi_0(t)$ on $[t_0, m_0]$. But $(-1)^k L_k y_{s_m}(t) \geq 0$ on $[t_0, m_0]$. Therefore $(-1)^k L_k \varphi_0(t_1) \geq 0$. It implies that $(-1)^k L_k \varphi_0(t) \geq 0$ for all $t \geq t_0$, $k = 0, 1, \dots, n-1$. \square

Lemma 10. *Let $\{y_m(t)\}_{m=n_0}^\infty$ be a sequence of solutions of (L) on $[t_0, \infty)$, where $a < t_0 < n_0$, n is an odd number, and $L_k y_m(m) = (-1)^{k-1}$ for all $m \geq n_0$, $k = 0, 1, \dots, n-1$. Let (A) hold, and let $P_k(t) \equiv 0$ on $[a, \infty)$ for all even integers $k \in [1, n]$. Let $-\infty < \int_{t_0}^\infty P_0(s) ds = P < 0$, $\int_{t_0}^\infty P_k(s) ds \geq -\frac{1}{2}$ for $k = 1, 2, \dots, n-1$, let P_k be nondecreasing functions for $k = 0, 1, \dots, n-1$, $\int_{t_0}^\infty 1/p_r(s) ds \leq \frac{1}{2}$ for $r = 1, 2, \dots, n-1$, and let K be a real positive constant such that $0 \leq f(t) \leq K$ for $t \in (-\infty, \infty)$. Then there exists a subsequence of $\{y_m(t)\}_{m=n_0}^\infty$ which converges to*

$\varphi_0(t)$. This function $\varphi_0(t)$ is a solution of (L) on $[t_0, \infty)$, and $(-1)^k L_k \varphi_0(t) \leq 0$ on $[t_0, \infty)$ for $k = 0, 1, \dots, n-1$.

Proof. The proof is similar to the proof of Lemma 9 (instead of Lemma 6, part a), and Lemma 7 we use Lemma 6, part b) and Lemma 8, respectively), so it is omitted. \square

Theorem 1. Let n be an even number. Let (A) hold, and let $P_k(t) \equiv 0$ on $[a, \infty)$ for all odd integers $k \in [1, n]$. Let $P_k(t)$ be nondecreasing functions on $[a, \infty)$ such that $\int_a^\infty P_k(s) ds > -\infty$ for $k = 0, 1, \dots, n-1$, $\int_a^\infty 1/p_r(s) ds < \infty$ for $r = 1, 2, \dots, n-1$, and let K be a real positive constant such that $0 \leq f(t) \leq K$ for all $t \in (-\infty, \infty)$. Then (L) admits a Kneser solution $y(t)$ on $[a, \infty)$, i.e. $y(t) > 0$, $(-1)^k L_k y(t) \geq 0$ on $[a, \infty)$ for $k = 1, 2, \dots, n-1$.

Proof. Let us take $t_0 \in (a, \infty)$ such that $\int_{t_0}^\infty P_k(s) ds \geq -\frac{1}{2}$, $\int_{t_0}^\infty 1/p_r(s) ds \leq \frac{1}{2}$ for $k = 1, 2, \dots, n-1$; $r = 1, 2, \dots, n-1$. According to Lemma 5, there exists a sequence $\{y_m(t)\}_{m=n_0}^\infty$ of solutions of (L) on $[t_0, \infty)$ such that $L_k y_m(m) = (-1)^k$ for all $m \geq n_0 > t_0$, $k = 0, 1, \dots, n-1$. Lemma 7 ensures validity of (B), and Lemma 6, part a), yields that $\{y_m(t)\}_{m=n_0}^\infty$ has the required properties from Lemma 9. According to the last-mentioned lemma, there exists a function $y(t)$ such that $L(y(t)) \equiv 0$ on $[t_0, \infty)$, $(-1)^k L_k y(t) \geq 0$ on $[t_0, \infty)$ for $k = 0, 1, \dots, n-1$. This solution $y(t)$ of (L) on $[t_0, \infty)$ can be continued onto $[a, \infty)$ by Lemma 5. According to Lemma 6, part a), $y(t)$ is a Kneser solution of (L) on $[a, \infty)$ because $y(t) > 0$ on $[a, \infty)$ (this follows from $f(0) \neq 0$). \square

Theorem 2. Let n be an odd number. Let (A) hold, and let $P_k(t) \equiv 0$ on $[a, \infty)$ for all even integers $k \in [1, n]$. Let $P_k(t)$ be nondecreasing functions on $[a, \infty)$ such that $\int_a^\infty P_k(s) ds > -\infty$ for $k = 0, 1, \dots, n-1$, $\int_a^\infty 1/p_r(s) ds < \infty$ for $r = 1, 2, \dots, n-1$ and let K be a real positive constant such that $0 \leq f(t) \leq K$ for all $t \in (-\infty, \infty)$. Then (L) admits a Kneser solution $y(t)$ on $[a, \infty)$, i.e. $y(t) < 0$, $(-1)^k L_k y(t) \leq 0$ on $[a, \infty)$ for $k = 1, 2, \dots, n-1$.

Proof. The proof is similar to that of the previous theorem (instead of Lemma 6, part a) and Lemma 9 we will use Lemma 6, part b) and Lemma 10, respectively) and so it is omitted. \square

3. EXAMPLES

Example 1. The equation

$$(t^4(t^3(t^2y')')')' - \frac{1}{t^2}(t^3(t^2y')') + \left[\left(\frac{72}{t^8} - \frac{1296}{t^4}\right)\sqrt{1+t^{-18}}\right] \frac{1}{\sqrt{1+y^2}} \equiv 0$$

admits a Kneser solution $y(t) = t^{-9}$ on $[1, \infty)$ according to Theorem 1 because $\int_1^\infty (1/p_r(t)) dt < \infty$ for $r = 1, 2, 3$, $P_0(t)$ is nonpositive and nondecreasing on $[1, \infty)$, $\int_1^\infty P_k(t) dt > -\infty$ for $k = 0, 1, 2, 3$, $0 \leq 1/\sqrt{1+y^2} \leq 1$, $f(0) \neq 0$.

Example 2. The equation of the n -th order (n is an even number)

$$L_n y + \sum_{k=1}^{\frac{n}{2}-1} P_{2k}(t) L_{2k} y + P_0(t) f(y) \equiv 0,$$

where $P_{2k}(t) = -t^{-2k-2}$ for $k = 0, 1, \dots, \frac{n}{2} - 1$, $p_r(t) = t^{3r}$ for $r = 1, 2, \dots, n-1$, $f(t) = e^{-t^2}$ admits a Kneser solution on $[1, \infty)$ according to Theorem 1 because $\int_1^\infty (1/p_r(t)) dt < \infty$ for $r = 1, 2, \dots, n-1$, $\int_1^\infty P_{2k}(t) dt > -\infty$ for $k = 0, 1, \dots, \frac{n}{2} - 1$, $0 \leq e^{-t^2} \leq 1$, $f(0) \neq 0$.

Example 3. The equation

$$L_5 y - \frac{1}{t^6} L_3 y - \frac{1}{t^2} L_1 y + (12t^{-13} + 1188t^{-12} - 14256t^{-3}) \frac{\sqrt{1+t^{-48}}}{\sqrt{1+y^4}} \equiv 0,$$

where $p_r(t) = t^{r+1}$ for $r = 1, 2, 3, 4$ admits a Kneser solution $y(t) = -t^{-12} < 0$ on $[1, \infty)$ according to Theorem 2 because $\int_1^\infty (1/p_r(t)) dt < \infty$ for $r = 1, 2, 3, 4$, $P_0(t)$ is nonpositive and nondecreasing on $[1, \infty)$, $\int_1^\infty P_k(t) dt > -\infty$ for $k = 0, 1, 2, 3, 4$, $0 \leq \frac{1}{\sqrt{1+y^4}} \leq 1$, $f(0) \neq 0$.

References

- [1] *Barrett, J. H.*: Two-point boundary problems for linear self-adjoint differential equations of the fourth-order with middle term. *Duke Math. J.* 29 (1962), 543–554.
- [2] *Hartman, P.*: Ordinary Differential Equations. John Wiley & Sons, New York, 1964.
- [3] *Leighton, W.; Nehari, Z.*: On the oscillation of solutions of self-adjoint linear differential equations of the fourth order. *Trans. Amer. Math. Soc.* 89 (1958), 325–377.
- [4] *Palumbíny, O.*: On existence of monotone solutions of a certain class of n -th order nonlinear differential equations. To appear.
- [5] *Philos, Ch. G.*: Oscillation and asymptotic behaviour of third order linear differential equations. *Bull. Inst. Math. Acad. Sinica* 11(2) (1983), 141–160.
- [6] *Regenda, J.*: Oscillatory and nonoscillatory properties of solutions of the differential equation $y^{(4)} + P(t)y'' + Q(t)y = 0$. *Math. Slovaca* 28 (1978), 329–342.
- [7] *Rovder, J.*: Comparison theorems for third-order linear differential equations. *Bull. Inst. Math. Acad. Sinica* 19 (1991), 43–52.
- [8] *Rovder, J.*: Kneser problem for third order nonlinear differential equation. *Zborník vedeckých prác MtF STU Trnava* (1993).
- [9] *Shair, A.*: On the oscillation of solutions of a class of linear fourth order differential equations. *Pacific J. Math.* 34 (1970), 289–299.
- [10] *Škerlík, A.*: Criteria of property **A** for third order superlinear differential equations. *Math. Slovaca* 43 (1993), 171–183.
- [11] *Švec, M.*: On various properties of the solutions of third and fourth order linear differential equations. *Proceedings of the conference held in Prague in September 1962.* pp. 187–198.
- [12] *Švec, M.*: Über einige neue Eigenschaften der oszillatorischen Lösungen der linearen homogenen Differentialgleichung vierter Ordnung. *Czechoslovak Math. J.* 4(79) (1954), 75–94.
- [13] *Tóthová, M.; Palumbíny, O.*: On monotone solutions of the fourth order ordinary differential equations. *Czechoslovak Math. J.* 45(120) (1995), 737–746.

Author's address: Oleg Palumbíny, Department of Mathematics, Faculty of Material Engineering and Technology, Slovak Technical University, Paulínska 16, 917 24 Trnava, Slovakia, e-mail: palum@mtf.stuba.sk.