

A CONTRIBUTION TO BEST APPROXIMATION IN THE L_2 -NORM

S.R.H. Rizvi and O.P. Juneja

1. Let E denote a closed and bounded Jordan region in the complex plane with transfinite diameter $d > 0$. Let $w(z)$ be a positive continuous function on E and let $\mathcal{H}_2(E)$ denote the Hilbert space of functions analytic in D with inner product.

$$(1.1) \quad (f, g) = \iint_E w(z) f(z) \overline{g(z)} dx dy, \quad f, g \in \mathcal{H}_2(E)$$

so that for any $f \in \mathcal{H}_2(E)$, we have

$$(1.2) \quad \|f\| = \left[\iint_E w(z) |f(z)|^2 dx dy \right]^{1/2} < \infty$$

If $\mathcal{A}(E) \equiv \{p_{n-1}(z)\}_{n=1}^\infty$, $p_n(z)$ being a polynomial of degree n , is a complete orthonormal sequence in $\mathcal{H}_2(E)$ ¹, set, for any $f \in \mathcal{H}_2(E)$ and for $n = 0, 1, 2, \dots$

$$(1.3) \quad \Delta_n(f) \equiv \Delta_n(f; E) = \inf_{\{c_i\}} \left[\iint_E w(z) |f(z) - c_0 - c_1 p_1(z) - \dots - c_n p_n(z)|^2 dx dy \right]^{1/2}$$

$$(1.4) \quad a_n \equiv a_n(f; E) = \iint_E w(z) f(z) \overline{p_n(z)} dx dy.$$

$\Delta_n(f; E)$ is called the minimum error of f in L^2 -norm with respect to the system $\mathcal{A}(E)$ and a_n is called the n th Fourier coefficient of f with respect to the

¹Such a sequence of polynomials always exists in $\mathcal{H}_2(E)$ as can be easily seen with the help of Faber polynomials ([2] and [8]).

system $\mathcal{A}(E)$. It is well known (see e.g. [3] that if U denotes the unit disc and $|w(z)| \equiv 1$, then $\mathcal{A}(U) \equiv \left\{ \sqrt{\frac{2}{\pi}} z^{n-1} \right\}_{n=1}^{\infty}$ forms a complete orthonormal sequence in $\mathcal{H}_2(U)$ and that if $f(z) = \sum_{n=0}^{\infty} b_n z^n (|z| < 1)$ is in $\mathcal{H}_2(t)$ then

$$(1.5) \quad b_n = \sqrt{\frac{n+1}{\pi}} a_n(f; U).$$

Hence it follows from (1.5) that if f can be extended to an entire function of order ρ , lower order λ and type T then in all the results that give ρ, λ, T in terms of the coefficients b_n 's (see [1], [4], [5] etc.), one can replace b_n by $a_n(f; U)$. In fact, much more than this can be said. Authors have recently shown [7] that if $f \in \mathcal{H}_2(E)$, the Fourier series $\sum_{k=0}^{\infty} a_k p_k(z)$ converges uniformly to $f(z)$ on E and f can be extended to an entire function, if and only if,

$$(1.6) \quad \lim |a_n(f; E)|^{1/n} = 0.$$

Further, if f is of order $\rho (0 < \rho < \infty)$, type T and lower order λ then

$$(1.7) \quad \rho = \limsup_{n \rightarrow \infty} \frac{n \log n}{\log |a_n(f; E)|^{-1}};$$

$$(1.8) \quad Td^\rho = \limsup_{n \rightarrow \infty} (n/e_\rho) |a_n(f; E)|^{\rho/n};$$

$$(1.9) \quad \lambda = \sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log |a_{n_k}(f; E)|^{-1}};$$

where supremum in (1.9) is taken over all increasing sequences $\{n_k\}$ of positive integers.

Attempts have also been made to connect the minimum error $\Delta(f; U)$ with the growth of the entire function f . Thus, for $w(z) \equiv 1$, Reddy [6] has obtained relations essentially involving $\Delta_n(f; U)$ and the order and the type of entire function f . The aim of the present paper is to generalize the results of Reddy to any bounded and closed Jordan region E and for any weight function $w(z)$, positive and continuous on E . We also obtain relations involving $\Delta_n(f; E)$ and the lower order of f the analogues of which for the unit disc U have not been obtained by Reddy. We shall assume throughout that E is a bounded and closed Jordan region with transfinite diameter $d > 0$ and that for $f \in \mathcal{H}_2(E)$, $\Delta_n(f)$ and a_n are given by (1.3) and (1.4) respectively.

2. We first prove few lemmas which will be required in the sequel.

LEMMA 1: Let $f \in \mathcal{H}_2(E)$, then

$$(2.1) \quad \Delta_n(f) = \left[\sum_{k=1}^{\infty} |a_k|_2 \right]^{1/2}, \quad n = 1, 2, 3, \dots$$

PROOF. By (1.3), we have

$$\Delta_n(f) \leq \left[\iint_E w(z) \left| f(z) - \sum_{k=0}^n a_k p_k(z) \right|^2 dx dy \right]^{1/2}.$$

Since $\sum_{k=0}^{\infty} a_k p_k(z)$ converges to $f(z)$ on E , we get,

$$\begin{aligned} \Delta_n(f) &\leq \left[\iint_E w(z) \left| \sum_{k=n+1}^{\infty} a_k p_k(z) \right|^2 dx dy \right]^{1/2} \\ (2.2) \qquad &= \left[\sum_{k=n+1}^{\infty} |a_k|_2 \right]^{1/2}. \end{aligned}$$

Again as for any $(n + 1)$ complex numbers $\{a_0', a_1', \dots, a_n'\}$

$$\begin{aligned} &\iint_E w(z) |f(z) - a_0' p_0(z) - a_1' p_1(z) - \dots - a_n' p_n(z)|^2 dx dy \\ &= \iint_E \left| \sum_{k=0}^n (a_k - a_k') p_k(z) + \sum_{k=n+1}^{\infty} a_k p_k(z) \right|^2 dx dy \\ &= \sum_{k=0}^n |a_k - a_k'|^2 + \sum_{k=n+1}^{\infty} |a_k|^2 \\ &\geq \sum_{k=n+1}^{\infty} |a_k|^2. \end{aligned}$$

Since this is true for any $(n + 1)$ complex numbers a_0', a_1', \dots, a_n' , we have

$$(2.3) \qquad [\Delta_n(f)]^2 \geq \sum_{k=n+1}^{\infty} |a_k|^2.$$

Combining (2.2) and (2.3) we get (2.1).

LEMMA 2 [9]: Let $w = \varphi(z)$ be the function mapping the complement of E onto $|w| > 1$ such that $\varphi(\infty) = \infty$ and $\varphi'(\infty) > 0$ and let $E_r = \{z: |\varphi(z)| = r\}$. Let f be an entire function of order ρ ($0 < \rho < \infty$), lower order λ , type T and let

$$(2.4) \qquad \overline{M}(r) = \max_{z \in E_r} |f(z)|,$$

then

$$(2.5) \qquad \rho = \lim_{r \rightarrow \infty} \sup \frac{\log \log \overline{M}(r)}{\log r}$$

$$(2.6) \quad d^\rho T = \limsup_{r \rightarrow \infty} \frac{\log \overline{M}(r)}{r^\rho}$$

LEMMA 3: Let f be an entire function and $\overline{M}(r)$ be given by (2.4), then for any $\varepsilon > 0$

$$(2.7) \quad \Delta_n(f) \leq A \frac{\overline{M}(r)}{r^n} e^{n\varepsilon}$$

for $r \geq r_0(\varepsilon)$, and $n \geq n_0(\varepsilon)$, where A is a constant independent of n and r .

PROOF. We known [9] that there exist polynomials $\{g_n(z)\}_{n=1}^\infty$ of respective degree less than or equal to n , such that for all $z \in E$

$$(2.8) \quad f(z) - g_n(z) \leq K \frac{\overline{M}(r)}{r^n} e^{n\varepsilon}$$

for $n \geq n_0(\varepsilon, E)$ and $r \geq r_0(\varepsilon)$ and where K is a constant independent of n and depends only on ε and E .

Now as each $g_n(z)$ can be written as a linear combination of $\{p_k(z)\}_{k=0}^n$, it follows that

$$[\Delta_n(f)]^2 \leq \iint_E w(z) |f(z) - g_n(z)|^2 dx dy.$$

Using (2.8) we get

$$\Delta_n(f) \leq A \frac{\overline{M}(r)}{r^n} e^{n\varepsilon}$$

where A is a independent of n and r .

3. In this section we give necessary and sufficient conditions, in terms of $\Delta_n(f)$, for an entire function to be of order ρ ($0 < \rho < \infty$) and Type T . First we have

THEOREM 1. Let $f \in \mathcal{H}_2(E)$. Then f can be extended to an entire function if and only if,

$$(3.1) \quad \lim_{n \rightarrow \infty} \Delta_n(f)^{1/n} = 0.$$

PROOF. Suppose f can be extended to an entire function. Then, by (1.6), we have

$$\lim_{n \rightarrow \infty} |a_n|^{1/n} = 0;$$

so that for any ε satisfying $0 < \varepsilon < 1$, there exists an $n_0 = n_0(\varepsilon)$ such that

$$|a_n| < \varepsilon^n \text{ for } n \geq n_0.$$

Using (2.1), we get

$$\Delta_n(f) \leq \frac{\varepsilon^n}{(1 - \varepsilon^2)^{1/2}}.$$

So

$$\limsup_{n \rightarrow \infty} \Delta_n(f)^{1/n} \leq \varepsilon$$

i.e.

$$\limsup_{k \rightarrow \infty} \Delta_n(f)^{1/n} = 0.$$

Conversely, if $\lim_{n \rightarrow \infty} \Delta_n(f)^{1/n} = 0$, then as (2.1) gives

$$\Delta_n(f) > |a_{n+1}|$$

for every n , we get

$$\lim_{n \rightarrow \infty} |a_n|^{1/n} = 0.$$

Hence (1.6) yields that f is entire.

THEOREM 2. *Let $f \in \mathcal{H}_2(E)$; then f can be extended to an entire function of finite order ρ , if and only if,*

$$(3.2) \quad \mu \equiv \limsup_{n \rightarrow \infty} \frac{n \log n}{\log \Delta_n(f)^{-1}} < \infty.$$

Further $\mu = \rho$ also holds.

PROOF. Suppose first that (3.2) holds. Let ε be any positive number, then there exists $n_0 = n_0(\varepsilon)$ such that

$$\frac{n \log n}{\log \Delta_n(f)^{-1}} < \mu + \varepsilon \text{ for } n \geq n_0,$$

or

$$\Delta_n(f) < n^{-n/\mu + \varepsilon} \text{ for } n \geq n_0,$$

which gives $\lim_{n \rightarrow \infty} \Delta_n(f)^{1/n} = 0$.

So, in view of Theorem 1, f can be extended to an entire function. Let its order be ρ . Since, by (2.1)

$$(3.3) \quad \Delta_n(f) > |a_{n+1}|$$

for every n , (3.3) gives

$$\limsup_{n \rightarrow \infty} \frac{n \log n}{\log \Delta_n(f)^{-1}} \geq \limsup_{n \rightarrow \infty} \frac{n \log n}{\log |a_n|^{-1}}.$$

So, using (1.7) we get

$$\rho \leq \limsup_{n \rightarrow \infty} \frac{n \log n}{\log \Delta_n(f)^{-1}}$$

i.e.

$$(3.4) \quad \rho \leq \mu$$

which shows that f is of finite order.

Conversely, suppose that f is an entire function of finite order ρ . Then (1.7) gives that, for any $\varepsilon > 0$ there exists $n_0(\varepsilon)$ such that

$$|a_n| \leq n^{-n/\rho+\varepsilon} \text{ for } n \geq n_0.$$

Using (2.1) we get

$$[\Delta_n(f)]^2 \leq \sum_{k=n+1}^{\infty} k^{-2k/\rho+\varepsilon} \text{ for } n \geq n_0$$

or

$$[\Delta_n(f)]^2 \leq (n+1)^{-\frac{2(n+1)}{\rho+\varepsilon}} [1 + o(1)] \text{ as } n \rightarrow \infty.$$

i.e.

$$\limsup_{n \rightarrow \infty} \frac{n \log n}{\log \Delta_n(f)^{-1}} \leq (\rho + \varepsilon)$$

As this is true for every $\varepsilon > 0$, we get

$$(3.5) \quad \mu \leq \rho$$

which shows that (3.2) holds.

Combining (3.4) and (3.5) we also get $\mu = \rho$.

THEOREM 3. *Let $f \in \mathcal{H}_2(E)$ and v defined by*

$$(3.6) \quad \limsup_{n \rightarrow \infty} n |\Delta_n(f)|^{\rho/n} = v \quad (0 < \rho < \infty)$$

satisfies $0 < v < \infty$). Then f can be extended to an entire function of order ρ and type T , if and only if

$$v = (e\rho T)d^\rho,$$

where d is the transfinite diameter of E .

PROOF. Suppose $v = (e\rho T)d^\rho$ and $0 < v < \infty$. Then (3.6) gives easily

$$\limsup_{n \rightarrow \infty} \frac{n \log n}{\log \Delta_n(f)^{-1}} = \rho$$

So, Theorem 2 gives that f can be extended to an entire function of order ρ . Now, we shall show that its type is $v/(e\rho d^\rho)$.

Let f be of type T . By (2.6), we have, for any $\varepsilon > 0$, there exists $r_0 = r_0(\varepsilon)$ such that

$$(3.7) \quad \log \overline{M}(r) < (T' + \varepsilon)r^\rho \quad \text{for } r \geq r_0,$$

where

$$T' = Td^\rho.$$

Using (2.7) and (3.7), we get

$$\log \Delta_n(f) \geq \log A + n\varepsilon - n \log r + (T' + \varepsilon)r^\rho$$

for $n \geq n_0$ and $r \geq r_0$.

Chose a sequence $r_n \rightarrow \infty$ as

$$r_n = \left[\frac{n}{\rho(T' + \varepsilon)} \right]^{1/\rho},$$

then for large n , we have

$$\log \Delta_n(f) \leq \log A + n\varepsilon - \frac{n}{\rho} [\log n - \log(T' + \varepsilon)\rho] + \frac{n}{\rho}$$

or,

$$n\Delta_n(f)^{\rho/n} \leq e\rho(T' + \varepsilon)e^{\rho\varepsilon} + o(1).$$

Hence

$$\limsup_{n \rightarrow \infty} n\Delta_n(f)^{\rho/n} \leq e\rho(T' + \varepsilon)e^{\rho\varepsilon} \quad \text{as } n \rightarrow \infty$$

as this true for every $\varepsilon > 0$ so we get

$$v \equiv \limsup_{n \rightarrow \infty} n\Delta_n(f)^{\rho/n} \leq e\rho T' = e\rho Td^\rho.$$

i.e.,

$$(3.8) \quad T \geq v/(e\rho d^\rho).$$

For reverse inequality, we observe that, since

$$\Delta_n(f) \geq |a_{a+1}| \quad \text{for all } n,$$

so

$$\limsup_{n \rightarrow \infty} n|a_n|^{\rho/n} \leq \limsup_{n \rightarrow \infty} n\Delta_n(f)^{\rho/n}.$$

Using (1.8) we get

$$e\rho Td^\rho \leq \limsup_{n \rightarrow \infty} n\Delta_n(f)^{\rho/n} = v$$

i. e.,

$$(3.9) \quad T \leq v/(e\rho d^p).$$

Combing (3.8) and (3.9) we get $e\rho d^p T = v$.

This proves the theorem.

4. We now obtain relations involving $\Delta_n(f)$ and the lower order of f . We thus have

THEOREM 4. *Let $f \in \mathcal{H}_2(E)$. Then f can be extended to an entire function of finite lower order λ , and only if,*

$$\lambda = \sup_{\{n_k\}} \liminf_{n \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}} < \infty$$

where supremum is taken over all increasing sequences $\{n_k\}$ of positive integers.

PROOF. Suppose f can be extended to an entire function of finite lower order λ . We show that λ satisfies (4.1). Let $\{n_k\}$ be an increasing sequence of positive integers. Since (2.1) gives

$$\Delta_{n_k}(f) \geq |a_{n_k+1}|$$

for every k , we get

$$\liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}} \geq \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log |a_{n_k}|^{-1}}.$$

Hence

$$\sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}} \geq \sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log |a_{n_k}|^{-1}}.$$

Using (1.9) we get

$$\lambda \leq \sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}}.$$

Now for reverse inequality we proceed as follows. For any increasing sequence $\{n_k\}$ of positive integers define

$$(4.3) \quad \alpha \equiv \alpha(\{n_k\}) = \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}}.$$

Obviously $0 \leq \alpha \leq \infty$. Consider first the case when $0 < \alpha < \infty$. Let $\varepsilon > 0$ be such that $0 < \varepsilon < \alpha$, then, by (4.3), there exists $k_0 = k_0(\varepsilon)$ such that

$$\frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}} > \alpha - \varepsilon \text{ for } k \geq k_0.$$

Or

$$(4.4) \quad \Delta_{n_k}(f) \geq n_{k-1}^{-n_k/(\alpha-\varepsilon)} \text{ for } k \geq k_0.$$

Using (2.7) and (4.4), we get, for $r \geq r_0$ and $k \geq k_0$,

$$\log A + \log \overline{M}(r) > -\frac{n_k}{\alpha - \varepsilon} \log n_{k-1} + n_k \log r - n_k \varepsilon.$$

Define a sequence $r_k \rightarrow \infty$ as

$$(4.5) \quad r_k = e^{1+\varepsilon} n_{k-1}^{1/\alpha-\varepsilon}.$$

Then, if k is large and $r_k \leq r \leq r_{k+1}$,

$$\log A + \log \overline{M}(r) > -\frac{n_k}{\alpha - \varepsilon} \log n_{k-1} + n_k \log r_k - n_k \varepsilon.$$

Or,

$$\log A + \log \overline{M}(r) \geq \left(\frac{r_{n_k+1}}{e^{1+\varepsilon}} \right)^{\alpha-\varepsilon}$$

i.e.,

$$\log A + \log \overline{M}(r) \geq \left(\frac{r}{e^{1+\varepsilon}} \right)^{\alpha-\varepsilon}.$$

Hence

$$\lambda \equiv \liminf_{r \rightarrow \infty} \frac{\log \log \overline{M}(r)}{\log r} \geq \alpha - \varepsilon.$$

Since ε is arbitrary so we get $\lambda \geq \alpha$; which holds good in case $\alpha = 0$. Moreover, when $\alpha = \infty$, λ can not be finite. Hence

$$\lambda \geq \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}}.$$

Since, this is true for any increasing $\{n_k\}$ of positive integers, we get

$$(4.6) \quad \lambda \geq \sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}}.$$

Combing (4.2) and (4.6) we get (4.1)

Conversely, suppose (4.1) is true while f can not be extended to an entire function (3.1) gives

$$\limsup_{n \rightarrow \infty} \Delta_n(f)^{1/n} = l = 0$$

Since for every $f \in \mathcal{H}_2(E)$, $\Delta_n(f) \rightarrow 0$, it is clear that $0 < l \leq 1$. Let $\varepsilon > 0$ be any positive number such that $0 < \varepsilon < l$, then there exists an increasing sequence $\{n_k\}$ of positive integers such that

$$\Delta_{n_k}(f) > (l - \varepsilon)^{n_k} \text{ for every } k.$$

Now on this sequence $\{n_k\}$

$$\liminf_{k \rightarrow \infty} \frac{n_k \log n_{k-1}}{\log \Delta_{n_k}(f)^{-1}} = \infty,$$

which contradicts the fact that (4.1) holds. Hence f can be extended to an entire function and then its lower order λ is given by (4.1).

The following theorem can also be proved in a similar manner.

THEOREM 5. *Let $f \in \mathcal{H}_2(E)$. Then f can be extended to an entire function of finite lower order λ , if and only if,*

$$(4.7) \quad \lambda = \sup_{\{n_k\}} \liminf_{k \rightarrow \infty} \frac{(n_k - n_{k-1}) \log n_{k-1}}{\log \left| \frac{\Delta_{n_{k-1}}(f)}{\Delta_{n_k}(f)} \right|} < \infty,$$

where supremum is taken over all increasing sequences $\{n_k\}$ of positive integers.

5. In this section we study about the sequence $\{n_k\}$ of those values of n for which $\Delta_{n-1}(f) > \Delta_n(f)$ i.e.,

$$\Delta_n(f) = \Delta_{n_k}(f) \text{ for } n_{k-1} \leq n < n_k.$$

We shall see that for function of regular growth this sequence can not grow too rapidly. In fact we have

THEOREM 6. Let $f \in \mathcal{H}_2(E)$ be such that its extension is an entire function of order ρ and lower order λ ($0 \leq \lambda \leq \rho < \infty$); and let $\{n_k\}$ be the sequence defined by (5.1); then

$$\lambda \leq \liminf_{k \rightarrow \infty} \frac{\log n_k}{\log n_{k+1}}.$$

PROOF. Set $h(z) = \sum_{k=1}^{\infty} \pi_k(f) z^{n_k}$, where $\pi_k(f) = \Delta_{n_{k-1}}(f) - \Delta_{n_k}(f)$. In view of (1.7), (1.9) and theorem 2 and 4 it can be easily seen that $h(z)$ is an entire function of order ρ and lower order λ .

So

$$\rho = \limsup_{k \rightarrow \infty} \frac{n_k \log n_k}{\log \pi_k^{-1}(f)}$$

and

$$\begin{aligned} \lambda &= \sup_{\{m_h\}} \liminf_{h \rightarrow \infty} \frac{n_{m_h} \log n_{m_h-1}}{\log \pi_{m_h}^{-1}(f)} \\ &\leq \sup_{\{m_h\}} \limsup_{h \rightarrow \infty} \frac{n_{m_h} \log n_{m_h}}{\log \pi_{m_h}^{-1}(f)} \times \sup_{\{m_h\}} \liminf_{h \rightarrow \infty} \frac{\log n_{m_h-1}}{\log n_{m_h}} \\ &= \rho \liminf_{k \rightarrow \infty} \frac{\log n_k}{\log n_{k+1}} \end{aligned}$$

which completes the proof.

COROLLARY. *For entire functions or regular growth*

$$\log n_n \sim \log n_{k-1} \text{ as } k \rightarrow \infty$$

Finally we have

THEOREM 7. *Let f be an entire function. Then there exist integers*

$$0 \leq n_1 < n_2 < \cdots < n_k < \cdots$$

for which $a_{n_{k+1}} \neq 0$ for every k and

$$\Delta_{n_k} \sim |a_{n_{k+1}}| \text{ as } k \rightarrow \infty.$$

PROOF. Since f is entire, (1.6) gives

$$\lim_{k \rightarrow \infty} |a_k|^{1/k} = 0.$$

Let $\{\varepsilon_k\}$ be an arbitrary sequence of positive numbers each less than one and $\varepsilon \rightarrow \infty$ as $k \rightarrow \infty$. Now we shall show that there exists a sequence $\{n_k\}$ such that

$$|a_{n_k+j}| \leq |a_{n_k+1}| \varepsilon_k^{j-1} \text{ for } k = 1, 2, \dots$$

Since for each $\varepsilon_k > 0$, there exists $N = N(k)$ such that

$$(5.4) \quad |a_n|^{1/n} \leq \varepsilon_k \text{ for } n \geq N(k);$$

let $\{n_k\}$ be the sequence such that

$$\max_{n \geq N=N(k)} |a_n|^{1/n} = |a_{n_k+1}|^{1/(n_k+1)},$$

then (5.4) gives

$$|a_{n_k+1}|^{1/(n_k+1)} < \varepsilon_k$$

for every k ; so

$$|a_{n_k+j}|^{1/(n_k+j)} \leq |a_{n_k+1}|^{1/(n_k+1)} < \varepsilon_k$$

for every j or

$$\begin{aligned} |a_{n_k+j}| &< |a_{n_k+1}|^{\frac{n_k+j}{n_k+1}} \varepsilon_k^{n_k+j} \\ &\leq |a_{n_k+1}| |a_{n_k+1}|^{\frac{j-1}{n_k+1}} \end{aligned}$$

Or

$$|a_{n_k+j}| < |a_{n_k+1}| \varepsilon_k^{j-1}.$$

Now as

$$(\Delta_n(f))^2 = \sum_{n=1}^{\infty} |a_n|^2.$$

So we get

$$(\Delta_{n_k}(f))^2 < \sum_{j=1}^{\infty} |a_{n_k+1}|^2 \varepsilon_k^{(j-1)/2}$$

or

$$(5.5) \quad \Delta_{n_k}(f) < \frac{|a_{n_k+1}|}{(1 - \varepsilon_k^2)^{1/2}}.$$

Combining (5.5) together with the fact that $\Delta_n(f) > |a_{n+1}|$ we get that

$$\lim_{k \rightarrow \infty} \frac{\Delta_{n_k}(f)}{|a_{n_k+1}|} = 1$$

i.e $\Delta_{n_k} \sim |a_{n_k+1}|$ as $k \rightarrow \infty$.

REFERENCES

- [1] Boas, R.P., *Entire functions*, Academic Press, New York, 1954.
- [2] Epstein, B., *Orthogonal families of analytic functions*, The Macmillan Co., New York, 1965.
- [3] Goffman C. and Pedrick, G. *First course in Functional Analysis*, Prentice Hall of India, New Delhi, 1974.
- [4] Juneja, O.P., *On the coefficients of an entire series*, J. Analyse Math., **24**, 395–401 (1971).
- [5] Juneja, O.P. and Kapoor, G.P., *On the lower order of entire functions*, J. London Math. Soc. (2), **5**, 310–312 (1972).
- [6] Reddy, A.R., *A contribution to best approximation in the L^2 norm*, J. Approximation Theory, **11**, 110–117 (1974).
- [7] Rizvi, S.R.H. and Juneja, O.P., *Fourier expansions of entire functions*, To appear,
- [8] Smirnov, V.I., and Lebedev, N.A., *Function of a complex variable*, Iliffe Books London, 1968.
- [9] Winiarski, T.N., *Approximation and Interpolation of entire function*, Annales Polonici Mathematici, **23**, 259–273 (1970).

Department of Mathematics
 Indian Institute of Technology
 Kanpur 208016,
 INDIA