

INTEGRABILITY OF TENSOR STRUCTURES OF ELECTROMAGNETIC TYPE

J. M. Hernando, E. Reyes, P. M. Gadea

Abstract. We study characterizations of the integrability of G -structures defined by tensor fields of electromagnetic type.

1. Introduction. In [3] were considered the G -structures defined by a (1,1) tensor \tilde{J} on a differentiable manifold M^n such that

$$(\tilde{J}^2 - f^2)(\tilde{J}^2 + g^2) = 0,$$

where f, g are C^∞ functions on M^n nowhere zero. This situation generalizes that of Hlavaty [4] and Mishra [7]. They consider the so called electromagnetic tensor fields (of first class) on a 4-manifold which is the space-time of General Relativity. In [3] it was proved that the G -structure P defined by such a tensor field \tilde{J} is identical to the G -structure defined by a (1, 1) tensor field J that satisfies the same conditions as \tilde{J} but with $f = g = 1$, and so we have $J^4 = 1$.

On the other hand, that situation generalizes also the almost product and almost complex structures simultaneously. In [9] the family of linear connections that parallelize J (and an adapted metric also) is given. Connections partially adapted to such a structure are studied in [11]. In this note we study several characterizations and conditions of integrability of the (G -structure defined by the) tensor field \tilde{J} .

Thus, we consider the following situation:

Let M^n be a differentiable manifold and \tilde{J} a (1,1) tensor field such that:

a) $(\tilde{J}^2 - f^2)(\tilde{J}^2 + g^2) = 0$, where f, g are C^∞ functions on M^n with f, g nowhere null;

b) The characteristic polynomial of \tilde{J} is $(x - f)^{r_1}(x + f)^{r_2}(x^2 + g^2)^s$, where r_1, r_2, s are constants greater than or equal to 1 such that $r_1 + r_2 + 2s = n$. Then,

since J which satisfies a) and b), but with $f = g = 1$, defines the same G -structure P as \tilde{J} (not an associated G -structure, but exactly the same P see [3]), we can characterize the integrability of P in terms of J .

2. Integrability in terms of the Nijenhuis tensor. We denote from now on by X, Y, \dots , vectors fields on M^n . We consider the complementary projection operators $l = (J^2 + 1)/2$, $l_3 = (1 - J^2)/2$, which verify

$$Jl = lJ; J^2l = l, Jl_3 = l_3J, J^2l_3 = -l_3;$$

denote by L and L_3 the corresponding distributions, and put $L = L_1 \oplus L_2$, where L_1 and L_2 are distributions corresponding to the projectors l_1 and l_2 on L given by the eigenvalues $+1$ and -1 of $J|_L$. Let us decompose the Nijenhuis tensor of in the following manner:

$$\begin{aligned} N(X, Y) &= lN(lX, lY) + l_3N(lX, lY) + N(lX, l_3Y) \\ &\quad + N(l_3X, lY) + lN(l_3X, l_3Y) + l_3N(l_3X, l_3Y). \end{aligned}$$

Then we have the following

PROPOSITION 2.1. a) L is integrable iff $(\forall X, Y)l_3N(lX, lY) = 0$;

b) L_3 is integrable iff $(\forall X, Y)lN(l_3X, l_3Y) = 0$;

c) If L is integrable, the almost product structure defined by $J|_L$ on each integral manifold of L is integrable iff $(\forall X, Y)N(lX, lY) = 0$;

d) If L_3 is integrable, the almost complex structure defined by $J|_{L_3}$ on each integral manifold of L_3 is integrable iff $(\forall X, Y)N(l_3X, l_3Y) = 0$.

Proof. a) $N(lX, lY) = [JlX, JlY] - J[JlX, lY] - J[lX, JlY] + J^2[lX, lY]$. Thus, if L is integrable, each bracket is an element of L and so $l_3N(lX, lY) = 0$. Conversely, suppose now that $l_3N(lX, lY) = 0$; then we obtain easily:

$$\begin{aligned} &l_3N(JlX, JlY) + Jl_3N(JlX, lY) + Jl_3N(lX, JlY) \\ &= 3l_3[lX, lY] + l_3(N(lX, lY) - J^2[lX, lY]) \\ &= 4l_3[lX, lY] + l_3N(lX, lY), \end{aligned}$$

and since by the hypothesis $l_3N(lX, lY) = 0$, L is integrable;

b) Analogous to a), if we consider now

$$lN(Jl_3X, Jl_3Y) + JlN(Jl_3X, l_3Y) + JlN(l_3X, Jl_3Y);$$

c) If L is integrable, then $J|_L$ induces on each integral manifold of L an almost product structure. As such a structure is integrable iff its Nijenhuis tensor is zero, that is, $N_{J|_L}(lX, lY) = 0$, and since $N_{J|_L}(lX, lY) = N(lX, lY)$, we obtain c);

d) Similar to c). □

Definition 2.2. We say that J is *partially integrable* iff L and L_3 are integrable, and also the almost product and almost complex structure induced by J on the integral manifolds of L and L_3 , respectively.

Thus J is partially integrable iff $N(X, Y) = N(lX, l_3Y) + N(l_3X, lY)$.

So, we consider now the condition $N(lX, l_3Y) = 0$. Since the Lie derivative $L_Y J$ verifies by definition $(L_Y J)X = J[X, Y] - [JX, Y]$, we deduce:

$$a) N(lX, l_3Y) = J(L_{l_3Y} J)lX - (L_{Jl_3Y} J)lX;$$

$$b) N(l_3X, lY) = J(L_{lY} J)l_3X - (L_{JlY} J)l_3X;$$

and from these expressions it is immediate that:

PROPOSITION 2.3. $lN(lX, l_3Y) = 0$ (resp. $l_3N(lX, l_3Y) = 0$) for every X, Y iff $l(L_{l_3Z} J)l = 0$ (resp. $l_3(L_{lZ} J)l_3 = 0$ for every Z).

COROLLARY 2.4. $N(lX, l_3Y) = 0$ iff $l(L_{l_3Z} J)l = l_3(l_{lZ} J)l_3 = 0$, for every X, Y, Z .

Now, we have

THEOREM 2.5. J is integrable iff $N_J = 0$.

Proof. J is integrable iff for every $x \in M^n$, there exists a neighbourhood U of x and a coordinate system in $U, \{x^i\}$, such that the basis $\{\partial/\partial x^i\}, i = 1, \dots, n$ is adapted in U . That is, J can be expressed as a linear combination of products $\partial/\partial x^i \otimes dx^j$ with constant coefficients, and so, trivially, $N = 0$.

Conversely, suppose $N = 0$. By a) and b) of Prop. 2.1, L and L_3 are integrable. Thus, for each $x \in M^n$ there exists a chart centered at $x, (U, \varphi)$, with coordinates $\{x^i, y^a\}, i = 1, \dots, r_1 + r_2, a = 1, \dots, 2s$, such that

$$\partial/\partial x^i \in L, \quad \partial/\partial y^a \in L_3.$$

So, in the local basis $\{\partial/\partial x^i, \partial/\partial y^a\}$, J has a matrix of the form

$$J = \begin{pmatrix} J_j^i & 0 \\ 0 & J_b^a \end{pmatrix};$$

that is, $J = J_j^i \partial/\partial x^i \otimes dx^j + J_b^a \partial/\partial y^a \otimes dy^b$. Moreover, $l = \partial/\partial x^i \otimes dx^i$, and $l_3 = \partial/\partial y^a \otimes dy^a$. Thus,

$$L_{\partial/\partial y^c} J = \frac{\partial J_j^i}{\partial y^c} \frac{\partial}{\partial x^i} \otimes dx^j + \frac{\partial J_b^a}{\partial y^c} \frac{\partial}{\partial y^a} \otimes dy^b.$$

Hence

$$l(L_{\partial/\partial y^c} J)l = \frac{\partial J_j^i}{\partial y^c} \frac{\partial}{\partial x^i} \otimes dx^j.$$

So, Corollary 2.4 implies $\partial J_j^i / \partial y^c = 0$ (and analogously $\partial J_b^a / \partial x^i = 0$).

Consider now the integral manifold L_x of L , of coordinates $y^a = 0$. Since the almost product structure on L_x is integrable, there exist coordinate functions u^i in a neighbourhood of $x \in L_x$ in L_x such that

$$\partial/\partial u^i \in L_1, \quad i = 1, \dots, r_1, \quad \partial/\partial u^i \in L_2, \quad i = r_1 + 1, \dots, r_1 + r_2,$$

where these fields are considered in the regular submanifold $L_x \cap U$.

We define new coordinates in a neighbourhood W of x in U : put, for $x' \in W$,

$$u^i(x') = u^i(\varphi^{-1}(x^1(x'), \dots, x^{r_1+r_2}(x')), 0, \dots, 0), \quad \bar{y}^a(x') = y^a(x).$$

Then, for the new coordinate system $\{u^i, y^{-a}\}$ we have

$$\frac{\partial}{\partial x^i} = \frac{\partial u^j}{\partial x^i} \frac{\partial}{\partial u^j}, \quad dx^i = \frac{\partial x^i}{\partial u^j} du^j, \quad \text{where} \quad \frac{\partial}{\partial \bar{y}^a} \left(\frac{\partial u^i}{\partial x^i} \right) = \frac{\partial}{\partial \bar{y}^a} \left(\frac{\partial x^i}{\partial u^j} \right) = 0$$

This is a new coordinate system adapted to L and L_3 , and we have

$$J = \bar{J}_j^i \frac{\partial}{\partial u^i} \otimes du^j + \bar{J}_b^a \frac{\partial}{\partial \bar{y}^a} \otimes d\bar{y}^b, \quad \text{with} \quad \frac{\partial \bar{J}_j^i}{\partial \bar{y}^a} = 0.$$

But for the points of coordinates $\bar{y}^a = 0$ we have by construction (see 2.1)

$$\bar{J}_j^i = \begin{pmatrix} I_{r_1} & 0 \\ 0 & -I_{r_2} \end{pmatrix}.$$

Hence, in certain neighbourhood of x we also have the same matrix expression for \bar{J}_j^i .

Similarly, since the structure defined by J in L_{3x} is almost complex, a change of coordinates analogous to the previous one gives for \bar{J}_b^a the expression

$$\begin{pmatrix} 0 & -I_s \\ I_s & 0 \end{pmatrix}.$$

In other words, we have deduced that J is integrable. \square

3. Integrability in terms of a linear connection. Now, let ∇ be a linear connection without torsion on M^b and let Q be the $(1, 2)$ tensor field on M^n defined by $Q(X, Y) = \{(\nabla_{JY}J)X + J(\nabla_YJ)X + 2J(\nabla_XJ)Y\}/4$.

We define a new connection D by means of the expression

$$D_X Y = l\nabla_X lX - l\nabla_{lY} l_3 X + l_3 \nabla_X l_3 Y - l_3 \nabla_{l_3 Y} lX + lQ(lX, lY) - l_3 Q(l_3 X, l_3 Y).$$

It is easily proved that:

$$\begin{aligned} i) \quad & D_X lY = l\nabla_X lY - l\nabla_{lX} l_3 X + lQ(lX, lY); \\ ii) \quad & D_X l_3 Y = l_3 \nabla_X l_3 Y - l_3 \nabla_{l_3 Y} lX - l_3 Q(l_3 X, l_3 Y); \\ iii) \quad & D_{lX} lY = l\nabla_{lX} lY + lQ(lX, lY); \\ iv) \quad & D_{l_3 X} lY = l\nabla_{l_3 X} lY - l\nabla_{lY} l_3 X; \\ v) \quad & D_{lX} l_3 Y = l_3 \nabla_{lX} l_3 Y - l_3 \nabla_{l_3 Y} lX; \\ vi) \quad & D_{l_3 X} l_3 X = l_3 \nabla_{l_3 X} l_3 Y - l_3 Q(l_3 X, l_3 Y); \\ vii) \quad & D_X l = D_X l_3 = 0. \end{aligned} \tag{3.1}$$

So we have

PROPOSITION 3.1. *The torsion T of D has the expression*

$$T(X, Y) = 1/4\{lN(lY, lX) + l_3N(l_3N(l_3X, l_3Y))\} - l[l_3X, l_3Y] - l_3[lX, lY].$$

Proof. Immediate from the expression for D and Q , applying that ∇ is torsionless, and proving that $Q(X, Y) - Q(Y, X) = N(Y, X)/4$ \square

From that we obtain

COROLLARY 3.2. $(\forall X, Y)T(lX, l_3Y) = 0$.

We now prove

LEMMA 3.3. a) *The distribution L is integrable iff $l_3T(lX, lY) = 0$;*

b) *The distribution L_3 is integrable iff $lT(l_3X, l_3Y) = 0$;*

c) *If L is integrable, then the almost product structure induced by $J|_L$ on each integral manifold of L is integrable iff $lT(lX, lY) = 0$;*

d) *If L_3 is integrable, then the almost complex structure induced by $J|_{L_3}$ on each integral manifold of L_3 is integrable iff $l_3T(l_3X, l_3Y) = 0$;*

e) *J is partially integrable iff $T(X, Y) = 0$;*

f) $lN(JlX, lY) = lN(lX, JlY)$; g) $lT(JX, lY) = lT(lX, JY)$;

h) $l_3N(Jl_3X, l_3Y) = l_3N(l_3X, Jl_3Y)$; i) $l_3T(JX, l_3Y) = l_3T(l_3X, JY)$;

j) $(D_{l_3X}J)l_3Y = 0$; k) $(D_{lX}J)lY = 0$.

Proof. a) It suffices to prove $l_3T(lX, lY) = -l_3[lX, lY]$;

b) analogous to a); c) it suffices to consider $lT(lX, lY) = lN(lY, lX)/4$ and a), c) of Prop. 2.1; d) it is deduced from $l_3T(l_3X, l_3Y) = l_3N(l_3X, l_3Y)/4$, and b), d) of Prop. 2.1.;

e) from Cor. 3.2 we obtain $T(X, Y) = T(lX, lY) + T(l_3X, l_3Y)$. If J is partially integrable, from a) and c) we deduce $T(lX, lY) = 0$ and from b) and d) that $T(l_3X, l_3Y) = 0$. Hence $T(X, Y) = 0$. Conversely, $T(X, Y) = 0$ implies $T(lX, lY) = T(l_3X, l_3Y) = 0$ and thus $lT(lX, lY) = l_3T(lX, lY) = lT(l_3X, l_3Y) = l_3T(l_3X, l_3Y) = 0$.

From these equalities and from a), b), c), d) we deduce that J is partially integrable; f) the proof is immediate and moreover, as a consequence, we obtain $lN(JlX, JlY) = lN(lX, lY)$; g) $lT(JX, lY) = lN(lY, JlX)/4$ and $lT(JX, lX) = lN(JlY, lX)/4$, and from f) we obtain the result; h) the proof is analogous to that of f) and we deduce here $l_3N(Jl_3X, Jl_3Y) = -l_3N(l_3X, l_3Y)$;

i) we have $l_3T(JX, l_3Y) = l_3N(Jl_3X, l_3Y)/4$ and $l_3T(l_3X, JY) = l_3N(l_3X, Jl_3Y)/4$, and the conclusion follows from h);

j) from (3.1), vi) we have $D_{l_3X}Jl_3Y = l_3\nabla_{l_3X}Jl_3Y - l_3Q(l_3X, Jl_3Y)$ and $Jl_3D_{l_3X}l_3Y = Jl_3\nabla_{l_3X}l_3Y - l_3JQ(l_3X, l_3Y)$. Subtracting we obtain

$$\begin{aligned} (D_{l_3X}J)l_3Y &= l_3(\nabla_{l_3X}J)l_3Y + l_3\{J(\nabla_{Jl_3Y}J)l_3X + J^2(\nabla_{l_3X}J)l_3X \\ &\quad + 2J^2(\nabla_{l_3X}J)l_3Y - (\nabla_{J^2l_3Y}J)l_3X - J(\nabla_{Jl_3X}J)l_3X \\ &\quad - 2J(\nabla_{l_3X}J)l_3Y\}/4 = l_3(\nabla_{l_3X}J)l_3Y - l_3(\nabla_{l_3X}J)l_3Y = 0; \end{aligned}$$

k) analogous to that of j), by using (3.1), iii). \square

THEOREM 3.4. *J is integrable iff there exists a linear connection without torsion that parallelizes J . If J is integrable, then D gives an explicit example of such a connection.*

Proof. Suppose J integrable. Then for the earlier connection D we have: 1) D is torsionless, and 2) $DJ = 0$. Indeed, if J is integrable then it is partially integrable and, from e) of Lemma 3.3. we obtain 1). On the other hand, if J is integrable, $N(X, Y) = 0$, and since from (3.1), v) we have

$$(D_{JlX}J)l_3Y + (D_{lX}J)Jl_3Y = l_3N(lX, l_3Y) = 0,$$

we deduce

$$(D_{JlX}J)l_3Y = -(D_{lX}J)Jl_3Y. \quad (3.2)$$

Substituting Y by JY we have

$$(D_{JlX}J)l_3JY = -(D_{lX}J)J^2l_3Y = (D_{lX}J)l_3Y, \quad (3.3)$$

and, if in (3.2) we substitute X by JX , we have

$$(D_{JlX}J)Jl_3Y = -(D_{J^2lX}J)l_3Y = -(D_lJ)l_3Y. \quad (3.4)$$

From (3.3) and (3.4) we deduce

$$(D_{lX}J)l_3Y = 0. \quad (3.5)$$

From (3.1), iv) we have $lN(l_3X, lY) = (D_{Jl_3X}J)lY + (D_{l_3X}J)JlY$; if J is integrable we obtain analogously

$$(D_{l_3X}J)lY = 0. \quad (3.6)$$

But in j) and k) of Lemma 3.3. we have

$$(D_{l_3X}J)l_3Y = 0 \text{ and } (D_{lX}J)lY = 0 \quad (3.7)$$

Hence, from (3.5) (3.6) and (3.7) follows 2).

Conversly, suppose now that there is a linear connection ∇ without torsion such that $\nabla J = 0$. If we consider Q from ∇ as before, we see that $Q = 0$. But in the proof of the Prop. 3.1 we have seen that $Q(X, Y) - Q(Y, X) = N(Y, X)/4$; that is, J is integrable.

Remark. As is well known, Lehmann-Lejeune [5] proves that, for 0-deformable (1,1) tensor fields, the integrability is equivalent to the existence of a torsionless structural local connection. In our case, we have a global connection and we also give its explicit expression when J is integrable.

4. Integrability in terms of the structure tensor. We have now at disposal two criteria of integrability of the G -structure P defined by \tilde{J} . The first

one in terms of the Nijenhuis tensor of the field J , the second one in terms of a linear connection. A third criterion is that which expresses the integrability in terms of the Guillemin structure tensors [2].

The field \tilde{J} is not 0-deformable, but the associated field J , which defines the same structure P , is 0-deformable and so, we can anew characterize the integrability of P in terms of J ; but from the results of Lehmann-Lejeune [5] it suffices to consider, in this case, the Chern-Ehresmann-Bernard tensor [1] and have the equivalence of the integrability with nullity of the 1-st structure tensor of P , as we express in the final theorem.

5. Integrability in terms of prolongations and complete lifts. Now, we consider of the one hand the complete lift J^c of J in the sense of Yano-Kobayashi [12], which is a (1,1) tensor field on TM^n defined from J and, on the other hand, the canonical prolongation \hat{J} of J in the sense of Morimoto [8], which is also a (1,1) tensor field on TM^n . Firstly, we have the following:

PROPOSITION 5.1. *The canonical prolongation to TN^n of the $(J^4 = 1)$ -structure J is a $(J^4 = 1)$ -structure \hat{J} which coincides with the complete lift J^c of J .*

Proof. The structural group G corresponding to J (and \tilde{J}) is that of matrices of the form [3]

$$\left[\begin{array}{cc|c} A & 0 & 0 \\ 0 & B & \\ \hline 0 & C & -D \\ & D & C \end{array} \right]$$

where $A \in Gl(r_1, \mathbf{R})$, $B \in Gl(r_2, \mathbf{R})$, $C + iD \in Gl(s, \mathbf{C})$, $r_1 + r_2 + 2s = n$. If we denote \hat{G} the structural group of the canonical prolongation \hat{P} of P defined by J (and \tilde{J}), it has as elements the \hat{g} obtained by means of

$$\hat{g} = j_n(\{g, X\}), \quad g \in G, \quad X \in \mathfrak{g},$$

where \mathfrak{g} denotes the Lie algebra of G , X the translated $R_{g^{-1}*}Y$, for a certain $Y \in T_g G$, and j_n the imbedding $j_n : TGl(n, \mathbf{R}) \rightarrow Gl(2n, \mathbf{R})$.

More precisely,

$$j_n(\{g, X\}) = \begin{bmatrix} g & 0 \\ Xg & g \end{bmatrix} = \left[\begin{array}{cc|cc|cc} A & 0 & & & & \\ 0 & B & & 0 & & \\ \hline & & C & -D & & 0 \\ 0 & & D & C & & \\ \hline \alpha & 0 & & & A & 0 \\ 0 & \beta & & 0 & 0 & B \\ \hline & & \gamma & -\delta & & \\ 0 & & \delta & \gamma & 0 & \\ \hline & & & & C & -D \\ & & & & D & C \end{array} \right] \quad (5.1)$$

since

$$Xg = \left[\begin{array}{cc|cc} M & 0 & & \\ 0 & N & & 0 \\ \hline & & P & -Q \\ 0 & & Q & P \end{array} \right] \left[\begin{array}{cc|cc} A & 0 & & \\ 0 & B & & 0 \\ \hline & & C & -D \\ 0 & & D & C \end{array} \right] = \left[\begin{array}{cc|cc} MA & 0 & & \\ 0 & NB & & 0 \\ \hline & & PC-QD & -PD-QC \\ 0 & & QC+PD & -QD+PC \end{array} \right]$$

where $M \in \mathcal{M}(r_1; \mathbf{R})$, $N \in \mathcal{M}(r_2; \mathbf{R})$, $P + iQ \in \mathcal{M}(s, \mathbf{C})$.

It is immediate that $j_n(\{g, X\})$, belongs to the matrix group $Gl(2r_1, \mathbf{R}) \times Gl(2r_2, \mathbf{R}) \times Gl(2s, \mathbf{C})$, by means of a convenient rearrangement of the boxes of the matrix (5.1). Hence, we have the structural group of a ($J^4 = 1$)-structure on TM^n .

On the other hand, for a given local coordinate system $\{U, x^1, \dots, x^n\}$ on M^n , and a section σ of the principal bundle of frames FM^n on U , expressed as


$$\sigma(x) = \left(\dots, \sum_{i=1}^n \sigma_j^i(x) \frac{\partial}{\partial x^i} \Big|_x, \dots \right), \quad x \in U,$$

Morimoto [8] proves that $\tilde{\sigma} = j_{M^n} \cdot T\sigma$ (where j_{M^n} is the canonical embedding $j_{M^n} : TFM^n \rightarrow FTM^n$), is a section of FTM^n on TU , which can be expressed as

$$\tilde{\sigma} \left(\sum_{i=1}^n y^i \frac{\partial}{\partial x^i} \Big|_x \right) = \left(\dots, \sum_{i=1}^n \sigma_j^i(x) \frac{\partial}{\partial x^i} \Big|_x + \sum_{i,k=1}^n \frac{\partial \sigma^i(x)}{\partial x^k} y^k \frac{\partial}{\partial y^i} \Big|_X \dots, \right. \\ \left. \sum_{i=1}^n \sigma_j^i(x) \frac{\partial}{\partial y^i} \Big|_X, \dots \right)$$

where $\{x^1, \dots, x^n, y^1, \dots, y^n\}$ is the local coordinate system induced in TU , and $X = \sum_{i=1}^n y^i \frac{\partial}{\partial x^i} \Big|_x \in TU$.

We now consider the diagram



with the explained notations. Let again $\{U, x^1, \dots, x^n\}$ a local coordinate system on M^n , and σ a section of the G -structure P on U ; then $\tilde{\sigma} = j_{M^n} \cdot T\sigma$ is a section of the canonical prolongation \hat{P} of P , since

$$\hat{\sigma}(TU) = j_{M^n} \cdot T\sigma(TU) \subset j_{M^n}(T(\sigma(U))) \subset j_{M^n}(TP) = \hat{P}.$$

Now, let $J_0 : \mathbf{R}^n \rightarrow \mathbf{R}^n$ be an automorphism such that $J_0^4 = 1$.

From the diagram

$$\begin{array}{ccc} T_x M^n & \xrightarrow{J_x} & T_x M^n \\ \sigma(x) \uparrow & & \uparrow \sigma(x) \\ \mathbf{R}^n & \xrightarrow{J_0} & \mathbf{R}^n \end{array}$$

we define J_x as $J_x = \sigma(x) \cdot J_0 \cdot \sigma(x)^{-1}$.

Then $J : x \sim \rightarrow J_x$ is the $(J^4 = 1)$ -structure associated to P globally defined.

Indeed:

a) $J^4 = 1$. Immediate from $J_0^4 = 1$;

b) globally defined: If $x \in U \cap U'$, where U, U' are coordinate neighbourhoods and σ' is a section of P on U' , then $J'_x = \sigma'(x) \cdot J_0 \cdot \sigma'(x)^{-1}$, but since

$$\sigma'(x) = g(x) \cdot \sigma(x), \quad g(x) \in G.$$

we deduce

$$J'_x = g(x) \cdot \sigma(x) \cdot J_0 \cdot \sigma(x)^{-1} \cdot g(x)^{-1} = g(x) \cdot J_x \cdot g(x)^{-1}.$$

We now consider $TJ_0 : T\mathbf{R}^n \rightarrow T\mathbf{R}^n$. Since

$$TJ_0 = \begin{bmatrix} J_0 & 0 \\ 0 & J_0 \end{bmatrix}$$

we have $(TJ_0)^4 = 1$, and we define

$$\hat{J}(X) = \hat{\sigma}(X) \cdot TJ_0 \cdot \hat{\sigma}(X)^{-1}, \quad \text{for every } X \in TU.$$

It is clear that $\hat{J}^4 = 1$, and \hat{J} is the $(J^4 = 1)$ -structure on TM^n canonical prolongation of J , since $\hat{\sigma} = j_{M^n} \cdot T\sigma$.

On the other hand, we can choose as a basis of $T_X TM^n$ the set

$$\left\{ \frac{\partial}{\partial x^1} \Big|_X, \dots, \frac{\partial}{\partial x^n} \Big|_X, \frac{\partial}{\partial y^1} \Big|_X, \dots, \frac{\partial}{\partial y^n} \Big|_X \right\}.$$

Then, using the earlier expressions for $\hat{\sigma}$ and TJ_0 , we obtain

$$\begin{aligned} \hat{J}(x) = \hat{\sigma}(x) \cdot TJ_0 \cdot \hat{\sigma}(x)^{-1} &= \begin{bmatrix} \sigma(x) & 0 \\ \partial\sigma(x) & \sigma(x) \end{bmatrix} \begin{bmatrix} J_0 & 0 \\ 0 & J_0 \end{bmatrix} \begin{bmatrix} \sigma(x) & 0 \\ \partial\sigma(x) & \sigma(x) \end{bmatrix}^{-1} \\ &= \begin{bmatrix} J_x & 0 \\ \partial J_x & J_x \end{bmatrix}, \end{aligned}$$

which is precisely the formula of the complete lift J^c of J (see [12]), being

$$\partial J_x = \left(\sum_{k=1}^n y^k \frac{\partial J_j^i}{\partial x^k} \right)$$

□

But Morimoto [8] proves that a G -structure P is integrable if and only if the canonical prolongation \hat{P} is integrable; hence we obtain.

PROPOSITION 5.2. *Let (M^n, J) be a $(J^4 = 1)$ -manifold. Then the following statements are equivalent:*

- a) *The G -structure P defined by J is integrable;*
- b) *The Nijenhuis tensor of the tensor \hat{J} corresponding to the canonical prolongation \hat{P} of P is zero;*
- c) *The Nijenhuis tensor of the complete lift J^c of J is zero.*

6. J -Lie groups. Now, we consider a sufficient condition in order to a $(j^4 = 1)$ -structure be integrable.

Let (M_1, J_1) and (M_2, J_2) be two $(J^4 = 1)$ -manifolds. We say that a differentiable map $f : M_1 \rightarrow M_2$ is a J -map if and only if the following diagram is commutative

$$\begin{array}{ccc} T_x M_1 & \xrightarrow{f_*} & T_{f(x)} M_2 \\ J_1 \downarrow & & \downarrow J_2 \quad \text{for every } x \in M_1. \\ T_x M_1 & \xrightarrow{f_*} & T_{f(x)} M_2 \end{array}$$

Definition 6.1. We call J -Lie group a Lie Group G with a $(J^4 = 1)$ -structure J such that the usual translations L_g and R_g are J -maps, for every $g \in G$.

Thus, we have

PROPOSITION 6.2. *If G is a J -Lie group, then J is integrable.*

Proof. Since L_g and R_g are J -maps, we have $\text{ad } g \cdot J = J \cdot \text{ad } g$.

In particular for $g = \exp tX$, $X \in \mathfrak{g}$, $t \in \mathbf{R}$, we have

$$\exp(\text{Ad } tX) \cdot JY = J(\exp(\text{Ad } tX)Y), \text{ for every } Y \in \mathfrak{g}. \quad (6.1)$$

Moreover, we obtain

$$\begin{aligned} \exp(\text{Ad } tX)JY &= JY + t[X, JY] + t^2[X, [X, JY]]/2 + \dots \\ J\exp(\text{Ad } tX)Y &= JY + tJ[X, Y] + t^2J[X, [X, Y]]/2 + \dots \end{aligned}$$

Hence, from (6.1) and taking the limit for $t \rightarrow 0$ we deduce $[X, JY] = J[X, Y]$, and also $J[X, Y] = -J[Y, X] = -[Y, JX] = [JX, Y]$.

Thus, it is immediate $N(X, Y) = 0$, $X, Y \in \mathfrak{g}$. □

7. Characterizations of the integrability. Finally, according to the earlier results we can give the following:

THEOREM 7.1. *Let M^n be a differentiable manifold with a tensor field of electromagnetic type and class \tilde{J} . Then the following statements are equivalent:*

- a) *The G -structure P defined by \tilde{J} is integrable;*
- b) *The Nijenhuis tensor of the associated tensor field J is zero;*
- c) *There exists a linear torsionless connection which parallelizes J ;*
- d) *The structure tensor of P is zero;*
- e) *The Nijenhuis tensor of the tensor field \hat{J} corresponding to the canonical prolongation \hat{P} of P is zero;*
- f) *The Nijenhuis tensor of the complete lift J^c of J is zero; moreover,*
- g) *If G is a J -Lie group, then J is integrable.*

When M^n is J -Kaehlerian [10], other conditions can be given.

We note that any linear connection which parallelizes \tilde{J} does not exist.

Acknowledgment. The authors would like to express their deep appreciation to Prof. A. Montesinos for his valuable help.

REFERENCES

- [1] D. Bernard, *Sur la géométrie différentielle des G -structures*, Ann. Inst. Fourier, Grenoble **10** (1960), 151–270.
- [2] V. Guillemin, *The integrability problem for G -structures*, Trans. Amer. Math. Soc. **116** (1965), 544–560.
- [3] J. M. Hernando, P. M. Gadea, A. Montesinos, *G -structures defined by tensor fields of electromagnetic type*, (to appear).
- [4] V. Hlavatý, *Geometry of Einstein's Unified Field Theory*, P. Noordhoff, 1958.
- [5] J. Lehmann-Lejeune, *Intégrabilité des G -structures définies par une 1-forme 0-déformable à valeurs dans le fibre tangent*, Ann. Inst. Fourier, Grenoble, **16** (1966), 329–387.
- [6] R. S. Millman, *Groups in the category of f -manifolds*, Fund. Math., **89** (1975), 1–4.
- [7] R. S. Mishra, *Structures in electromagnetic tensor fields*, Tensor (N.S) **30** (1976), 145–156.

- [8] A. Morimoto, *Prolongations of G-structures to tangent bundles*, Nagoya Math. J. **32** (1968), 67–108.
- [9] E. Reyes, A. Montesinos, P. M. Gadea, *Connections making parallel a metric ($J^4 = 1$)-structure*, Anal. St. Univ. Al. I. Cuza. **28** (1982), 49–56.
- [10] E. Reyes, A. Montesinos, P. M. Gadea, *($J^4 = 1$)-Kähler manifolds*—to appear.
- [11] E. Reyes, A. Montesinos and P. M. Gadea, *Connections partially adapted to a ($J^4 = 1$)-structure*, to appear.
- [12] K. Yano, S. Kobayashi, *Prolongations of tensor fields and connections to tangent bundles. I. General theory*, J. Math. Soc. Japan **18** (1966), 194–210.

Departamento de Geometria
Facultad de Ciencias, Valladolid and
Instituto Jorge Juan
C.S.I.C., Madrid, Spain.

(Received 27 02 1984)