

SPECTRAL RADIUS AND SPECTRUM OF THE COMPRESSION OF A SLANT TOEPLITZ OPERATOR

Taddesse Zegeye and S. C. Arora

Communicated by Miroslub Jevtić

ABSTRACT. A slant Toeplitz operator A_φ with symbol φ in $L^\infty(T)$, where T is the unit circle on the complex plane, is an operator whose representing matrix $M = (a_{ij})$ is given by $a_{ij} = \langle \varphi, z^{2i-j} \rangle$, where $\langle \cdot, \cdot \rangle$ is the usual inner product in $L^2(T)$. The operator B_φ denotes the compression of A_φ to $H^2(T)$ (Hardy space). In this paper, we prove that the spectral radius of B_φ is greater than the spectral radius of A_φ , and if φ and φ^{-1} are in H^∞ , then the spectrum of B_φ contains a closed disc and the interior of this disc consists of eigenvalues with infinite multiplicity.

1. Introduction

Let $\varphi \in L^\infty(T)$. Then $\varphi(z) \sim \sum_{i=-\infty}^{\infty} a_i z^i$, where $a_i = \langle \varphi, z^i \rangle$ is the i -th Fourier coefficient of φ and $\{z^i : i \in Z\}$ is the usual basis, and Z is the set of integers. The slant Toeplitz operator A_φ is defined as follows: $A_\varphi(z^k) = \sum_{i=-\infty}^{\infty} a_{2i-k} z^i$. Furthermore, by [4, Proposition 1] $A_\varphi = WM_\varphi$, where M_φ is a multiplication operator and $Wz^{2n} = z^n$, $Wz^{2n-1} = 0$, for $n \in Z$.

B_φ , the compression of A_φ to $H^2(T)$, is by definition $B_\varphi = PA_\varphi|_{H^2}$. Equivalently, $B_\varphi P = PA_\varphi P$, where P is the orthogonal projection from L^2 on to H^2 . By [4, p. 846], $B_\varphi = WT_\varphi$, where T_φ is the Toeplitz operator on $H^2(T)$.

2. Spectral radius

Our aim is to prove that the spectral radius of B_φ is greater than the spectral radius of A_φ . To do this we need the following lemmas.

2000 *Mathematics Subject Classification*. Primary 47B35; Secondary 47A10.

Key words and phrases. Toeplitz Operator, Slant Toeplitz Operator, Compression, Spectrum.

LEMMA 2.1. $(I - P)M_{z^n} \rightarrow 0$, as $n \rightarrow \infty$ in the strong operator topology, where M_{z^n} is the multiplication by z^n on $L^2(T)$.

PROOF. Let $f \in L^2(T)$ and $f(z) \sim \sum_{i=-\infty}^{\infty} a_i z^i$ be its Fourier expansion. Then $(I - P)M_{z^n} f = (I - P) \left(\sum_{i=-\infty}^{\infty} a_i z^{i+n} \right) = \sum_{i=-\infty}^{-n-1} a_i z^{i+n}$. Since, $\left\| \sum_{i=-\infty}^{-n-1} a_i z^{i+n} \right\|^2 = \sum_{i=-\infty}^{-n-1} |a_i|^2 \rightarrow 0$, as $n \rightarrow \infty$, the assertion follows. \square

The proof of the following lemma is similar to that of [1, Theorem 5].

LEMMA 2.2. $M_{\bar{z}^n} B_\varphi P M_{z^{2n}} \rightarrow A_\varphi$, as $n \rightarrow \infty$, in the strong operator topology.

PROOF. From Lemma 2.1, we know $(I - P)M_{z^n} \rightarrow 0$. This implies that $M_{\bar{z}^n} (I - P)M_{z^n} \rightarrow 0$ which is equivalent to $M_{\bar{z}^n} P M_{z^n} \rightarrow I$. Consider

$$M_{\bar{z}^n} B_\varphi P M_{z^{2n}} = M_{\bar{z}^n} P A_\varphi P M_{z^{2n}} = (M_{\bar{z}^n} P M_{z^n})(M_{\bar{z}^n} A_\varphi M_{z^{2n}})(M_{z^{2n}} P M_{z^{2n}}).$$

Since, for each $n = 1, 2, \dots$, $M_{\bar{z}^n} A_\varphi P M_{z^{2n}} = A_\varphi$, [4, Proposition 3], and the first and the last factors converge to I , as $n \rightarrow \infty$, the assertion follows. \square

The following theorem is proved in [4, p. 851] but we give here a different proof.

THEOREM 2.3. $\|A_\varphi\| = \|B_\varphi\|$.

PROOF. For each $n = 1, 2, \dots$, we have $\|M_{\bar{z}^n} B_\varphi P M_{z^{2n}}\| \leq \|B_\varphi\|$. So from Lemma 2.2, we get $\|A_\varphi\| \leq \|B_\varphi\|$. Since B_φ is the compression of A_φ , we get $\|A_\varphi\| \geq \|B_\varphi\|$. The proof is complete. \square

We are now ready to prove our main result.

THEOREM 2.4. *The spectral radius of B_φ is greater than the spectral radius of A_φ .*

PROOF. First, we prove the following claim by induction.

Claim: For $k = 1, 2, 3, \dots$, $M_{\bar{z}^n} B_\varphi^k P M_{z^{2k n}} \rightarrow A_\varphi^k$, as $n \rightarrow \infty$ in the strong operator topology.

For $k = 1$, the claim is true by Lemma 2.2. Let m be any positive integer and assume that it is true for $k \leq m$, and consider

$$\begin{aligned} & \|M_{\bar{z}^n} B_\varphi^{m+1} P M_{z^{2^{m+1} n}} - A_\varphi^{m+1}\| \\ &= \|M_{\bar{z}^n} (B_\varphi^{m+1} P M_{z^{2^{m+1} n}} - M_{z^n} A_\varphi^{m+1})\| = \|B_\varphi^{m+1} P M_{z^{2^{m+1} n}} - M_{z^n} A_\varphi^{m+1}\| \\ &\leq \|B_\varphi^{m+1} P M_{z^{2^{m+1} n}} - P M_{z^n} A_\varphi^{m+1}\| + \|P M_{z^n} A_\varphi^{m+1} - M_{z^n} A_\varphi^{m+1}\| \end{aligned}$$

By Lemma 2.1, the second term tends to 0 as n approaches infinity. As to the first term, we use $M_{z^n} A_\varphi = A_\varphi M_{z^{2n}}$ [4, Proposition 3] and get the following

approximation

$$\begin{aligned}
\|B_\varphi^{m+1}PM_{z^{2^{m+1}n}} - PM_{z^{2^n}}A_\varphi^{m+1}\| &= \|PA_\varphi B_\varphi^m PM_{z^{2^{m+1}n}} - PA_\varphi M_{z^{2^n}}A_\varphi^m\| \\
&\leq \|A_\varphi\| \|B_\varphi^m PM_{z^{2^{m+1}n}} - M_{z^{2^n}}A_\varphi^m\| \\
&= \|A_\varphi\| \|(B_\varphi^m PM_{z^{2^m n}} - M_{z^{2^n}}A_\varphi^m)M_{z^{2^m n}}\| \\
&\leq \|A_\varphi\| \|B_\varphi^m PM_{z^{2^m n}} - M_{z^{2^n}}A_\varphi^m\| \\
&= \|A_\varphi\| \|M_{z^{2^n}}B_\varphi^m PM_{z^{2^m n}} - A_\varphi^m\|
\end{aligned}$$

By the induction assumption, the last expression tends to 0 as n approaches infinity. Therefore, the claim is proved. The fact that $\|M_{z^{2^n}}B_\varphi^k PM_{z^{2^k n}}\| \leq \|B_\varphi^k\|$, for $k = 1, 2, \dots$, and the above claim imply that $\|B_\varphi^k\| \geq \|A_\varphi^k\|$. This in turn implies that $r(B_\varphi) \geq r(A_\varphi)$, where r represents spectral radius. \square

DEFINITION 2.5. $H^\infty = \{\varphi \in L^\infty(T) : \langle \varphi, z^n \rangle = 0 \text{ for } n < 0\}$. The elements of H^∞ are called analytic and their conjugates are called coanalytic.

COROLLARY 2.6. $r(A_\varphi) = r(B_\varphi)$, if φ is analytic or coanalytic.

PROOF. If φ is analytic, then $B_\varphi = A_\varphi|_{H^2}$. Therefore, for each $k = 1, 2, 3, \dots$, $\|B_\varphi^k\| \leq \|A_\varphi^k\|$. This implies $r(B_\varphi) \leq r(A_\varphi)$. This together with Theorem 2.4, gives the required assertion. If φ is coanalytic, then $B_\varphi^* = A_\varphi^*|_{H^2}$. By the same argument, we get $r(A_\varphi^*) = r(B_\varphi^*)$. Consequently $r(A_\varphi) = r(B_\varphi)$. \square

The following fact is indicated in [4, p. 856], but we give a different proof below.

THEOREM 2.7. If φ is invertible in $L^\infty(T)$, then $r(A_\varphi) \geq [r(A_{\varphi^{-1}})]^{-1}$.

PROOF. First, we show that $\varphi(z)$ is invertible if and only if $\varphi(z^2)$ is invertible. Suppose φ is invertible. Then $\varphi\varphi^{-1} = 1$. Therefore, by [4, p. 846] $W^*\varphi W^*\varphi^{-1} = 1$. Equivalently $\varphi(z^2)\varphi^{-1}(z^2) = 1$. Hence $\varphi(z^2)$ is invertible. Conversely, if $\varphi(z^2)$ is invertible, then $\varphi(z^2)\varphi^{-1}(z^2) = 1$. This and [4, p. 847] implies $W\varphi(z^2)W\varphi^{-1}(z^2) = 1$, which is equivalent to $\varphi\varphi^{-1} = 1$. Therefore φ is invertible.

Let $h(z) = \varphi(z^2)$ be invertible. Then $hh^{-1} = 1$. This and [4, p. 847] implies that $(Wh)(Wh^{-1}) = 1$. Therefore $(Wh)^{-1} = W(h^{-1})$. This in turn implies that, for each $n = 1, 2, 3, \dots$

$$(\varphi^{-1})_n = (\varphi_n)^{-1},$$

where

$$\varphi_n = \overbrace{W(W(\dots(W|h|^2)|h|^2\dots)|h^2|)}^{n \text{ times}}.$$

From this and [4, p. 851], we get

$$\begin{aligned}
r(A_{h^{-1}}) &= \lim_{n \rightarrow \infty} \|(\varphi^{-1})_n\|_\infty^{1/2n} = \lim_{n \rightarrow \infty} \|(\varphi_n)^{-1}\|_\infty^{1/2n} \\
&\geq \left[\lim_{n \rightarrow \infty} \|\varphi_n\|_\infty^{1/2n} \right]^{-1} = [r(A_h)]^{-1}
\end{aligned}$$

Therefore, $r(A_h) \geq [r(A_{h^{-1}})]^{-1}$.

Since $\sigma(A_\varphi) = \sigma(A_{\varphi(z^2)})$, where σ denotes the spectrum [4, Lemma 9], it follows that $r(A_\varphi) \geq [r(A_{\varphi^{-1}})]^{-1}$. \square

COROLLARY 2.8. *$r(B_\varphi) \geq r(B_{\varphi^{-1}})^{-1}$, if φ and φ^{-1} are analytic or φ and φ^{-1} are coanalytic.*

PROOF. This is an immediate consequence of Corollary 2.6 and Theorem 2.7. \square

3. Spectrum

Ho [4] showed that, for invertible φ in L^∞ , the spectrum of A_φ contains a closed disk consisting of eigenvalues of A_φ . We also show, for φ and φ^{-1} in H^∞ , that the spectrum of B_φ contains a closed disc and the interior of this disc consists of eigenvalues with infinite multiplicity, by using the idea of the proof of Proposition 10 in [4].

THEOREM 3.1. *Let φ and φ^{-1} be in H^∞ . Then the spectrum of B_φ contains a closed disc and the interior of this disc consists of eigenvalues with infinite multiplicity.*

PROOF. Assume first that $\lambda \neq 0$. Suppose that $B_{\varphi(z^2)}^* - \lambda$ is onto. Since $B_{\varphi(z^2)} = T_\varphi W$, we have, for f in H^2 ,

$$(B_{\varphi(z^2)}^* - \lambda)f = (W^*T_\varphi - \lambda)f = (W^*T_\varphi - \lambda P_e)f \oplus (-\lambda P_0 f)$$

where P_e is the projection on the closed span of $\{z^{2n} : n = 0, 1, 2, \dots\}$ in $H^2(T)$ and $P_0 = I - P_e$. Now let $0 \neq g_0$ be in $P_0(H^2)$. Since $B_{\varphi(z^2)}^* - \lambda$ is onto, there exists a nonzero vector f in $H^2(T)$ such that $(B_{\varphi(z^2)}^* - \lambda)f = g_0$. But then from the computations above, we have $(W^*T_\varphi - \lambda P_e)f = 0$, because $g_0 \neq 0$. Since $\lambda \neq 0$ and T_φ is invertible [2, Theorem 7.1], it follows that $\lambda W^*T_\varphi(\lambda^{-1} - T_{\varphi^{-1}}W)f = 0$, and the fact that W^* is an isometry implies that $(\lambda^{-1} - T_{\varphi^{-1}}W)f = 0$. This in turn implies $\lambda^{-1} \in \sigma_p(B_{\varphi^{-1}(z^2)})$, where σ_p denotes the point spectrum. Since $\dim P_0(H^2) = \infty$, it follows that λ^{-1} is of infinite multiplicity. Now, for $\lambda \in \rho(B_{\varphi(z^2)}^*)$, the resolvent of $B_{\varphi(z^2)}^*$, the operator $B_{\varphi(z^2)}^* - \lambda$ is invertible (hence onto), so we have

$$D = \{\lambda^{-1} : \lambda \in \rho(B_{\varphi(z^2)}^*)\} \subseteq \sigma_\rho(B_{\varphi^{-1}(z^2)}),$$

Since $B_{\varphi(z^2)} = T_\varphi W$, $B_\varphi = WT_\varphi$ and T_φ is invertible, we have $\sigma_\rho(B_{\varphi(z^2)}) = \sigma_\rho(B_\varphi)$ [3, Problem 61]. Therefore $D \subseteq \sigma_\rho(B_{\varphi^{-1}})$. So by replacing φ^{-1} with φ , we have shown that for any invertible φ in H^∞ , the spectrum of B_φ contains a disc consisting of eigenvalues with infinite multiplicity. Therefore, by the fact that the spectrum of any operator is compact, $\sigma(B_\varphi)$ contains a closed disc and the interior of this disc consists of eigenvalues with infinite multiplicity. \square

REMARK 3.2. If φ and φ^{-1} are coanalytic, then $T_\varphi T_\varphi^{-1} = T_\varphi^{-1} T_\varphi = I$ [2, Theorem 7.1]. Therefore, one can repeat the proof above and arrive at the same conclusion as Theorem 3.1, that is, if φ and φ^{-1} are coanalytic, then the spectrum

of B_φ contains a closed disc and the interior of this disc consists of eigenvalues of infinite multiplicity.

REMARK 3.3. The radius of the closed disc contained in $\sigma_p(B_\varphi)$ is equal to $(r(B_{\varphi^{-1}}))^{-1}$, because if $D_0 = \{0\} \cup \{\lambda^{-1} : |\lambda| > r(B_{\varphi^{-1}})\}$, then $D_0 \subseteq \{\lambda^{-1} : \lambda \in \rho(B_{\varphi^{-1}}^*)\} \cup \{0\} \subseteq \sigma_p(B_\varphi)$ and the radius of the disc D_0 is equal to $(r(B_{\varphi^{-1}}))^{-1}$. Hence $r(B_\varphi) \geq [r(B_{\varphi^{-1}})]^{-1}$. This relation is also proved in Theorem 2.7.

REMARK 3.4. If $\varphi(z) = 1$, then $r(B_{\varphi^{-1}}) = r(B_\varphi) = 1$ by the spectral radius formula for A_φ [4, p. 851] and Corollary 2.6. Hence, the spectrum of B_φ is the closed unit disc by Theorem 3.1 and Remark 3.3. Since the eigenvalues are of infinite multiplicity, it follows that the essential spectrum of B_φ is the same as the spectrum of B_φ .

References

- [1] A. Brown and P. R. Halmos, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. **213** (1964), 89–102.
- [2] R. G. Douglas, *Banach Algebra Techniques in Operator Theory*, Academic Press, New York and London, 1972.
- [3] P. R. Halmos, *A Hilbert Space Problem Book*, Springer-Verlag, New York, 1979.
- [4] M. C. Ho, *Properties of slant Toeplitz operators*, Indiana Univ. Math. J.(c) **45** (1996), 843–862.

Bahir Dar University
 Department of Mathematics
 P.O. Box 79, Bahir Dar
 Ethiopia

(Received 13 01 2000)
 (Revised 10 09 2001)

`bdtc@telecom.net.et`

Department of Mathematics
 University of Delhi
 Delhi 110007
 India