

**WEIGHTED NORM INEQUALITY
FOR THE POISSON INTEGRAL ON THE SPHERE**

BENJAMIN BORDIN, IARA A.A. FERNANDES and SERGIO A. TOZONI

Abstract: We obtain, for each p , $1 < p < \infty$, a necessary and sufficient condition for the Poisson integral of functions defined on the sphere S^n , to be bounded from a weighted space $L^p(S^n, Wd\sigma)$ into a space $L^p(\mathbb{B}, \nu)$, where σ is the Lebesgue measure on S^n and ν is a positive measure on the unit ball \mathbb{B} of \mathbb{R}^{n+1} .

Introduction

In this paper we consider a homogeneous space $X = G/H$ where G is a locally compact Hausdorff topological group and H is a compact subgroup of G which is provided with a quasi-distance d and with a measure μ induced on X by a Haar measure on the topological group G . If $x \in X$ and $r > 0$, $B(x, r)$ will denote the ball $\{y \in X : d(x, y) < r\}$ in X . We also write $\tilde{X} = X \times [0, \infty)$ and if $B = B(x, r)$ we write $\tilde{B} = B(x, r) \times [0, r]$.

We define the maximal operator \mathcal{M} by

$$\mathcal{M}f(x, r) = \sup_{s \geq r} \frac{1}{\mu(B(x, s))} \int_{B(x, s)} |f(y)| d\mu(y)$$

for all real-valued locally integrable function f on X and $(x, r) \in \tilde{X}$. If $r = 0$ the above supremum is taken over all $s > 0$ and $\mathcal{M}f(x, 0) = f^*(x)$ is the Hardy-Littlewood maximal function.

A weight is a positive locally integrable function $W(x)$ on X and we will write $W(A) = \int_A W d\mu$. We say that W is a weight in the class $A_\infty(X)$ if there exist

Received: September 4, 2000; *Revised:* February 23, 2001.

AMS Subject Classification: 42B25, 43A85.

Keywords: Poisson integral; Carleson condition; homogeneous space; weight; sphere.

positive constants C_W and δ such that

$$\frac{\mu(A)}{\mu(B)} \leq C_W \left(\frac{W(A)}{W(B)} \right)^\delta,$$

for all ball $B = B(x, r)$, $x \in X$, $r > 0$, and all Borel subsets A of B . We observe that the above inequality is equivalent to a similar one where μ appears instead of W and conversely (see [5, 1]). We write $L^p(W) = L^p(X, W(x) d\mu(x))$, $1 \leq p < \infty$.

Let $1 < p < \infty$, p' such that $1/p + 1/p' = 1$, let β be a positive measure on the Borel subsets of \tilde{X} and W a weight on X . In Section 2 we introduce a maximal operator of dyadic type \mathcal{M}_d^b , where b is an integer, using partitions of dyadic type for the homogeneous space X introduced in Section 1.

In Section 3 we prove the following theorem.

Theorem 3.1. *Let G be a compact or an Abelian group, let $1 < p < \infty$ and let W be a weight on X such that $W^{1-p'} \in A_\infty(X)$. Then the following conditions are equivalent:*

- (i) *There exists a constant $C > 0$, such that, for all $f \in L^p(W)$,*

$$\int_{\tilde{X}} [\mathcal{M}f(x, r)]^p d\beta(x, r) \leq C \int_X |f(x)|^p W(x) d\mu(x).$$

- (ii) *There exists a constant $C > 0$, such that, for all balls $B = B(z, t)$, $0 \leq t < \infty$,*

$$\int_{\tilde{B}} [\mathcal{M}(W^{1-p'} \chi_B)(x, r)]^p d\beta(x, r) \leq C \int_B W^{1-p'}(x) d\mu(x) < \infty.$$

The above result for $X = \mathbb{R}^n$ was proved in Ruiz-Torrea [7]. A similar result for the fractional maximal operator was obtained in Bernardis-Salinas [1]. The condition (ii) of Theorem 3.1 implies the condition

$$\frac{\beta(\tilde{B})^{1/p}}{\mu(B)} \left(\int_B W^{1-p'}(x) d\mu(x) \right)^{1/p'} \leq C < \infty$$

for all balls B . It was proved in Ruiz-Torrea [8] that the above condition is a necessary and sufficient condition for \mathcal{M} to be a bounded operator from $L^p(X, W(x) d\mu(x))$ into weak $-L^p(\tilde{X}, \beta)$. In the particular case $W(x) \equiv 1$, the condition (ii) of Theorem 3.1 is equivalent to the Carleson's condition for the homogeneous space X :

$$\beta(\tilde{B}) \leq C \mu(B)$$

for all balls B and for a constant $C > 0$.

Now, if $x \in \mathbb{R}^{n+1}$, we write $|x| = (x \cdot x)^{1/2}$ and $d(x, y) = |x - y|$, where $x \cdot y$ is the usual scalar product of x and y in \mathbb{R}^{n+1} . Here S^n will denote the unit n -sphere $\{y \in \mathbb{R}^{n+1} : |y| = 1\}$ in \mathbb{R}^{n+1} , σ the normalized Lebesgue measure on S^n and $h : [1 - \sqrt{2}, 1] \rightarrow [0, 2]$ will be the function defined by $h(r) = \sqrt{2}(1 - r)$.

The Poisson kernel for the sphere S^n is given by

$$P_{ry}(x) = \frac{1}{\omega_n} \frac{1 - r^2}{|ry - x|^{n+1}}$$

for $x, y \in S^n$ and $0 \leq r < 1$, where ω_n is the area of the sphere S^n . For a real-valued integrable function f we denote by $u_f(ry)$ the Poisson integral

$$u_f(ry) = \int_{S^n} P_{ry}(x) f(x) d\sigma(x) .$$

and we define the maximal function u_f^* by

$$u_f^*(ry) = \sup_{0 \leq s \leq r} |u_f(sy)|, \quad 0 \leq r < 1, \quad y \in S^n .$$

If B is the open ball $B(z, t) = \{x \in S^n : |x - z| < t\}$, $0 < t \leq 2$, we define

$$\bar{B} = \{sx : h^{-1}(t) \leq s \leq 1, x \in B\} \quad \text{if } 0 < t \leq \sqrt{2} ;$$

$$\bar{B} = \{sx : 0 \leq s \leq 1, x \in B\} \quad \text{if } \sqrt{2} \leq t \leq 2 .$$

We observe that \bar{B} is a truncated cone in the ball $\mathbb{B} = \{y \in \mathbb{R}^{n+1} : |y| \leq 1\}$ in \mathbb{R}^{n+1} if $0 < t \leq \sqrt{2}$ and a cone if $\sqrt{2} \leq t \leq 2$.

In Section 4 we prove the following result.

Theorem 4.1. *Let $1 < p < \infty$, let W be a weight on S^n such that $W^{1-p'} \in A_\infty(S^n)$ and let ν be a Borel positive measure on S^n . Then the following conditions are equivalent:*

- (i) *There exists a constant $C > 0$, such that, for all $f \in L^p(W)$,*

$$\int_{\mathbb{B}} [u_f^*(y)]^p d\nu(y) \leq C \int_{S^n} |f(x)|^p W(x) d\sigma(x) .$$

- (ii) *There exists a constant $C > 0$, such that, for all balls $B = B(z, t)$, $0 < t \leq 2$,*

$$\int_{\bar{B}} [u_{W^{1-p'} \chi_B}^*(y)]^p d\nu(y) \leq C \int_B W^{1-p'}(x) d\sigma(x) < \infty .$$

We point out that the Theorem 4.1 for $W \equiv 1$ and $n = 1$ was proved in Carleson [2].

1 – Preliminaries

In this section we introduce some notations, definitions and basic facts.

Let G be a locally compact Hausdorff topological group with unit element e , H be a compact subgroup and $\pi : G \mapsto G/H$ the canonical map. Let dg denote a left Haar measure on G , which we assume to be normalized in the case of G to be compact. If A is a Borel subset of G we will denote by $|A|$ the Haar measure of A . The homogeneous space $X = G/H$ is the set of all left cosets $\pi(g) = gH, g \in G$, provided with the quotient topology. The Haar measure dg induces a measure μ on the Borel σ -field on X . For $f \in L^1(X)$,

$$\int_X f(x) d\mu(x) = \int_G f \circ \pi(g) dg .$$

We observe that the group G acts transitively on X by the map $(g, \pi(h)) \mapsto g\pi(h) = \pi(gh)$, that is, for all $x, y \in X$, there exists $g \in G$ such that $gx = y$. We also observe that the measure μ on X is invariable on the action of G , that is, if $f \in L^1(X)$, $g \in G$ and $R_g f(x) = f(g^{-1}x)$, then

$$\int_X f(x) d\mu(x) = \int_X R_g f(x) d\mu(x) .$$

Definition 1.1. A quasi-distance on X is a map $d : X \times X \mapsto [0, \infty)$ satisfying:

- (i) $d(x, y) = 0$ if and only if $x = y$;
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (iii) $d(gx, gy) = d(x, y)$ for all $g \in G, x, y \in X$;
- (iv) there exists a constant $K \geq 1$ such that, for all $x, y, z \in X$,

$$d(x, y) \leq K[d(x, z) + d(z, y)] ;$$

- (v) the balls $B(x, r) = \{y \in X : d(x, y) < r\}$, $x \in X, r > 0$, are relatively compact and measurable, and the balls $B(\mathbb{1}, r)$, $r > 0$, form a basis of neighborhoods of $\mathbb{1} = \pi(e)$;
- (vi) there exists a constant $A \geq 1$ such that, for all $r > 0$ and $x \in X$,

$$\mu(B(x, 2r)) \leq A\mu(B(x, r)) . \square$$

In this paper X will denote a homogeneous space provided with a quasi-distance d .

Given a quasi-distance d on X , there exists a distance ρ on X and a positive real number γ such that d is equivalent to ρ^γ (see [5]). Therefore the family of d -balls is equivalent to the family of ρ^γ -balls and ρ^γ -balls are open sets.

It follows by Definition 1.1(iii) that $B(gx, r) = gB(x, r)$ for all $g \in G$, $x \in X$ and $r > 0$, and hence $\mu(B(gx, r)) = \mu(B(x, r))$. Thus we can write $X = \bigcup_{j \geq 1} g_j B(x, r)$ where (g_j) is a sequence of elements of G and consequently $\mu(B(x, r)) > 0$. In particular, X is separable.

Lemma 1.1. *Let b be a positive integer and let $\lambda = 8K^5$. Then for each integer k , $-b \leq k \leq b$, there exist an enumerable Borel partition \mathcal{A}_k^b of X and a positive constant C depending only on X , such that:*

(i) *for all $Q \in \mathcal{A}_k^b$, $-b \leq k \leq b$, there exists $x_Q \in Q$ such that*

$$B(x_Q, \lambda^k) \subset Q \subset B(x_Q, \lambda^{k+1})$$

and

$$\mu(B(x_Q, \lambda^{k+1})) \leq C\mu(Q) ;$$

(ii) *if $-b \leq k < b$, $Q_1 \in \mathcal{A}_{k+1}^b$, $Q_2 \in \mathcal{A}_k^b$ and $Q_1 \cap Q_2 \neq \emptyset$, then $Q_2 \subset Q_1$, and*

$$0 < \mu(Q_1) \leq C\mu(Q_2) ;$$

(iii) *for all $x \in X$ and r , $\lambda^{-b-1} \leq r \leq \lambda^b$, there exist $Q \in \mathcal{A}_k^b$ for some $-b \leq k \leq b$ and $g \in G$ such that $d(gx, x) \leq \lambda^{k+1}$, $B(x, r) \subset gQ$ and*

$$\mu(Q) \leq C\mu(B(x, r)) .$$

Proof: The properties (i) and (ii) follow by Lemma 3.21, p. 852 of [9] and by Definition 1.1.

Let us prove (iii). Given $x \in X$ and $\lambda^{-b-1} \leq r \leq \lambda^b$, let $-b \leq k \leq b$ such that $\lambda^{k-1} \leq r \leq \lambda^k$. There exists an unique $Q \in \mathcal{A}_k^b$ such that $x \in Q$. Consider x_Q as in (i) and $g \in G$ such that $x = gx_Q$. If a is an integer such that $2^{a-1} < \lambda \leq 2^a$, then by (i) we have

$$B(x, r) \subset B(gx_Q, \lambda^k) \subset gQ \subset B(x, \lambda^{k+1})$$

and hence by Definition 1.1(vi) we have

$$\begin{aligned}\mu(Q) &\leq \mu(B(x_Q, 2^a \lambda^k)) \\ &\leq A^a \mu(B(x_Q, \lambda^k)) \\ &\leq A^{2a} \mu(B(x, r)) .\end{aligned}$$

We also have that $d(gx, x) = d(x, x_Q) \leq \lambda^{k+1}$. ■

Let $(\Omega, \mathcal{F}, \nu)$ be a σ -finite measure space, let $(\mathcal{F}_k)_{k \in \mathbb{Z}}$ be an increasing sequence of sub- σ -fields of \mathcal{F} , and for each $k \in \mathbb{Z}$, consider a real-valued \mathcal{F}_k -measurable function f_k . We say that the sequence $(f_k)_{k \in \mathbb{Z}}$ is a martingale with respect to the sequence $(\mathcal{F}_k)_{k \in \mathbb{Z}}$ if, for all $k \in \mathbb{Z}$ and all $A \in \mathcal{F}_k$ such that $\nu(A) < \infty$, we have that

$$\int_A |f_k| d\nu < \infty, \quad \int_A f_k d\nu = \int_A f_{k+1} d\nu .$$

Now, consider a σ -finite measure ν on the Borel σ -field of X and let \mathcal{F}_k be the σ -field generated by the partition \mathcal{A}_{-k}^b for $-b \leq k \leq b$, by \mathcal{A}_{-b}^b for $k \geq b$ and by \mathcal{A}_b^b for $k \leq -b$. If $f \in L^1(X, \nu)$,

$$f_k(x) = E[f | \mathcal{F}_k](x) = \sum_{Q \in \mathcal{A}_k^b} \left(\frac{1}{\nu(Q)} \int_Q f(y) d\nu(y) \right) \chi_Q(x), \quad -b \leq k \leq b ;$$

and $f_k = f_b$ for $k \geq b$, $f_k = f_{-b}$ for $k \leq -b$, then $(f_k)_{k \in \mathbb{Z}}$ is a martingale with respect to the sequence $(\mathcal{F}_k)_{k \in \mathbb{Z}}$. We define the maximal operator M_ν^b , for all $f \in L^1(X, \nu)$ by

$$M_\nu^b f(x) = \sup_{k \in \mathbb{Z}} E[|f| | \mathcal{F}_k](x) = \sup_{\substack{x \in Q \\ Q \in \mathcal{A}^b}} \frac{1}{\nu(Q)} \int_Q |f(y)| d\nu(y) .$$

where $\mathcal{A}^b = \bigcup_{-b \leq k \leq b} \mathcal{A}_k^b$.

The next result can be found in Dellacherie-Meyer [3], number 40, p. 37.

Theorem 1.1. *If $1 < p < \infty$ and $f \in L^p(X, \nu)$, then*

$$\|M_\nu^b f\|_{L^p(X, \nu)} \leq p' \|f\|_{L^p(X, \nu)} .$$

2 – A maximal operator of dyadic type

Let b be a fixed positive integer. Given $Q \in \mathcal{A}^b = \bigcup_{-b \leq k \leq b} \mathcal{A}_k^b$, where \mathcal{A}_k^b are the partitions of X in Lemma 1.1, \tilde{Q} will denote the subset $Q \times [0, \alpha^{-1}(\mu(Q))]$ of $\tilde{X} = X \times [0, \infty)$, where $\alpha : [0, \infty) \rightarrow [0, \infty)$ is the function defined by $\alpha(r) = \mu(B(\mathbb{1}, r))$, $\mathbb{1} = \pi(e)$.

If f is a real-valued locally integrable function on X , we define, for each $(x, r) \in \tilde{X}$,

$$\mathcal{M}_d^b f(x, r) = \sup_{\substack{x \in Q \in \mathcal{A}^b \\ \mu(Q) \geq \alpha(r)}} \frac{1}{\mu(Q)} \int_Q |f(y)| d\mu(y) .$$

If $\mu(Q) < \alpha(r)$ for all $Q \in \mathcal{A}^b$ such that $x \in Q$, we define $\mathcal{M}_d^b f(x, r) = 0$.

Lemma 2.1. *Let W be a weight and let A be a measurable subset of X . If $1 < p < \infty$ and $W^{-1}\chi_A \notin L^{p'}(W)$, then there exists a positive function $f \in L^p(W)$ such that*

$$\int_A f(x) d\mu(x) = \infty .$$

Proof: Let ψ be the linear functional on $L^p(W)$ given by $\psi(g) = \int_A g d\mu$. Since $W^{-1}\chi_A \notin L^{p'}(W)$, it follows by the Riesz representation theorem that ψ is not continuous. Therefore, there exists $\varepsilon > 0$, such that, for each positive integer m , there exists $g_m \in L^p(W)$ such that $\|g_m\|_{L^p(W)} \leq 2^{-m}$ and $|\psi(g_m)| \geq \varepsilon$. We set $f_m(x) = |g_1(x)| + \dots + |g_m(x)|$ and then, for all $m, k \geq 1$,

$$\|f_{m+k} - f_m\|_{L^p(W)} \leq \|g_{m+1}\|_{L^p(W)} + \dots + \|g_{m+k}\|_{L^p(W)} < 2^{-m} .$$

Hence (f_m) is a Cauchy sequence in $L^p(W)$ and therefore there exists $f \in L^p(W)$ such that $f_m \rightarrow f$ in $L^p(W)$. On the other hand

$$\psi(f_m) \geq |\psi(g_1)| + \dots + |\psi(g_m)| \geq m\varepsilon .$$

But $f_m \uparrow f$ a.e. and thus by the monotone convergence theorem we obtain

$$\int_A f d\mu = \lim_{m \rightarrow \infty} \psi(f_m) = \infty . \blacksquare$$

Theorem 2.1. *Given a weight W on X , a positive measure β on \tilde{X} , and $1 < p < \infty$, the following conditions are equivalent:*

- (i) There exists a constant $C > 0$, such that, for all $f \in L^p(W)$ and all positive integer b ,

$$\int_{\tilde{X}} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) \leq C \int_X |f(x)|^p W(x) d\mu(x) .$$

- (ii) There exists a constant $C > 0$, such that, for all $Q \in \mathcal{A}^b$ and all positive integer b ,

$$\int_{\tilde{Q}} [\mathcal{M}_d^b (W^{1-p'} \chi_Q)(x, r)]^p d\beta(x, r) \leq C \int_Q W^{1-p'}(x) d\mu(x) < \infty .$$

Proof: The proof of (i) \Rightarrow (ii) is exactly as the proof of (i) \Rightarrow (ii) in Theorem 3.1.

Proof of (ii) \Rightarrow (i): Let us fix $f \in L^p(W)$ and for each $k \in \mathbb{Z}$, let Ω_k be the set

$$\Omega_k = \left\{ (x, r) \in \tilde{X} : \mathcal{M}_d^b f(x, r) > 2^k \right\} .$$

For each $k \in \mathbb{Z}$, we denote by C_k^0 the family formed by all $Q \in \mathcal{A}^b$ such that

$$|f|_Q = \frac{1}{\mu(Q)} \int_Q |f(y)| d\mu(y) > 2^k .$$

Since for every $Q \in \mathcal{A}_k^b$, $-b \leq k < b$, there exists $Q' \in \mathcal{A}_{k+1}^b$ such that $Q \subset Q'$, then every element $Q \in C_k^0$ is contained in a maximal element $Q' \in C_k^0$. We denote by C_k the family $\{Q_j^k : j \in J_k\}$ formed by all maximal elements $Q \in C_k^0$. Since \mathcal{A}_k^b is a partition of X and all elements of C_k are maximal, we can conclude that the sets $Q_j^k, j \in J_k$, are pairwise disjoint. Therefore the sets $\tilde{Q}_j^k, j \in J_k$, are also pairwise disjoint and,

$$\Omega_k = \bigcup_{j \in J_k} \tilde{Q}_j^k .$$

Now, for each $k \in \mathbb{Z}$ and each $j \in J_k$, let

$$E_j^k = \tilde{Q}_j^k \setminus \Omega_{k+1} .$$

Then the sets E_j^k and E_i^h are disjoint for $(k, j) \neq (h, i)$ and

$$\left\{ (x, r) : \mathcal{M}_d^b f(x, r) > 0 \right\} = \bigcup_{k \in \mathbb{Z}} (\Omega_k \setminus \Omega_{k+1}) = \bigcup_{k \in \mathbb{Z}} \bigcup_{j \in J_k} E_j^k .$$

Therefore

$$\begin{aligned}
 \int_{\tilde{X}} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) &= \sum_{k,j} \int_{E_j^k} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) \\
 &\leq 2^p \sum_{k,j} \beta(E_j^k) (2^k)^p \\
 (2.1) \qquad \qquad \qquad &\leq 2^p \sum_{k,j} \beta(E_j^k) \left(\frac{1}{\mu(Q_j^k)} \int_{Q_j^k} |f(x)| d\mu(x) \right)^p .
 \end{aligned}$$

Now, we introduce the following notations:

$$\begin{aligned}
 \nu(x) &= W^{1-p'}(x), \quad \nu(A) = \int_A \nu(x) d\mu(x) , \\
 \gamma_{k,j} &= \beta(E_j^k) \left(\frac{\nu(Q_j^k)}{\mu(Q_j^k)} \right)^p, \quad g_{k,j} = \left(\frac{1}{\nu(Q_j^k)} \int_{Q_j^k} \frac{|f(x)|}{\nu(x)} \nu(x) d\mu(x) \right)^p, \\
 Y &= \{(k, j) : k \in \mathbb{Z}, j \in J_k\}, \quad \Gamma(\lambda) = \{(k, j) \in Y : g_{k,j} > \lambda\} .
 \end{aligned}$$

Let γ be the measure on Y such that $\gamma(\{(k, j)\}) = \gamma_{k,j}$ and let g be the function defined on Y by $g((k, j)) = g_{k,j}$. We have that

$$\gamma_{k,j} g_{k,j} = \beta(E_j^k) \left(\frac{1}{\mu(Q_j^k)} \int_{Q_j^k} |f(x)| d\mu(x) \right)^p$$

and hence it follows by (2.1) that

$$\begin{aligned}
 \int_{\tilde{X}} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) &\leq 2^p \sum_{k,j} \gamma_{k,j} g_{k,j} \\
 &= 2^p \int_0^\infty \gamma(\Gamma(\lambda)) d\lambda \\
 &= 2^p \int_0^\infty \left(\sum_{(k,j) \in \Gamma(\lambda)} \gamma_{k,j} \right) d\lambda \\
 (2.2) \qquad \qquad \qquad &= 2^p \int_0^\infty \sum_{(k,j) \in \Gamma(\lambda)} \int_{E_j^k} \left(\frac{\nu(Q_j^k)}{\mu(Q_j^k)} \right)^p d\beta(x, r) d\lambda .
 \end{aligned}$$

For each $\lambda > 0$, let $\{Q_i^\lambda : i \in I_\lambda\}$ be the family formed by all maximal elements of the family

$$\left\{ Q_j^k : (k, j) \in \Gamma(\lambda) \right\} = \left\{ Q_j^k : \frac{1}{\nu(Q_j^k)} \int_{Q_j^k} \frac{|f(x)|}{\nu(x)} \nu(x) d\mu(x) > \lambda^{1/p} \right\} .$$

If $Q_j^k \subset Q_i^\lambda$ and $(x, r) \in E_j^k$, then $x \in Q_j^k$, $\mu(Q_j^k) \geq \alpha(r)$ and thus

$$\mathcal{M}_d^b(\nu\chi_{Q_i^\lambda})(x, r) = \sup_{\substack{x \in Q \in \mathcal{A}^b \\ \mu(Q) \geq \alpha(r)}} \frac{\nu(Q \cap Q_i^\lambda)}{\mu(Q)} \geq \frac{\nu(Q_j^k)}{\mu(Q_j^k)}.$$

Therefore, if $Q_j^k \subset Q_i^\lambda$ we obtain

$$(2.3) \quad \int_{E_j^k} \left(\frac{\nu(Q_j^k)}{\mu(Q_j^k)} \right)^p d\beta(x, r) \leq \int_{E_j^k} [\mathcal{M}_d^b(\nu\chi_{Q_i^\lambda})(x, r)]^p d\beta(x, r).$$

Taking into account that the sets E_j^k are disjoint, it follows from (2.2), (2.3) and by the hypothesis that

$$\begin{aligned} \int_{\tilde{X}} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) &\leq 2^p \int_0^\infty \sum_{i \in I_\lambda} \sum_{\substack{(k,j) \in \Gamma(\lambda) \\ Q_j^k \subset Q_i^\lambda}} \int_{E_j^k} [\mathcal{M}_d^b(\nu\chi_{Q_i^\lambda})(x, r)]^p d\beta(x, r) \\ &\leq 2^p \int_0^\infty \sum_{i \in I_\lambda} \int_{\tilde{Q}_i^\lambda} [\mathcal{M}_d^b(\nu\chi_{Q_i^\lambda})(x, r)]^p d\beta(x, r) \\ &\leq C 2^p \int_0^\infty \sum_{i \in I_\lambda} \int_{Q_i^\lambda} \nu(x) d\mu(x) \\ (2.4) \quad &= C 2^p \int_0^\infty \nu \left(\bigcup_{(k,j) \in \Gamma(\lambda)} Q_j^k \right) d\mu(x). \end{aligned}$$

It follows by the definition of the maximal operator M_ν^b in Section 1 and by the definition of $\Gamma(\lambda)$ that

$$(2.5) \quad \bigcup_{(k,j) \in \Gamma(\lambda)} Q_j^k \subset \left\{ x \in X : M_\nu^b \left(\frac{|f|}{\nu} \right) (x) > \lambda^{1/p} \right\}.$$

Then, by (2.4), (2.5) and Theorem 1.1,

$$\begin{aligned} \int_{\tilde{X}} [\mathcal{M}_d^b f(x, r)]^p d\beta(x, r) &\leq C 2^p \int_0^\infty \nu \left(\left\{ x : \left(M_\nu^b \left(\frac{|f|}{\nu} \right) (x) \right)^p > \lambda \right\} \right) d\lambda \\ &= C 2^p \int_X \left(M_\nu^b \left(\frac{|f|}{\nu} \right) (x) \right)^p \nu(x) d\mu(x) \\ &\leq C 2^p (p')^p \int_X \frac{|f(x)|^p}{(\nu(x))^p} \nu(x) d\mu(x) \\ &= C 2^p (p')^p \int_X |f(x)|^p W(x) d\mu(x) . \blacksquare \end{aligned}$$

Remark 2.1. Let us fix $g \in G$ and let $g^{-1}\mathcal{A}_k^b = \{g^{-1}Q : Q \in \mathcal{A}_k^b\}$, $g^{-1}\mathcal{A}^b = \{g^{-1}Q : Q \in \mathcal{A}^b\}$. Then for each $-b \leq k \leq b$, $g^{-1}\mathcal{A}_k^b$ is a partition of X and Lemma 1.1 and Theorem 1.1 in Section 1 also hold, with the same constants, when we change \mathcal{A}_k^b for $g^{-1}\mathcal{A}_k^b$. If f is a real-valued locally integrable function on X , we define

$$\mathcal{M}_d^{b,g}f(x, r) = \sup_{\substack{x \in Q \in g^{-1}\mathcal{A}^b \\ \mu(Q) \geq \alpha(r)}} \frac{1}{\mu(Q)} \int_Q |f(y)| d\mu(y) .$$

Then

$$\mathcal{M}_d^b(R_g f)(gx, r) = \mathcal{M}_d^{b,g}f(x, r)$$

where $R_g f(x) = f(g^{-1}x)$. The Theorem 2.1 also hold, with the same proof, when we change the operator \mathcal{M}_d^b for $\mathcal{M}_d^{b,g}$ and the family \mathcal{A}^b for $g^{-1}\mathcal{A}^b$. \square

3 – The boundedness of the operator \mathcal{M}

Given a positive integer b and a real-valued locally integrable function f on X , we define for $(x, r) \in \tilde{X}$,

$$\mathcal{M}^b f(x, r) = \sup_{\max\{\lambda^{-b-1}, r\} \leq s \leq \lambda^b} \frac{1}{\mu(B(x, s))} \int_{B(x, s)} |f(y)| d\mu(y) .$$

We define $\mathcal{M}^b f(x, r) = 0$ if $r > \lambda^b$ and we observe that $\mathcal{M}^b f(x, r) \uparrow \mathcal{M}f(x, r)$ if $b \uparrow \infty$ for all $(x, r) \in \tilde{X}$.

Let us denote

$$G_b = \left\{ g \in G : d(gx, x) \leq \lambda^{b+1} \text{ for all } x \in X \right\} .$$

If $d(g\mathbb{1}, \mathbb{1}) = d(gx, x)$ for all $x \in X$ and $g \in G$, in particular if G is an Abelian group, then

$$G_b = \left\{ g \in G : d(g\mathbb{1}, \mathbb{1}) \leq \lambda^{b+1} \right\} ,$$

and hence G_b is relatively compact in G and $0 < |G_b| < \infty$ (see [4]).

Lemma 3.1. *Let b be a positive integer, $g \in G$, let $\mathcal{M}_d^{b,g}$ be the maximal operator defined in Remark 2.1, let f be a real-valued locally integrable function on X and let $(x, r) \in \tilde{X}$. Then*

$$(3.1) \quad \mathcal{M}_d^{b,g}f(x, r) \leq C\mathcal{M}f(x, r) .$$

If G is a compact or an Abelian group, then

$$(3.2) \quad \mathcal{M}^b f(x, r) \leq \frac{C}{|G_b|} \int_{G_b} \mathcal{M}_d^{b,g} f(x, r) dg .$$

The constants C in (3.1) and in (3.2) depend only on X and if X is compact we can change G_b for G .

Proof: Let us fix $(x, r) \in \tilde{X}$ and $g \in G$. If $\mu(Q) < \alpha(r)$ for all $Q \in \mathcal{A}^b$ such that $x \in g^{-1}Q$, we have $\mathcal{M}_d^{b,g} f(x, r) = 0$. Thus to prove (3.1), it is enough to consider $Q \in \mathcal{A}_k^b$, $-b \leq k \leq b$, such that $x \in g^{-1}Q$ and $\mu(Q) \geq \alpha(r)$. By Lemma 1.1(i) there exist $x_Q \in Q$ such that $Q \subset B(x_Q, \lambda^{k+1})$ and $\mu(B(x_Q, \lambda^{k+1})) \leq C\mu(Q)$. For $t = 2K\lambda^{k+1}$ we have $B(g^{-1}x_Q, \lambda^{k+1}) \subset B(x, t)$ and hence

$$\alpha(t) = \mu(B(x, t)) \geq \mu(B(g^{-1}x_Q, \lambda^{k+1})) \geq \mu(Q) \geq \alpha(r) .$$

If $2^{a-1} < K \leq 2^a$, it follows by Definition 1.1(vi) that

$$\mu(B(x, t)) \leq A^{a+1} \mu(B(x_Q, \lambda^{k+1})) \leq A^{a+1} C \mu(g^{-1}Q) .$$

Therefore

$$\frac{1}{\mu(g^{-1}Q)} \int_{g^{-1}Q} |f(y)| d\mu(y) \leq \frac{A^{a+1}C}{\mu(B(x, t))} \int_{B(x, t)} |f(y)| d\mu(y) \leq A^{a+1} C \mathcal{M}f(x, r)$$

and hence we obtain (3.1).

Let us fix $(x, r) \in \tilde{X}$. If $r > \lambda^b$ we have $\mathcal{M}^b f(x, r) = 0$ and thus we can suppose $r \leq \lambda^b$. Given s such that $\lambda^{-b-1} \leq s \leq \lambda^b$ and $s \geq r$, by Lemma 1.1(iii), there exist $Q \in \mathcal{A}_k^b$ for some $-b \leq k \leq b$ and $g \in G_b$, such that $B(x, s) \subset g^{-1}Q$ and $\mu(Q) \leq C\mu(B(x, s))$. Then

$$\frac{1}{\mu(B(x, s))} \int_{B(x, s)} |f(y)| d\mu(y) \leq \frac{C}{\mu(g^{-1}Q)} \int_{g^{-1}Q} |f(y)| d\mu(y) \leq C \mathcal{M}_d^{b,g} f(x, r)$$

since $\mu(Q) \geq \mu(B(x, s)) \geq \alpha(r)$. Therefore, integrating both sides of the above inequality on G_b , we have that

$$\frac{1}{\mu(B(x, s))} \int_{B(x, s)} |f(y)| d\mu(y) \leq \frac{C}{|G_b|} \int_{G_b} \mathcal{M}_d^{b,g} f(x, r) dg$$

and hence we obtain (3.2). ■

Proof of Theorem 3.1: First we prove the implication (i) \Rightarrow (ii). Suppose that there exists $B = B(z, t)$, $0 < t < \infty$ such that

$$\int_B W^{1-p'}(x) d\mu(x) = \infty .$$

Then $W^{-1}\chi_B \notin L^{p'}(W)$ and thus, by Lemma 2.1, there exists a positive function $f \in L^p(W)$ such that

$$\int_B f(x) d\mu(x) = \infty .$$

Therefore given $(x, r) \in \tilde{X}$, there exists $s \geq r$ such that $B \subset B(x, s)$ and hence $\mathcal{M}f(x, r) = \infty$. Since β is a positive measure, we have a contradiction of the condition (i). Thus

$$\int_B W^{1-p'}(x) d\mu(x) < \infty .$$

To obtain the inequality in (ii) it is sufficient to choose $f(x) = W^{1-p'}(x) \chi_B(x)$ in the hypothesis.

Let us prove (ii) \Rightarrow (i). We fix a positive integer $b, g \in G$ and $Q \in \mathcal{A}_k^b, -b \leq k \leq b$. Then, by Lemma 1.1(i) there exist $x_Q \in Q$ such that $Q \subset B(x_Q, \lambda^{k+1})$ and $\mu(B(x_Q, \lambda^{k+1})) \leq C\mu(Q)$. We write $B = B(g^{-1}x_Q, \lambda^{k+1}), Q' = g^{-1}Q$ and $\nu = W^{1-p'}$. Since $\nu \in A_\infty(X)$, there exist positive constants C_ν and δ , depending only on ν , such that

$$\frac{\mu(Q')}{\mu(B)} \leq C_\nu \left(\frac{\nu(Q')}{\nu(B)} \right)^\delta .$$

Therefore

$$\nu(B) \leq C_\nu^{1/\delta} \left(\frac{\mu(B)}{\mu(Q')} \right)^{1/\delta} \nu(Q') \leq C_1 \nu(Q') .$$

Then by the hypothesis and (3.1) we obtain

$$\begin{aligned} \int_{\tilde{Q}'} [\mathcal{M}_d^{b,g}(W^{1-p'} \chi_{Q'})(x, r)]^p d\beta(x, r) &\leq C_2 \int_{\tilde{B}} [\mathcal{M}(\nu \chi_B)(x, r)]^p d\beta(x, r) \\ &\leq C_3 \nu(B) \\ &\leq C_4 \int_{Q'} W^{1-p'}(x) d\mu(x) . \end{aligned}$$

Since the constant C_4 depends only on p, W and β , then by Theorem 2.1 and Remark 2.1, there exists a constant C_5 such that,

$$(3.3) \quad \int_{\tilde{X}} [\mathcal{M}_d^{b,g} f(x, r)]^p d\beta(x, r) \leq C_5 \int_X |f(x)|^p W(x) d\mu(x)$$

for all $f \in L^p(W)$ and all $g \in G$. Then, it follows by (3.2), (3.3) and by Jensen's inequality that

$$\int_{\tilde{X}} [\mathcal{M}^b f(x, r)]^p d\beta(x, r) \leq \int_{\tilde{X}} \left(\frac{C_6}{|G_b|} \int_{G_b} \mathcal{M}_d^{b,g} f(x, r) dg \right)^p d\beta(x, r)$$

$$\begin{aligned} &\leq C_6^p \int_{G_b} \int_{\tilde{X}} [\mathcal{M}_d^{b,g} f(x, r)]^p d\beta(x, r) \frac{dg}{|G_b|} \\ &\leq C_6^p C_5 \int_X |f(x)|^p W(x) d\mu(x) . \end{aligned}$$

The result follows by the Monotone Convergence Theorem. ■

Remark 3.1. (a) For $W \equiv 1$, the condition (ii) of Theorem 3.1 is given by

$$(3.4) \quad \int_{\tilde{B}} [\mathcal{M}(\chi_B)(x, r)]^p d\beta(x, r) \leq C \mu(B)$$

for all balls B . Let us fix $B = B(z, t)$, $0 < t < \infty$. Then, it follows as in the proof of inequality (3.1) of Lemma 3.1 that there exists a constant $C > 0$ such that

$$C \leq \mathcal{M}(\chi_B)(x, r) \leq 1$$

for all $(x, r) \in \tilde{B}$. Therefore, from (3.4) we obtain

$$C^p \beta(\tilde{B}) \leq \int_{\tilde{B}} [\mathcal{M}(\chi_B)(x, r)]^p d\beta(x, r) \leq C \mu(B) .$$

Then, the condition (3.4) implies the condition:

$$(3.5) \quad \beta(\tilde{B}) \leq C \mu(B)$$

for a constant $C > 0$ and all balls B . But, from the condition (3.5) we obtain

$$\int_{\tilde{B}} [\mathcal{M}(\chi_B)(x, r)]^p d\beta(x, r) \leq \beta(\tilde{B}) \leq C \mu(B) ,$$

and therefore the conditions (3.4) and (3.5) are equivalent. The condition (3.5) is the Carleson condition for the homogeneous space X (see [8]).

(b) Let $B = B(z, t)$, $0 < t < \infty$ and $\nu = W^{1-p'}$. Then

$$C \frac{\nu(B)}{\mu(B)} \leq \mathcal{M}(\nu\chi_B)(x, r)$$

for all $(x, r) \in \tilde{B}$. Therefore, from the condition (ii) of Theorem 3.1 we obtain

$$\begin{aligned} \beta(\tilde{B})^{1/p} &= \left(\frac{\mu(B)}{\nu(B)} \right) \left[\int_{\tilde{B}} \left(\frac{\nu(B)}{\mu(B)} \right)^p d\beta(x, r) \right]^{1/p} \\ &\leq C \left(\frac{\mu(B)}{\nu(B)} \right) \left[\int_{\tilde{B}} [\mathcal{M}(\nu\chi_B)(x, r)]^p d\beta(x, r) \right]^{1/p} \\ &\leq C' \left(\frac{\mu(B)}{\nu(B)} \right) \nu(B)^{1/p} . \end{aligned}$$

Then, the condition (ii) of Theorem 3.1 implies the condition:

$$(3.6) \quad \frac{\beta(\tilde{B})^{1/p}}{\mu(B)} \left(\int_B W^{1-p'}(x) d\mu(x) \right)^{1/p'} \leq C$$

for a constant $C > 0$ and all balls B . It was proved in Ruiz-Torrea [8] that the condition (3.6) is a necessary and sufficient condition for \mathcal{M} to be a bounded operator from $L^p(X, W(x) d\mu(x))$ into weak - $L^p(\tilde{X}, \beta)$. \square

4 – The boundedness of the Poisson integral

Let $\xi : [0, \pi]^{n-1} \times [0, 2\pi] \rightarrow S^n$ be the function defined by $\xi(\theta) = \xi(\theta_1, \dots, \theta_n) = (x_1, \dots, x_{n+1})$, where

$$x_1 = \cos \theta_1; \quad x_i = \cos \theta_i \prod_{j=1}^{i-1} \sin \theta_j, \quad 2 \leq i \leq n; \quad x_{n+1} = \prod_{j=1}^n \sin \theta_j .$$

We identify $S^n \times [0, 1]$ with the ball $\mathbb{B} = \{y \in \mathbb{R}^{n+1} : |y| \leq 1\}$ using the application $(y, r) \mapsto ry$. If f is a real and integrable function on S^n we define $\overline{\mathcal{M}}f(y) = \mathcal{M}f(y', h(|y|))$ for $y \in \mathbb{B}$, $y \neq 0$, $y' = y/|y|$.

In Rauch [6] it was proved that

$$u_f^*(y') = \sup_{0 \leq r < 1} |u_f(ry')| \leq C_n \overline{\mathcal{M}}f(y'), \quad y' \in S^n, \quad f \in L^1(S^n) .$$

The inequality in the following lemma generalizes the above inequality.

Lemma 4.1. *There exists a constant $C > 0$ such that, for all real-valued integrable function f on S^n and all $y \in \mathbb{B}$, $0 < |y| < 1$, we have*

$$u_f^*(y) \leq C \overline{\mathcal{M}}f(y) .$$

Proof: We may assume $y = r\mathbb{1} = r(1, 0, \dots, 0)$, $0 \leq r < 1$. Let us denote $\theta = (\theta_1, \dots, \theta_n)$, $\theta' = (\theta_2, \dots, \theta_n)$, $\omega(\theta') = \sin^{n-2} \theta_2 \cdots \sin \theta_{n-1}$ and

$$(4.1) \quad p(\theta_1, r) = P_{r\mathbb{1}}(\xi(\theta)) = \frac{1}{\omega_n} \frac{1 - r^2}{(1 - 2r \cos \theta_1 + r^2)^{(n+1)/2}} .$$

Then

$$u_f(r\mathbb{1}) = \int_0^\pi d\theta_1 \cdots \int_0^{2\pi} p(\theta_1, r) f(\xi(\theta)) \sin^{n-1} \theta_1 \omega(\theta') d\theta_n .$$

If $0 \leq r \leq 1/2$, we have that $p(\theta_1, r) \leq 2^{n+1}/\omega_n$ and hence

$$|u_f(r\mathbb{1})| \leq \frac{2^{n+1}}{\omega_n} \int_{S^n} |f(x)| d\sigma(x) \leq 2^{n+1} \overline{\mathcal{M}} f(r\mathbb{1}) .$$

Now, let us suppose $1/2 \leq r < 1$. If $m(r) = \arccos r(2-r)$, then, integrating by parts with respect to θ_1 , we obtain

$$\begin{aligned} I_r &= \left| \int_{S^n \setminus B(\mathbb{1}, h(r))} P_{r\mathbb{1}}(x) f(x) d\sigma(x) \right| \\ &\leq p(\pi, r) \int_0^\pi d\theta_1 \cdots \int_0^{2\pi} |f(\xi(\theta))| \sin^{n-1} \theta_1 \omega(\theta') d\theta_n \\ &\quad + p(m(r), r) \int_0^{m(r)} d\theta_1 \int_0^\pi d\theta_2 \cdots \int_0^{2\pi} |f(\xi(\theta))| \sin^{n-1} \theta_1 \omega(\theta') d\theta_n \\ &\quad + \int_{m(r)}^\pi \left| \frac{\partial p(\theta_1, r)}{\partial \theta_1} \right| \left[\int_0^{\theta_1} \left(\int_0^\pi d\theta_2 \cdots \int_0^{2\pi} |f(\xi(t, \theta'))| \sin^{n-1} t \omega(\theta') d\theta_n \right) dt \right] d\theta_1 \\ &= I_r^1 + I_r^2 + I_r^3 . \end{aligned}$$

We have that

$$I_r^1 = \frac{1}{\omega_n} \frac{1-r}{(1+r)^n} \int_{S^n} |f(x)| d\sigma(x) \leq \overline{\mathcal{M}} f(r\mathbb{1}) .$$

Since

$$\frac{\omega_{n-1}}{n 2^{n-1}} (1-r)^n \leq \sigma(B(\mathbb{1}, h(r))) \leq \frac{2^n \omega_{n-1}}{n} (1-r)^n$$

then for $1/2 \leq r < 1$, it follows that

$$I_r^2 \leq \frac{2}{\omega_n (1-r)^n} \int_{B(\mathbb{1}, h(r))} |f(x)| d\sigma(x) \leq \frac{2^{n+1} \omega_{n-1}}{n \omega_n} \overline{\mathcal{M}} f(r\mathbb{1}) .$$

Using properties of the Poisson kernel and integration by parts, we obtain

$$\int_0^\pi \left| \frac{\partial p(\theta_1, r)}{\partial \theta_1} \right| \left(\int_0^{\theta_1} \sin^{n-1} t dt \right) d\theta_1 = \frac{1}{\omega_{n-1}} \left(1 - \frac{1-r}{(1+r)^n} \right) \leq \frac{1}{\omega_{n-1}}$$

and thus

$$I_r^3 \leq \frac{1}{\omega_{n-1}} \overline{\mathcal{M}} f(r\mathbb{1}) .$$

Therefore, there exists a constant $D > 0$, such that

$$I_r \leq I_r^1 + I_r^2 + I_r^3 \leq D \overline{\mathcal{M}} f(r\mathbb{1})$$

for all $1/2 \leq r < 1$. Consequently

$$\begin{aligned} |u_f(r\mathbb{1})| &\leq \frac{2}{\omega_n} \frac{1}{(1-r)^n} \int_{B(\mathbb{1}, h(r))} |f(x)| d\sigma(x) + I_r \\ &\leq \frac{2^{n+1} \omega_{n-1}}{n \omega_n} \frac{1}{\sigma(B(\mathbb{1}, h(r)))} \int_{B(\mathbb{1}, h(r))} |f(x)| d\sigma(x) + D\overline{\mathcal{M}}f(r\mathbb{1}) \\ &\leq \frac{2^{n+1} \omega_{n-1}}{n \omega_n} \overline{\mathcal{M}}f(r\mathbb{1}) + D\overline{\mathcal{M}}f(r\mathbb{1}) = C\overline{\mathcal{M}}f(r\mathbb{1}) . \blacksquare \end{aligned}$$

Proof of Theorem 4.1: The Proof of (i) \Rightarrow (ii) is exactly as the proof of (i) \Rightarrow (ii) in Theorem 2.1 and Theorem 3.1.

Let us prove (ii) \Rightarrow (i). Let f be a real-valued positive integrable function on S^n . There exists a constant $C > 0$, depending only on n , such that

$$P_{ry'}(x) \geq \frac{C}{\sigma(B(y', h(r)))}$$

for all $0 \leq r < 1$, $y' \in S^n$ and $x \in B(y', h(r))$. Therefore

$$u_f(ry') \geq \frac{C}{\sigma(B(y', h(r)))} \int_{B(y', h(r))} f(x) d\sigma(x)$$

and hence

$$(4.2) \quad u_f^*(ry') \geq C\overline{\mathcal{M}}f(ry')$$

Consider the function $k: \mathbb{B} \rightarrow \tilde{S}^n$ defined by $k(x) = (x/|x|, h(|x|))$, $x \neq 0$, $k(0) = (\mathbb{1}, 0)$, $\mathbb{1} = (1, 0, \dots, 0)$. Then applying Theorem 3.1 to $X = S^n$ and to the image measure β of ν by k , $\beta(A) = \nu(k^{-1}(A))$ and using the inequalities (4.1) and (4.2), we obtain the wanted proof. \blacksquare

REFERENCES

- [1] BERNARDIS, A. and SALINAS, O. – Two-weight norm inequalities for the fractional maximal operator on spaces of homogeneous type, *Studia Math.*, 108 (1994), 201–207.
- [2] CARLESON, L. – Interpolation by bounded analytic functions and the corona problem, *Annals of Math.*, 76 (1962), 547–559.
- [3] DELLACHERIE, C. and MEYER, P.A. – *Probabilités et potentiel. Théorie des Martingales*, Hermann, Paris, 1980.
- [4] KORÁNYI, A. and VÁGI, S. – Singular integrals on homogeneous spaces and some problems of classical analysis, *Annali della Scuola Norm. Sup. di Pisa*, 25 (1971), 575–648.

- [5] MACIAS, R. and SEGOVIA, C. – A well behaved quasi-distance for spaces of homogeneous type, *Trabajos de Matemática*, 32, Inst. Argentino de Matemática, 1981.
- [6] RAUCH, H.E. – Harmonic and analytic functions of several variables and the maximal theorem of Hardy and Littlewood, *Canad. J. Math.*, 8 (1956), 171–183.
- [7] RUIZ, F.J. and TORREA, J.L. – A unified approach to Carleson measures and A_p weights II, *Pacific J. Math.*, 120 (1985), 189–197.
- [8] RUIZ, F.J. and TORREA, J.L. – Vector-valued Calderón–Zygmund theory and Carleson measures on spaces of homogeneous nature, *Studia Math.*, 88 (1988), 221–243.
- [9] SAWYER, E. and WHEEDEN, R.L. – Weighted inequalities for fractional integrals on euclidean and homogeneous spaces, *Amer. J. Math.*, 114 (1992), 813–874.

Benjamin Bordin, Iara A.A. Fernandes and Sergio A. Tozoni,
Instituto de Matemática, Universidade Estadual de Campinas,
Caixa Postal 6065, 13.081-970 Campinas - SP – BRAZIL

E-mail: bordin@ime.unicamp.br
iara@ime.unicamp.br
tozoni@ime.unicamp.br