

SOME VARIANTS OF THE EXPONENTIAL FORMULA, WITH APPLICATION TO THE MULTIVARIATE TUTTE POLYNOMIAL (ALIAS POTTS MODEL)

ALEXANDER D. SCOTT

Mathematical Institute, University of Oxford
24–29 St. Giles, Oxford OX1 3LB, England.
E-mail: scott@maths.ox.ac.uk

ALAN D. SOKAL[†]

Department of Physics, New York University
4 Washington Place, New York, NY 10003 USA.
E-mail: sokal@nyu.edu

Dedicated to the memory of Pierre Leroux

ABSTRACT. We prove some variants of the exponential formula and apply them to the multivariate Tutte polynomials (also known as Potts-model partition functions) of graphs. We also prove some further identities for the multivariate Tutte polynomial, which generalize an identity for counting connected graphs found by Riordan, Nijenhuis, Wilf and Kreweras and in more general form by Leroux and Gessel, and an identity for the inversion enumerator of trees found by Mallows, Riordan and Kreweras. Finally, we prove a generalization of Möbius inversion on the partition lattice.

1. INTRODUCTION

Let $\{c_n\}_{n=1}^{\infty}$ be a sequence of coefficients, and let

$$C(x) = \sum_{n=1}^{\infty} c_n \frac{x^n}{n!} \tag{1.1}$$

be the corresponding exponential generating function, considered as a formal power series in the indeterminate x . Now let q be another (commuting) indeterminate¹, and

2000 *Mathematics Subject Classification*. 05A19 (Primary); 05A15, 05A18, 05C15, 05C30, 05C99, 06A07, 82B20 (Secondary).

Key words and phrases. Exponential formula, generating function, sequence of binomial type, convolution family, graph, connected component, Tutte polynomial, Potts model, complete graph, inversion enumerator for trees, Abel's binomial identity, Chu–Vandermonde convolution, Rothe's identity, Möbius inversion, partition lattice.

[†] Also at Department of Mathematics, University College London, London WC1E 6BT, England.

¹Let us stress that this q has nothing to do with the q of q -series; rather, it is connected with the q of the q -state Potts model [44, 68, 69] and of the multivariate Tutte polynomial [57].

define

$$A_q(x) = \exp[qC(x)] = \sum_{n=0}^{\infty} a_n(q) \frac{x^n}{n!}, \quad (1.2)$$

so that $A_q(x) = A(x)^q$ [we write $A(x) = A_1(x)$]. It is easy to see that:

- (a) Each coefficient $a_n(q)$ is a polynomial in q , of degree at most n ; indeed we have the explicit formula²

$$a_n(q) = \sum_{\ell=0}^n \frac{q^\ell}{\ell!} \sum_{\substack{n_1, \dots, n_\ell \geq 1 \\ \sum n_j = n}} \binom{n}{n_1, \dots, n_\ell} \prod_{j=1}^{\ell} c_{n_j}. \quad (1.3)$$

- (b) $a_0(q) = 1$.

- (c) For $n \neq 0$, the polynomial $a_n(q)$ has zero constant term, so that we can write $a_n(q) = q\hat{a}_n(q)$ where $\hat{a}_n(q)$ is a polynomial of degree at most $n - 1$; moreover we have $\hat{a}_n(0) = c_n$.

For some purposes it is convenient to introduce the modified generating function

$$\hat{A}_q(x) = \frac{A_q(x) - 1}{q} = \frac{\exp[qC(x)] - 1}{q} = \sum_{n=1}^{\infty} \hat{a}_n(q) \frac{x^n}{n!}. \quad (1.4)$$

The coefficients $\{a_n(q)\}_{n=0}^{\infty}$ furthermore satisfy the following identities [23]:³

$$\sum_{k=0}^n \binom{n}{k} a_k(q_1) a_{n-k}(q_2) = a_n(q_1 + q_2) \quad (1.5)$$

$$\sum_{k=1}^n \binom{n}{k} k \hat{a}_k(q_1) a_{n-k}(q_2) = n \hat{a}_n(q_1 + q_2) \quad \text{for } n \geq 1 \quad (1.6)$$

Indeed, (1.5) follows from writing the trivial identity

$$A_{q_1}(x)A_{q_2}(x) = A_{q_1+q_2}(x) \quad (1.7)$$

and extracting the coefficient of x^n , while (1.6) follows analogously from

$$A_{q_2}(x) \left[x \frac{d}{dx} \frac{A_{q_1}(x) - 1}{q_1} \right] = x \frac{d}{dx} \frac{A_{q_1+q_2}(x) - 1}{q_1 + q_2} \quad (1.8)$$

[23, eq. (4.3)], which is an easy consequence of (1.2). Specializing (1.6) to $q_1 = 0$, or alternatively applying $x \frac{d}{dx}$ directly to (1.2), we obtain the well-known recursion relation [67, Theorem 3.10.1]

$$a_n(q) = q \sum_{k=1}^n \binom{n-1}{k-1} c_k a_{n-k}(q) \quad \text{for } n \geq 1, \quad (1.9)$$

²This formula holds also for $n = 0$ because the integer 0 has exactly one ordered partition (n_1, \dots, n_ℓ) into zero or more strictly positive parts: namely, we have $\ell = 0$ parts, and the multinomial coefficient and the empty product both equal 1, correctly giving $a_0(q) = 1$.

³See Gould [23] for fascinating comments on the history of identities equivalent to (1.6) [or to (1.10) below], which apparently go back to Euler (1748), Rothe (1793), Hindenburg (1796), von Ettingshausen (1826) and Hanstead (1881), among others.

which allows the $\{a_n(q)\}$ to be calculated given the $\{c_n\}$, or vice versa. More generally, we can form the linear combination $n \times (1.5) - (q_1 + q_2) \times (1.6)$ to obtain the recursion relation⁴

$$a_n(q_2) = \sum_{k=1}^n \left[\binom{n-1}{k-1} (q_1 + q_2) - \binom{n}{k} q_1 \right] \widehat{a}_k(q_1) a_{n-k}(q_2) \quad (1.10a)$$

$$= \sum_{k=1}^n \left[\binom{n-1}{k-1} q_2 - \binom{n-1}{k} q_1 \right] \widehat{a}_k(q_1) a_{n-k}(q_2) \quad (1.10b)$$

for $n \geq 1$, which allows the $\{a_n(q_2)\}$ to be calculated given the $\{\widehat{a}_n(q_1)\}$ for any fixed q_1 , and which reduces to (1.9) when $q_1 = 0$. We do not know whether there exist useful analogues of (1.5)/(1.6) with higher powers of k on the left-hand side.

Combinatorially, all these identities are versions of the *exponential formula* [5, 13, 14, 60, 64, 67], which relates the weights c_n of “connected” objects to the weights $a_n(q)$ of “all” objects, when the weight of an object is defined to be the product of the weights of its “connected components” times a factor q for each connected component.⁵ This idea can be formalized in the following combinatorial model: Let Π_n be the set of partitions of $[n] \equiv \{1, \dots, n\}$ into (zero or more) nonempty blocks; give a partition $\pi = \{\pi_1, \dots, \pi_\ell\}$ a weight $w(\pi) = q^\ell \prod_{i=1}^{\ell} c_{|\pi_i|}$, where $\{c_k\}_{k=1}^{\infty}$ are arbitrary coefficients⁶; and let $a_n(q) = \sum_{\pi \in \Pi_n} w(\pi)$. Then (1.3) is immediate: the multinomial coefficient counts the *ordered* partitions of $[n]$ into blocks of sizes n_1, \dots, n_ℓ , and the $1/\ell!$ converts this to *unordered* partitions. As for (1.5)/(1.6), we first observe that $a_n(q_1 + q_2)$ can be computed by allowing each block to be colored either “1” (with weight q_1) or “2” (with weight q_2). Identity (1.5) follows by defining $S \subseteq [n]$ to be the set of elements colored “1”, and setting $k = |S|$. Analogously, $\widehat{a}_n(q_1 + q_2)$ can be computed by coloring and weighting blocks as before, except that the block containing one specified element $i \in [n]$ (say, $i = 1$) must be colored “1” and it receives weight 1 instead of weight q_1 . Defining again $S \subseteq [n]$ to be the set of elements colored “1”, and setting $k = |S|$, we

⁴See [23, eq. (1.3)] or [59, Exercise 1.39, pp. 50 and 62] for the case $q_1 = 1$, which is of course trivially equivalent to the general case of $q_1 \neq 0$.

⁵In the *theory of combinatorial species* [5], the exponential formula is merely a special case of the more general concept of *composition* (or *substitution*) of species $F \circ G$ [5, Section 1.4], which is defined so that the exponential generating function of $F \circ G$ is the composition of the exponential generating functions of F and G . When F is taken to be the species E of sets (*ensembles* in French) and H is some species for which there exists a reasonable notion of connectedness, then, letting H^c be the subspecies of connected H -structures, we have the isomorphism of species $H = E \circ H^c$; since the exponential generating function of E is \exp , this immediately implies the exponential formula $H(x) = \exp(H^c(x))$. This result also generalizes to weighted and multi-sort species. See [34] for results closely related to those of the present paper, formulated in the language of species.

⁶The coefficients c_k may in turn be generating functions for some additional structure on the set $[k]$ — for instance, a graph, a digraph, a hypergraph, etc. — under the condition that this structure is “connected”.

obtain (1.6) in the form⁷

$$\widehat{a}_n(q_1 + q_2) = \sum_{k=1}^n \binom{n-1}{k-1} \widehat{a}_k(q_1) a_{n-k}(q_2) \quad \text{for } n \geq 1. \quad (1.11)$$

Sequences of polynomials $\{a_n(q)\}$ satisfying (1.2) [or equivalently satisfying (1.5) and not identically zero] have been termed *sequences of binomial type* by Rota and collaborators [12, 16, 39, 48, 49, 51] and studied by means of the umbral calculus [18, 49, 52].⁸ The corresponding sequences $\{a_n(q)/n!\}$ have been termed sequences of binomial type by Labelle [32] and *convolution families* by Knuth [30] and Zeng [70]; these authors used elementary formal-power-series methods closely resembling those used here. A purely combinatorial approach to sequences of binomial type, employing the theory of species, has been developed by Labelle [33] (see also [5, Section 3.1]).

Further identities, generalizing (1.5)/(1.6), can be obtained by a powerful transformation suggested by Knuth [30]. Start with any formal power series $A(x)$ with constant term 1; set $A_q(x) = A(x)^q$ and define polynomials $\{a_n(q)\}$ by (1.2). Now let t be a parameter, and define a formal power series $A(x; t)$ by the implicit equation

$$A(x; t) = A(xA(x; t)^t). \quad (1.12)$$

Since $A(x; 0) = A(x)$, we can view the family $\{A(x; t)\}_{t \in \mathbb{R}}$ as a special class of “perturbations” of $A(x)$. Now define polynomials $\{a_n(q; t)\}$ by the obvious formula

$$A(x; t)^q = \sum_{n=0}^{\infty} a_n(q; t) \frac{x^n}{n!}. \quad (1.13)$$

For each t , these polynomials form a family of the type (1.2) and thus satisfy the identities (1.5)/(1.6). On the other hand, a straightforward calculation using the Lagrange inversion formula [60] yields the remarkable relation⁹

$$a_n(q; t) = \frac{q}{q + nt} a_n(q + nt) = q \widehat{a}_n(q + nt), \quad (1.14)$$

or even more simply

$$\widehat{a}_n(q; t) = \widehat{a}_n(q + nt). \quad (1.15)$$

[In particular, $a_n(q; t)$ and $\widehat{a}_n(q; t)$ are *polynomials* jointly in q and t , so we can treat t as an indeterminate if we wish.] It follows that, for any family $\{a_n(q)\}$ of type (1.2),

⁷See also [45, Theorem 2.9]; and see [5, p. 188, Exercise 3.1.20(a)] for the special case $q_1 = 1$.

⁸It is not hard to see by induction that if $\{a_n(q)\}$ satisfies (1.5) [or even the special case of (1.5) with $q_1 = q_2$] and is not identically zero, then $\deg a_n \leq n$ with $a_0 = 1$ and $a_n(0) = 0$ for $n \geq 1$; moreover, with a little extra work it can be shown that $A_q(x)$ is necessarily of the form (1.2): see [16, Theorem 4.1] [30, 70] [60, Exercise 5.37, pp. 87–88 and 131–132].

Let us also remark that Rota *et al.* [39, 51] and many subsequent authors [12, 16, 48, 49], in defining “sequence of binomial type”, impose the additional condition that $\deg a_n = n$ exactly (i.e., $c_1 \neq 0$); but since this condition is irrelevant for our purposes, we prefer not to impose it.

⁹See [60, Exercises 5.37(e) and 5.58, pp. 87–88, 99, 131–133 and 148], where the argument is attributed to Eric Rains and Ira Gessel. Please note also that if we write $f(x) = x/A(x)^t$ and $g(x) = xA(x; t)^t$, then f and g are compositional inverses (as observed in [30, 32] for $t = 1$).

we have the following identities generalizing (1.5)/(1.6):¹⁰

$$\sum_{k=0}^n \binom{n}{k} q_1 \widehat{a}_k(q_1 + kt) q_2 \widehat{a}_{n-k}(q_2 + (n-k)t) = (q_1 + q_2) \widehat{a}_n(q_1 + q_2 + nt) \quad (1.16)$$

$$\sum_{k=1}^n \binom{n}{k} k \widehat{a}_k(q_1 + kt) q_2 \widehat{a}_{n-k}(q_2 + (n-k)t) = n \widehat{a}_n(q_1 + q_2 + nt) \quad \text{for } n \geq 1 \quad (1.17)$$

where we understand that $q \widehat{a}_0(q) = 1$. Multiplying (1.17) by t and summing it with (1.16), we obtain the alternate form

$$\sum_{k=0}^n \binom{n}{k} a_k(q_1 + kt) q_2 \widehat{a}_{n-k}(q_2 + (n-k)t) = a_n(q_1 + q_2 + nt). \quad (1.18)$$

If in (1.18) we interchange $k \leftrightarrow n-k$ and $q_1 \leftrightarrow q_2$ and then replace q_2 by $q_2 - nt$, we obtain the slightly simpler form

$$\sum_{k=0}^n \binom{n}{k} q_1 \widehat{a}_k(q_1 + kt) a_{n-k}(q_2 - kt) = a_n(q_1 + q_2). \quad (1.19)$$

A combinatorial proof and interpretation of (1.16)/(1.18) in terms of the theory of species has been given by Labelle [33].¹¹ We call (1.16)–(1.19) *Abel-type extensions* of (1.5)/(1.6) because in the simplest case $A(x) = e^x$, $a_n(q) = q^n$ they reduce to Abel's celebrated extensions of the binomial theorem (also discovered simultaneously by Cauchy) [28, 46, 61].

These Abel-type formulae can alternatively be derived as immediate consequences of the Pfaff–Cauchy derivative identities (generalizations of Leibniz's rule for the n th derivative of the product of two functions) recently presented by Johnson [28]. Indeed, setting $v = A(x)^{q_1}$, $w = A(x)^{q_2}$, $\phi = A(x)^t$ in [28, eq. (1.1)] yields (1.19), while making the same substitution in [28, eq. (1.4)] yields (1.16).

Some classical examples of families $\{a_n(q)\}$ and the corresponding identities (1.5)/(1.6) and (1.16)–(1.19) are listed in Table 1.

In this note we begin by proving some easy generalizations of the foregoing formulae to the case in which the integer index n is replaced by a multi-index \mathbf{n} over a finite ground set V . We then show some applications of the latter formulae to the multivariate Tutte polynomial [57] of a graph $G = (V, E)$. Finally, we derive some further identities for the multivariate Tutte polynomial, which generalize an identity for counting connected graphs found by Riordan, Nijenhuis, Wilf and Kreweras [31, 40, 47] and in more general form by Leroux [37] and Gessel [17] and an identity for the inversion enumerator of trees found by Mallows, Riordan and Kreweras [31, 38]. In the appendix we prove an apparently new generalization of Möbius inversion on the partition lattice.

¹⁰Already Rota *et al.* [51, p. 711, Proposition 4] and Reiner [45, Theorem 2.10] noticed that, for any t , the family $\{a_n(q; t)\}$ defined by (1.14) is of type (1.2). But they do not seem to have noticed the relation (1.12), which to our knowledge goes back to Knuth [30].

¹¹See also [5, p. 189, Exercise 3.1.22] for a brief summary.

$A(x)$	$a_n(q)$	(1.5)/(1.6)	(1.16)–(1.19)
e^x	q^n	binomial theorem	Abel–Cauchy binomial identities
$1+x$	q^n	Chu–Vandermonde convolution	Rothe’s generalized Chu–Vandermonde convolution
$(1-x)^{-1}$	$q^{\bar{n}}$		
$(1+\alpha x)^\beta$	$\alpha^n (\beta q)^{\underline{n}}$		
$\exp(e^x - 1)$	$B_n(q) = \sum_{k=0}^n \{n\}_k q^k$ (Bell polynomial [11])		
$\exp[-x/(1-x)]$	$L_n^{(-1)}(q) = \sum_{k=0}^n \frac{n!}{k!} \binom{n-1}{n-k} (-q)^k$ (Laguerre polynomial [1])		

TABLE 1. Some classical identities that are special cases of (1.5)/(1.6) and (1.16)–(1.19). See [21–24, 28, 46, 61] for more information concerning Abel–Cauchy and Chu–Vandermonde–Rothe identities. The identity (1.5) is well known for the Bell [11, p. 136, eq. 3n] and Laguerre [2, p. 192, eq. 41] [36, p. 96, Problem 20] polynomials. We do not know any references for (1.6) or (1.16)–(1.19) in these cases.

We stress that many (though not all) of the formulae derived in this paper are already known. But we think that our approach provides a unified perspective that may be of interest.

2. VARIANTS OF THE EXPONENTIAL FORMULA

Let V be a finite ground set; we shall use multi-indices $\mathbf{n} = (n_i)_{i \in V} \in \mathbb{N}^V$ and commuting indeterminates $\mathbf{x} = (x_i)_{i \in V}$. We write $|\mathbf{n}| = \sum_{i \in V} n_i$ and $\mathbf{x}^{\mathbf{n}} = \prod_{i \in V} x_i^{n_i}$. We use the notation $\mathbf{n}! = \prod_{i \in V} n_i!$ and analogously for binomial and multinomial coefficients. Finally, we denote by $\mathbf{0}$ (respectively $\mathbf{1}$) the multi-index with all entries 0 (respectively 1); and for any subset $W \subseteq V$, we denote by $\mathbf{1}_W$ the multi-index taking the value 1 on W and 0 on $V \setminus W$.

Let R be a commutative ring containing the rationals as a subring; we shall use polynomials and formal power series whose coefficients lie in R . (In applications, R will usually be either the rationals, reals or complex numbers, or a ring of polynomials or formal power series over one of these fields.) One possible approach is to treat the coefficients $\mathbf{c} = (c_{\mathbf{n}})_{\mathbf{n} \in \mathbb{N}^V \setminus \{\mathbf{0}\}}$ as indeterminates, and to take R to be the polynomial ring $\mathbb{Q}[\mathbf{c}]$; this approach has the advantage of exhibiting the polynomial dependence on the coefficients $c_{\mathbf{n}}$. But we need not commit ourselves to any specific choice of the ring R ; all of our identities will be valid in complete generality.

So let $\{c_{\mathbf{n}}\}_{\mathbf{n} \in \mathbb{N}^V}$ be a sequence of coefficients in the ring R , with $c_{\mathbf{0}} = 0$; and let

$$C(\mathbf{x}) = \sum_{\mathbf{n}} c_{\mathbf{n}} \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \quad (2.1)$$

be the corresponding exponential generating function, considered as a formal power series in the indeterminates \mathbf{x} . Now let q be another (commuting) indeterminate, and define

$$A_q(\mathbf{x}) = \exp[qC(\mathbf{x})] = \sum_{\mathbf{n}} a_{\mathbf{n}}(q) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!}. \quad (2.2)$$

Using the Taylor series for \exp , we deduce immediately that

- (a) Each coefficient $a_{\mathbf{n}}(q)$ is a polynomial in q , of degree at most $|\mathbf{n}|$; indeed we have the explicit formula¹²

$$a_{\mathbf{n}}(q) = \sum_{\ell=0}^{|\mathbf{n}|} \frac{q^{\ell}}{\ell!} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_{\ell} \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_{\ell}} \prod_{j=1}^{\ell} c_{\mathbf{n}_j}. \quad (2.3)$$

- (b) $a_{\mathbf{0}}(q) = 1$.

- (c) For $\mathbf{n} \neq \mathbf{0}$, the polynomial $a_{\mathbf{n}}(q)$ has zero constant term, so that we can write $a_{\mathbf{n}}(q) = q\widehat{a}_{\mathbf{n}}(q)$ where $\widehat{a}_{\mathbf{n}}(q)$ is a polynomial of degree at most $|\mathbf{n}| - 1$; moreover we have $\widehat{a}_{\mathbf{n}}(0) = c_{\mathbf{n}}$.

Conversely, it is not hard to see that if $(c_{\mathbf{n}})_{\mathbf{n} \in \mathbb{N}^V \setminus \{\mathbf{0}\}}$ is any sequence and we define $(a_{\mathbf{n}}(q))_{\mathbf{n} \in \mathbb{N}^V}$ by (2.3), then the latter family satisfies (2.2).

Using $qC(\mathbf{x}) = \log A_q(\mathbf{x})$ together with the Taylor series for $\log(1+z)$, we obtain the inverse formula

$$qc_{\mathbf{n}} = \sum_{\ell=1}^{|\mathbf{n}|} \frac{(-1)^{\ell-1}}{\ell} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_{\ell} \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_{\ell}} \prod_{j=1}^{\ell} a_{\mathbf{n}_j}(q) \quad (2.4)$$

or equivalently

$$c_{\mathbf{n}} = \sum_{\ell=1}^{|\mathbf{n}|} \frac{(-q)^{\ell-1}}{\ell} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_{\ell} \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_{\ell}} \prod_{j=1}^{\ell} \widehat{a}_{\mathbf{n}_j}(q). \quad (2.5)$$

It is a nontrivial fact that the right-hand side of (2.5) is independent of q .

Now let r be another indeterminate, and let us use the identity $A_{rq}(\mathbf{x}) = A_q(\mathbf{x})^r$ together with the Taylor series

$$(1+z)^r \equiv \exp[r \log(1+z)] = \sum_{\ell=0}^{\infty} \frac{r^{\ell}}{\ell!} z^{\ell} \quad (2.6)$$

¹²This formula holds also for $\mathbf{n} = \mathbf{0}$ because the multi-index $\mathbf{0}$ has exactly one ordered partition $(\mathbf{n}_1, \dots, \mathbf{n}_{\ell})$ into zero or more nonvanishing parts: namely, we have $\ell = 0$ parts, and the multinomial coefficient and the empty product both equal 1, correctly giving $a_{\mathbf{0}}(q) = 1$. This same observation applies to all subsequent formulae of a similar type, such as (2.7) ff.

where $r^\ell = r(r-1)\cdots(r-\ell+1)$ denotes the falling factorial. (If we like, we can alternatively use the notation $\binom{r}{\ell} = r^\ell/\ell!$, which is a well-defined polynomial in the indeterminate r .) We obtain

$$a_{\mathbf{n}}(rq) = \sum_{\ell=0}^{|\mathbf{n}|} \frac{r^\ell}{\ell!} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} a_{\mathbf{n}_j}(q). \quad (2.7)$$

In view of the identity $(-r)^\ell = (-1)^\ell r^{\bar{\ell}}$ where $r^{\bar{\ell}} = r(r+1)\cdots(r+\ell-1) = (r+\ell-1)^\ell$ denotes the rising factorial, we also have

$$a_{\mathbf{n}}(-rq) = \sum_{\ell=0}^{|\mathbf{n}|} \frac{(-1)^\ell r^{\bar{\ell}}}{\ell!} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} a_{\mathbf{n}_j}(q). \quad (2.8)$$

Since both sides of the identities (2.7)/(2.8) are polynomials in r , we can specialize these identities to r integer or rational, or more generally to r being any element of the ring R . If r is a positive integer, the sum in (2.7) can obviously be restricted to $\ell \leq r$. Slightly (but not much) less trivially, if r is a positive integer, then by using

$$\frac{r^{\bar{\ell}}}{\ell!} = \frac{(r+\ell-1)!}{\ell!(r-1)!} = \frac{(\ell+1)^{\overline{r-1}}}{(r-1)!} \quad (2.9)$$

we can rewrite (2.8) as

$$a_{\mathbf{n}}(-rq) = \sum_{\ell=0}^{|\mathbf{n}|} (-1)^\ell \frac{(\ell+1)^{\overline{r-1}}}{(r-1)!} \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} a_{\mathbf{n}_j}(q), \quad (2.10)$$

in which the coefficient $(\ell+1)^{\overline{r-1}}/(r-1)!$ is a polynomial of degree $r-1$ in ℓ . In particular, for $r=1$ we have

$$a_{\mathbf{n}}(-q) = \sum_{\ell=0}^{|\mathbf{n}|} (-1)^\ell \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} a_{\mathbf{n}_j}(q). \quad (2.11)$$

By formally setting $r = q_2/q_1$ (this is justifiable if we work in a ring of formal Laurent series in the indeterminate q_1), we obtain the identity

$$a_{\mathbf{n}}(q_2) = \sum_{\ell=0}^{|\mathbf{n}|} \frac{1}{\ell!} \left(\prod_{j=0}^{\ell-1} (q_2 - jq_1) \right) \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} \widehat{a}_{\mathbf{n}_j}(q_1), \quad (2.12)$$

which is nice because both sides are manifestly polynomials in the indeterminates q_1 and q_2 . Alternatively, for $\mathbf{n} \neq \mathbf{0}$ we can pull out the $j=0$ factor and write

$$\widehat{a}_{\mathbf{n}}(q_2) = \sum_{\ell=1}^{|\mathbf{n}|} \frac{1}{\ell!} \left(\prod_{j=1}^{\ell-1} (q_2 - jq_1) \right) \sum_{\substack{\mathbf{n}_1, \dots, \mathbf{n}_\ell \neq \mathbf{0} \\ \sum \mathbf{n}_j = \mathbf{n}}} \binom{\mathbf{n}}{\mathbf{n}_1, \dots, \mathbf{n}_\ell} \prod_{j=1}^{\ell} \widehat{a}_{\mathbf{n}_j}(q_1). \quad (2.13)$$

Conversely, it is not hard to see that if q_1 is any element of the ring R and $(\widehat{a}_{\mathbf{n}}(q_1))_{\mathbf{n} \in \mathbb{N}^V}$ is any sequence, and we define $(a_{\mathbf{n}}(q))_{\mathbf{n} \in \mathbb{N}^V}$ by (2.12), then the latter family satisfies (2.2).

The following special cases of (2.12)/(2.13) are of particular interest:

- (a) For $q_1 = 0$, the identity (2.12) reduces to (2.3).
- (b) For $q_2 = 0$, we obtain the inverse formula (2.4)/(2.5).
- (c) For $q_2 = -q_1$, we obtain the closely related formula (2.11).

It is easy to show that the coefficients $\{a_{\mathbf{n}}(q)\}_{\mathbf{n} \in \mathbb{N}^V}$ satisfy the identities

$$\sum_{\mathbf{k}} \binom{\mathbf{n}}{\mathbf{k}} a_{\mathbf{k}}(q_1) a_{\mathbf{n}-\mathbf{k}}(q_2) = a_{\mathbf{n}}(q_1 + q_2) \quad (2.14)$$

$$\sum_{\mathbf{k} \neq \mathbf{0}} \binom{\mathbf{n}}{\mathbf{k}} k_i \widehat{a}_{\mathbf{k}}(q_1) a_{\mathbf{n}-\mathbf{k}}(q_2) = n_i \widehat{a}_{\mathbf{n}}(q_1 + q_2) \quad \text{for } \mathbf{n} \neq \mathbf{0} \quad (2.15)$$

Indeed, (2.14) follows by expanding out the identity $A_{q_1}(\mathbf{x})A_{q_2}(\mathbf{x}) = A_{q_1+q_2}(\mathbf{x})$ and extracting the coefficient of $\mathbf{x}^{\mathbf{n}}$, while (2.15) follows analogously from

$$A_{q_2}(\mathbf{x}) \left[x_i \frac{\partial}{\partial x_i} \frac{A_{q_1}(\mathbf{x}) - 1}{q_1} \right] = x_i \frac{\partial}{\partial x_i} \frac{A_{q_1+q_2}(\mathbf{x}) - 1}{q_1 + q_2}, \quad (2.16)$$

which is an easy consequence of (2.2). In particular, specializing (2.15) to $q_1 = 0$, we obtain the recursion relation

$$a_{\mathbf{n}}(q) = q \sum_{\mathbf{k} \geq \boldsymbol{\delta}_i} \binom{\mathbf{n} - \boldsymbol{\delta}_i}{\mathbf{k} - \boldsymbol{\delta}_i} c_{\mathbf{k}} a_{\mathbf{n}-\mathbf{k}}(q) \quad \text{whenever } n_i \geq 1, \quad (2.17)$$

which allows the $\{a_{\mathbf{n}}(q)\}$ to be calculated given the $\{c_{\mathbf{n}}\}$, or vice versa. [Here $\boldsymbol{\delta}_i$ is the vector with entry 1 at element i and 0 elsewhere.] More generally, we can form the linear combination $n_i \times (2.14) - (q_1 + q_2) \times (2.15)$ to obtain the recursion relation

$$a_{\mathbf{n}}(q_2) = \sum_{\mathbf{k} \neq \mathbf{0}} \left[\binom{\mathbf{n} - \boldsymbol{\delta}_i}{\mathbf{k} - \boldsymbol{\delta}_i} (q_1 + q_2) - \binom{\mathbf{n}}{\mathbf{k}} q_1 \right] \widehat{a}_{\mathbf{k}}(q_1) a_{\mathbf{n}-\mathbf{k}}(q_2) \quad \text{whenever } n_i \geq 1, \quad (2.18)$$

which allows the $\{a_{\mathbf{n}}(q_2)\}$ to be calculated given the $\{\widehat{a}_{\mathbf{n}}(q_1)\}$ for any fixed q_1 , and which reduces to (2.17) when $q_1 = 0$.

We do not know whether there exist useful analogues of (2.14)/(2.15) with higher powers of \mathbf{k} on the left-hand side.

Finally, we can deduce the Abel-type extensions of (2.14)/(2.15):

$$\sum_{\mathbf{k}} \binom{\mathbf{n}}{\mathbf{k}} q_1 \widehat{a}_{\mathbf{k}}(q_1 + \mathbf{k} \cdot \mathbf{t}) q_2 \widehat{a}_{\mathbf{n}-\mathbf{k}}(q_2 + (\mathbf{n} - \mathbf{k}) \cdot \mathbf{t}) = (q_1 + q_2) \widehat{a}_{\mathbf{n}}(q_1 + q_2 + \mathbf{n} \cdot \mathbf{t}) \quad (2.19)$$

$$\sum_{\mathbf{k} \neq \mathbf{0}} \binom{\mathbf{n}}{\mathbf{k}} k_i \widehat{a}_{\mathbf{k}}(q_1 + \mathbf{k} \cdot \mathbf{t}) q_2 \widehat{a}_{\mathbf{n}-\mathbf{k}}(q_2 + (\mathbf{n} - \mathbf{k}) \cdot \mathbf{t}) = n_i \widehat{a}_{\mathbf{n}}(q_1 + q_2 + \mathbf{n} \cdot \mathbf{t}) \quad \text{for } \mathbf{n} \neq \mathbf{0} \quad (2.20)$$

where $\mathbf{t} = (t_i)_{i \in V}$ are commuting indeterminates and we understand $q \widehat{a}_{\mathbf{0}}(q) = 1$. These formulae can be proved by repeated use of the Lagrange inversion formula,

once in each variable. For details we refer to the paper of Zeng [70, pp. 225–226]. For further information on classical specializations of (2.19)/(2.20), see [42, 43, 58, 61, 70].

Remark. We wonder whether (2.19)/(2.20) might have an extension to identities involving a not-necessarily-diagonal *matrix* $T = (t_{ij})_{i,j \in V}$ of indeterminates, based on considering the implicit equation

$$e^{C(\mathbf{x};T)} = A(e^{C(\mathbf{x};T)T} \mathbf{x}) , \quad (2.21)$$

which generalizes (1.12). If true, this could be a powerful extension. See Strehl [61] for possibly related work in which a matrix of indeterminates is employed.

3. APPLICATION TO THE MULTIVARIATE TUTTE POLYNOMIAL

In this section we apply the formulae of Section 2 to deduce some identities for multivariate Tutte polynomials.

3.1. Definitions and basic properties. Let $G = (V, E)$ be a finite undirected graph with vertex set $V \neq \emptyset$ and edge set E ; loops and multiple edges are allowed unless explicitly stated otherwise. Then the *multivariate Tutte polynomial* [57] of G is, by definition, the polynomial

$$Z_G(q, \mathbf{v}) = \sum_{A \subseteq E} q^{k(A)} \prod_{e \in A} v_e , \quad (3.1)$$

where q and $\mathbf{v} = (v_e)_{e \in E}$ are commuting indeterminates, and $k(A)$ denotes the number of connected components in the subgraph (V, A) . It is also convenient to pull out one factor of q by defining

$$\widehat{Z}_G(q, \mathbf{v}) = \sum_{A \subseteq E} q^{k(A)-1} \prod_{e \in A} v_e ; \quad (3.2)$$

this is still a polynomial since $k(A) \geq 1$ for all A . Finally, let us make the convention that if $G = \emptyset$ (the graph with empty vertex set and empty edge set), then $Z_\emptyset = 1$ and \widehat{Z}_\emptyset is undefined.

If we specialize to $v_e = v$ for all edges e , we obtain a two-variable polynomial $Z_G(q, v)$ that is essentially equivalent to the classical Tutte polynomial [57, Section 2.5].

Note that at $q = 1$ we have the trivial formula

$$Z_G(1, \mathbf{v}) = \widehat{Z}_G(1, \mathbf{v}) = \prod_{e \in E} (1 + v_e) . \quad (3.3)$$

More interestingly, at $q = 0$ we have the specialization

$$\widehat{Z}_G(0, \mathbf{v}) = C_G(\mathbf{v}) \equiv \sum_{\substack{A \subseteq E \\ k(A) = 1}} \prod_{e \in A} v_e , \quad (3.4)$$

i.e., the generating polynomial of connected spanning subgraphs of G .

In statistical physics, $Z_G(q, \mathbf{v})$ is known as the partition function of the q -state Potts model [44, 68, 69] in the Fortuin–Kasteleyn representation [15, 26, 29]. This arises by virtue of the following identity [15, 29]:

Theorem 3.1 (Fortuin–Kasteleyn representation of the Potts model [15, 29]). *For integer $q \geq 1$, we have*

$$Z_G(q, \mathbf{v}) = \sum_{\sigma: V \rightarrow [q]} \prod_{e=ij \in E} \left[1 + v_e \delta(\sigma(i), \sigma(j)) \right] \quad (3.5)$$

where δ denotes the Kronecker delta.

Proof. On the right-hand side of (3.5), expand out the product over $e \in E$, and let $A \subseteq E$ be the set of edges for which the term $v_e \delta(\sigma(i), \sigma(j))$ is taken. Now perform the sum over maps $\sigma: V \rightarrow [q]$: in each connected component of the subgraph (V, A) the “color” $\sigma(i)$ must be constant, and there are no other constraints. Therefore, the right-hand side equals

$$\sum_{A \subseteq E} q^{k(A)} \prod_{e \in A} v_e, \quad (3.6)$$

as was to be proved. □

Specializing to $v_e = -1$ for all edges e , we obtain the Birkhoff–Whitney [4, 66] expansion for the chromatic polynomial of G :

Corollary 3.2. *For integer $q \geq 1$, the number of proper q -colorings of G is $P_G(q) \equiv Z_G(q, -1)$.*

3.2. Some generating functions. For the remainder of this section, let us assume that G is a *loopless* graph. For notational simplicity it is convenient to assume also that G has no multiple edges; the trivial changes to allow multiple edges can be left to the reader.

For any multi-index $\mathbf{n} \in \mathbb{N}^V$, let us define $G[\mathbf{n}]$ to be the graph obtained from G by expanding each vertex i to an independent set consisting of n_i vertices. That is, the vertices of $G[\mathbf{n}]$ are pairs $\langle i, \alpha \rangle$ with $i \in V$ and $\alpha \in [n_i]$, and the edges of $G[\mathbf{n}]$ are all pairs $\langle \langle i, \alpha \rangle, \langle j, \beta \rangle \rangle$ with $ij \in E$, $\alpha \in [n_i]$ and $\beta \in [n_j]$. Given a set of weights $\mathbf{v} = (v_e)_{e \in E}$ associated to the edges of G , we assign the weight v_{ij} to each edge $\langle \langle i, \alpha \rangle, \langle j, \beta \rangle \rangle$ in $G[\mathbf{n}]$.

In a similar way, let us define $G'[\mathbf{n}]$ to be the graph obtained from G by expanding each vertex i to a clique consisting of n_i vertices. That is, the vertex set of $G'[\mathbf{n}]$ is the same as that of $G[\mathbf{n}]$, and the edges of $G'[\mathbf{n}]$ consist of those of $G[\mathbf{n}]$ together with all pairs $\langle \langle i, \alpha \rangle, \langle i, \beta \rangle \rangle$ with $i \in V$, $\alpha, \beta \in [n_i]$ and $\alpha \neq \beta$. Given a set of weights $\mathbf{v} = (v_e)_{e \in E}$ associated to the edges of G and another set of weights $\mathbf{w} = (w_i)_{i \in V}$ associated to the vertices of G , we assign the weight v_{ij} to each edge $\langle \langle i, \alpha \rangle, \langle j, \beta \rangle \rangle$ ($i \neq j$) in $G'[\mathbf{n}]$ and the weight w_i to each edge $\langle \langle i, \alpha \rangle, \langle i, \beta \rangle \rangle$ in $G'[\mathbf{n}]$.

Note in particular that if $\mathbf{n} = \mathbf{1}_W$ for some subset $W \subseteq V$, then $G[\mathbf{n}] = G'[\mathbf{n}] = G[W]$, the induced subgraph of G on W .

The following “master formula” generates the multivariate Tutte polynomials (with weights as assigned above) of all the graphs $G[\mathbf{n}]$ and $G'[\mathbf{n}]$:

Theorem 3.3. *Let $G = (V, E)$ be a simple graph, and let $\mathbf{v} = (v_e)_{e \in E}$ and $\mathbf{w} = (w_i)_{i \in V}$ be indeterminates. We then have the following exponential generating functions:*

$$\sum_{\mathbf{n}} Z_{G[\mathbf{n}]}(q, \mathbf{v}) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} = \left(\sum_{\mathbf{n}} \left(\prod_{ij \in E} (1 + v_{ij})^{n_i n_j} \right) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \right)^q \quad (3.7)$$

$$\sum_{\mathbf{n}} Z_{G'[\mathbf{n}]}(q, \mathbf{v}, \mathbf{w}) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} = \left(\sum_{\mathbf{n}} \left(\prod_{ij \in E} (1 + v_{ij})^{n_i n_j} \right) \left(\prod_{i \in V} (1 + w_i)^{n_i(n_i-1)/2} \right) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \right)^q \quad (3.8)$$

where all sums run over $\mathbf{n} \in \mathbb{N}^V$.

We remark that the expression in large parentheses on the right-hand side of (3.7) [respectively (3.8)] is the grand partition function for a lattice gas on the graph G [respectively on the graph G° obtained from G by adjoining a loop at each vertex] in which arbitrary nonnegative integer occupation numbers $\mathbf{n} = (n_i)_{i \in V}$ are allowed, with fugacities \mathbf{x} on the vertices and two-particle Boltzmann weights $1 + v_{ij}$ on edges ij [and two-particle Boltzmann weights $1 + w_i$ on the loops]. See e.g. [53, 54] for definitions concerning lattice gases.

When $G = K_1$ (the graph with one vertex and no edges), (3.8) reduces to the well-known [6, 17, 19, 63, 65] exponential generating function for the Tutte polynomials of the complete graphs K_n .¹³

$$\sum_{n=0}^{\infty} Z_n(q, v) \frac{x^n}{n!} = \left(\sum_{n=0}^{\infty} (1 + v)^{n(n-1)/2} \frac{x^n}{n!} \right)^q. \quad (3.9)$$

When $G = K_2$, (3.7) reduces to the exponential generating function for the Tutte polynomials of the complete bipartite graphs K_{n_1, n_2} ,

$$\sum_{n_1, n_2=0}^{\infty} Z_{n_1, n_2}(q, v) \frac{x^{n_1}}{n_1!} \frac{y^{n_2}}{n_2!} = \left(\sum_{n_1, n_2=0}^{\infty} (1 + v)^{n_1 n_2} \frac{x^{n_1}}{n_1!} \frac{y^{n_2}}{n_2!} \right)^q, \quad (3.10)$$

which in turn specializes for $v = -1$ to the well-known [60, Exercise 5.6, pp. 73 and 107–108] exponential generating function for the chromatic polynomials of the complete bipartite graphs¹⁴,

$$\sum_{n_1, n_2=0}^{\infty} P_{K_{n_1, n_2}}(q) \frac{x^{n_1}}{n_1!} \frac{y^{n_2}}{n_2!} = (e^x + e^y - 1)^q. \quad (3.11)$$

More generally, when $G = K_r$, (3.7) gives the exponential generating function for the Tutte polynomials of the complete r -partite graphs K_{n_1, \dots, n_r} .

Note also that if we only want to know $Z_{G[\mathbf{n}]}(q, \mathbf{v})$ for $\mathbf{n} \leq \mathbf{1}$ — that is, we want to know $Z_{G[W]}(q, \mathbf{v})$ for subsets $W \subseteq V$ — then it suffices to get correct the terms

¹³Unfortunately, this formula is usually written in terms of the classical Tutte polynomial $T_G(x, y)$, which is related to $Z_G(q, v)$ by a change of variables [57, Section 2.5] that obscures the very different roles played by q and v .

¹⁴(3.11) can alternatively be derived from the explicit expression [62] for $P_{K_{n_1, n_2}}(q)$ by using a well-known Stirling-number identity [25, eq. (7.49)] together with the binomial formula.

$\mathbf{n} \leq \mathbf{1}$ on the right-hand side of (3.7); all the other terms can be dropped or altered arbitrarily. It follows that

$$Z_{G[W]}(q, \mathbf{v}) = [\mathbf{x}^{1W}] \left(\sum_{\mathbf{n} \leq \mathbf{1}} \left(\prod_{ij \in E} (1 + v_{ij})^{n_i n_j} \right) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \right)^q. \quad (3.12)$$

Note that the expression in large parentheses is now the grand partition function for a lattice gas with *hard-core self-repulsion*, i.e., at most one atom can occupy each site. In the special case $\mathbf{v} = -\mathbf{1}$, this becomes

$$P_{G[W]}(q) = [\mathbf{x}^{1W}] I_G(\mathbf{x})^q, \quad (3.13)$$

where $I_G(\mathbf{x})$ is the multivariate generating polynomial for *independent sets* of vertices in G . The formula (3.13) was found previously by Lass [35, Proposition 3.1].

Since the formula (3.7) for $G[\mathbf{n}]$ is just a specialization of the formula (3.8) for $G'[\mathbf{n}]$, obtained by taking $\mathbf{w} = \mathbf{0}$, it suffices to prove the latter.

First Proof of Theorem 3.3. Since the coefficient of $\mathbf{x}^{\mathbf{n}}$ on each side of (3.8) is a polynomial in q , it suffices to prove the identity for infinitely many values of q in the ring R ; in particular, it suffices to prove it for all positive integers q . By Theorem 3.1, for integer $q \geq 1$ we have

$$Z_{G'[\mathbf{n}]}(q, \mathbf{v}, \mathbf{w}) = \sum_{\sigma: V(G'[\mathbf{n}]) \rightarrow [q]} \prod_{rs \in E(G'[\mathbf{n}])} \left[1 + v_{rs} \delta(\sigma(r), \sigma(s)) \right] \quad (3.14a)$$

$$\begin{aligned} &= \sum_{\sigma: V(G'[\mathbf{n}]) \rightarrow [q]} \left(\prod_{ij \in E} \prod_{\substack{\alpha \in [n_i] \\ \beta \in [n_j]}} \left[1 + v_{ij} \delta(\sigma(i, \alpha), \sigma(j, \beta)) \right] \right) \\ &\quad \times \left(\prod_{i \in V} \prod_{\substack{\alpha, \beta \in [n_i] \\ \alpha < \beta}} \left[1 + w_i \delta(\sigma(i, \alpha), \sigma(i, \beta)) \right] \right) \end{aligned} \quad (3.14b)$$

Now, because of the symmetries in the construction of $G'[\mathbf{n}]$, the summand in (3.14b) depends only on the numbers $\{n_{i,\tau}\}_{i \in V, \tau \in [q]}$ defined by

$$n_{i,\tau} = \#\{\alpha \in [n_i]: \sigma(i, \alpha) = \tau\}, \quad (3.15)$$

and we have

$$\begin{aligned} Z_{G'[\mathbf{n}]}(q, \mathbf{v}, \mathbf{w}) &= \sum_{\substack{\{n_{i,\tau}\} \\ \sum_{\tau=1}^q n_{i,\tau} = n_i \forall i}} \left(\prod_{i \in V} \binom{n_i}{n_{i,1}, \dots, n_{i,q}} \right) \left(\prod_{ij \in E} \prod_{\tau=1}^q (1 + v_{ij})^{n_{i,\tau} n_{j,\tau}} \right) \\ &\quad \times \left(\prod_{i \in V} \prod_{\tau=1}^q (1 + w_i)^{n_{i,\tau} (n_{i,\tau} - 1) / 2} \right). \end{aligned} \quad (3.16)$$

Now multiply by $\mathbf{x}^{\mathbf{n}}/\mathbf{n}!$ and sum over \mathbf{n} ; the sums over the $n_{i,\tau}$ now become unrestricted, and the sums for the different values of τ decouple; we end up with the

product of q terms each of which is

$$\sum_{\mathbf{n}} \left(\prod_{ij \in E} (1 + v_{ij})^{n_i n_j} \right) \left(\prod_{i \in V} (1 + w_i)^{n_i(n_i-1)/2} \right) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!}. \quad (3.17)$$

□

It is instructive to give an alternate proof of Theorem 3.3 in which q is treated directly as an indeterminate:

Second Proof of Theorem 3.3. We begin by proving the special case (3.9). From the binomial formula we have

$$\begin{aligned} \left(1 + \sum_{n=1}^{\infty} (1+v)^{n(n-1)/2} \frac{x^n}{n!} \right)^q &= 1 + \sum_{b=1}^{\infty} \frac{q^b}{b!} \left(\sum_{n=1}^{\infty} (1+v)^{n(n-1)/2} \frac{x^n}{n!} \right)^b \quad (3.18a) \\ &= 1 + \sum_{b=1}^{\infty} \frac{q^b}{b!} \sum_{n_1, \dots, n_b \geq 1} \frac{x^{\sum n_j}}{\prod n_j!} \prod_{j=1}^b (1+v)^{n_j(n_j-1)/2}. \end{aligned} \quad (3.18b)$$

Now write $n = \sum_{j=1}^b n_j$, and consider n_1, \dots, n_b as the block sizes in an *ordered* partition $\pi = (\pi_1, \dots, \pi_b)$ of $[n]$ into b nonempty blocks. The number of ordered partitions with those block sizes is precisely the multinomial coefficient $n! / \prod_{j=1}^b n_j!$, so

(3.18) is equal to

$$1 + \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{\pi \in \Pi'_n} \frac{q^{|\pi|}}{|\pi|!} \prod_{j=1}^{|\pi|} (1+v)^{|\pi_j|(|\pi_j|-1)/2}, \quad (3.19)$$

where Π'_n denotes the set of ordered partitions of $[n]$, and $|\pi|$ denotes the number of blocks in the partition π . Passing to *unordered* partitions absorbs the factor $1/|\pi|!$ (as these are partitions of a *labelled* set), and we get

$$1 + \sum_{n=1}^{\infty} \frac{x^n}{n!} \sum_{\pi \in \Pi_n} q^{|\pi|} \prod_{j=1}^{|\pi|} (1+v)^{|\pi_j|(|\pi_j|-1)/2}. \quad (3.20)$$

The factor $(1+v)^{|\pi_j|(|\pi_j|-1)/2}$ corresponds to a sum over (simple undirected) graphs on the vertex set π_j , with a weight v for each edge. We can now reinterpret (3.20) as a sum over (simple undirected) graphs H on the vertex set $[n]$, with a weight $v^{|E(H)|}$, together with a sum over partitions $\pi \in \Pi_n$ that are *compatible* with H in the sense that each block of π is a union of (vertex sets of) connected components of H . If H has r connected components, then these can be grouped into b nonempty blocks in $\left\{ \begin{smallmatrix} r \\ b \end{smallmatrix} \right\}$ ways (where $\left\{ \begin{smallmatrix} r \\ b \end{smallmatrix} \right\}$ is a Stirling number of the second kind); and it is well known [25, eq. (6.10)] that

$$\sum_{b=1}^r \left\{ \begin{smallmatrix} r \\ b \end{smallmatrix} \right\} q^b = q^r. \quad (3.21)$$

Therefore, each graph H gets a weight $q^{k(H)} v^{|E(H)|}$, and the sum over graphs gives $Z_n(q, v)$. This proves (3.9).

The more general formulae (3.7)/(3.8) are derived by a variant of this proof in which sums over integers $n \in \mathbb{N} \setminus \{0\}$ are replaced by sums over multi-indices $\mathbf{n} \in \mathbb{N}^V \setminus \{\mathbf{0}\}$. In the one-dimensional case, we were considering partitions of $[n]$, a collection of n distinct objects of a single type. We are now working with multi-indices in \mathbb{N}^V , and so we will be partitioning collections of objects of $|V|$ different types. We begin with some notation. Given $\mathbf{n} = (n_i)_{i \in V} \in \mathbb{N}^V$, we write $[\mathbf{n}]$ for the set of ordered pairs $\{(i, \alpha) : i \in V \text{ and } \alpha \in [n_i]\}$. Thus $|[\mathbf{n}]| = \sum_{i \in V} n_i$. We shall think of the subset $[\mathbf{n}]_i := \{(i, \alpha) : \alpha \in [n_i]\}$ as a copy of $[n_i]$ with type i ; thus, $[\mathbf{n}]$ is a collection of $|\mathbf{n}|$ objects, with n_i objects of type i for each $i \in V$. Note that $G'[\mathbf{n}]$ is simply the complete graph on the vertex set $[\mathbf{n}]$. Now, given a partition $\boldsymbol{\pi} = (\boldsymbol{\pi}_1, \dots, \boldsymbol{\pi}_k)$ of $[\mathbf{n}]$ (with all $\boldsymbol{\pi}_j$ nonempty), we write $|\boldsymbol{\pi}| = k$. The *block sizes* $B(\boldsymbol{\pi}_j)$ are elements of $\mathbb{N}^V \setminus \{\mathbf{0}\}$ defined by

$$B(\boldsymbol{\pi}_j) = (\boldsymbol{\pi}_j \cap [\mathbf{n}]_i)_{i \in V}. \quad (3.22)$$

Thus $B(\boldsymbol{\pi}_j)$ lists the number of objects of each type contained in the j th block of $\boldsymbol{\pi}$.

Now let us define the weight

$$W(\mathbf{n}) := \left(\prod_{ii' \in E} (1 + v_{ii'})^{n_i n_{i'}} \right) \left(\prod_{i \in V} (1 + w_i)^{n_i(n_i-1)/2} \right). \quad (3.23)$$

The right-hand side of (3.8) is then

$$\left(1 + \sum_{\mathbf{n} \neq \mathbf{0}} W(\mathbf{n}) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \right)^q = 1 + \sum_{b \geq 1} \frac{q^b}{b!} \left(\sum_{\mathbf{n} \neq \mathbf{0}} W(\mathbf{n}) \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \right)^b \quad (3.24a)$$

$$= 1 + \sum_{b \geq 1} \frac{q^b}{b!} \sum_{\mathbf{n}_1, \dots, \mathbf{n}_b \neq \mathbf{0}} \frac{\mathbf{x}^{\sum \mathbf{n}_j}}{\prod \mathbf{n}_j!} \prod_{j=1}^b W(\mathbf{n}_j). \quad (3.24b)$$

Now let $\mathbf{n} = \sum_{j=1}^b \mathbf{n}_j$, and consider $\mathbf{n}_1, \dots, \mathbf{n}_b$ as the block sizes in an ordered partition $\boldsymbol{\pi}$ of $[\mathbf{n}]$ into b nonempty blocks. The number of ordered partitions with those block sizes is the (multidimensional) multinomial coefficient $\mathbf{n}! / \prod_{j=1}^b \mathbf{n}_j!$, so (3.24) is equal to

$$1 + \sum_{\mathbf{n} \neq \mathbf{0}} \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \sum_{\boldsymbol{\pi}} \frac{q^{|\boldsymbol{\pi}|}}{|\boldsymbol{\pi}|!} \prod_{j=1}^{|\boldsymbol{\pi}|} W(B(\boldsymbol{\pi}_j)), \quad (3.25)$$

where the second sum is over ordered partitions of $[\mathbf{n}]$. Passing to *unordered* partitions, this becomes

$$1 + \sum_{\mathbf{n} \neq \mathbf{0}} \frac{\mathbf{x}^{\mathbf{n}}}{\mathbf{n}!} \sum_{\boldsymbol{\pi}} q^{|\boldsymbol{\pi}|} \prod_{j=1}^{|\boldsymbol{\pi}|} W(B(\boldsymbol{\pi}_j)). \quad (3.26)$$

The factor $W(B(\boldsymbol{\pi}_j))$ corresponds to a sum over graphs with vertex set $\boldsymbol{\pi}_j$, such that edges within $[\mathbf{n}]_i$ have weight w_i and edges between $[\mathbf{n}]_i$ and $[\mathbf{n}]_{i'}$ have weight $v_{ii'}$. As before, we can reinterpret (3.26) as a sum over graphs H with vertex set $[\mathbf{n}]$, with these same edge weights. The final step of the argument — summing over partitions $\boldsymbol{\pi}$ that are compatible with H — is identical to that in the one-dimensional case: it makes no difference whether the vertex set is $[n]$ or $[\mathbf{n}]$. We therefore see that each graph H gets weight $q^{k(H)}$ times the product of edge weights, and we are done. \square

3.3. Consequences for $Z_G(\mathbf{q}, \mathbf{v})$. In view of Theorem 3.3, we can apply all the results of Section 2 with the identifications

$$a_{\mathbf{n}}(q) = Z_{G'[\mathbf{n}]}(q, \mathbf{v}, \mathbf{w}) \quad (3.27a)$$

$$\widehat{a}_{\mathbf{n}}(q) = \widehat{Z}_{G'[\mathbf{n}]}(q, \mathbf{v}, \mathbf{w}) \quad (3.27b)$$

$$c_{\mathbf{n}} = C_{G'[\mathbf{n}]}(\mathbf{v}, \mathbf{w}) \quad (3.27c)$$

$$a_{\mathbf{n}}(1) = \left(\prod_{ij \in E} (1 + v_{ij})^{n_i n_j} \right) \left(\prod_{i \in V} (1 + w_i)^{n_i(n_i-1)/2} \right) \quad (3.27d)$$

We refrain from writing out all the formulae, which are obtained by substituting (3.27a–d) into (2.3), (2.4)/(2.5), (2.7)–(2.13), (2.14), (2.15), (2.17), (2.18), (2.19) and (2.20). Let us simply show a few important results that arise from the partition formula (2.7)/(2.12) and from the convolution formula (2.14)–(2.15).

3.3.1. Partition formulae. By specializing (2.7)/(2.12) to $\mathbf{n} = \mathbf{1}$, we can obtain a partition formula that expresses the multivariate Tutte polynomial of G at $q = q_2$ in terms of the multivariate Tutte polynomials of induced subgraphs of G at $q = q_1$, for arbitrary choices of q_1 and q_2 :

Proposition 3.4. *Let $G = (V, E)$ be a finite loopless graph. Then*

$$Z_G(q_2, \mathbf{v}) = \sum_{\pi \in \Pi(V)} (q_2/q_1)^{|\pi|} \prod_{B \in \pi} Z_{G[B]}(q_1, \mathbf{v}), \quad (3.28)$$

where the sum runs over (unordered) partitions π of the vertex set V , the product runs over blocks B of π , and $G[B]$ denotes the induced subgraph of G on the vertex set B .¹⁵ Equivalently we can write

$$Z_G(q_2, \mathbf{v}) = \sum_{\pi \in \Pi(V)} \left(\prod_{j=0}^{|\pi|-1} (q_2 - j q_1) \right) \prod_{B \in \pi} \widehat{Z}_{G[B]}(q_1, \mathbf{v}) \quad (3.29)$$

or

$$\widehat{Z}_G(q_2, \mathbf{v}) = \sum_{\pi \in \Pi(V)} \left(\prod_{j=1}^{|\pi|-1} (q_2 - j q_1) \right) \prod_{B \in \pi} \widehat{Z}_{G[B]}(q_1, \mathbf{v}). \quad (3.30)$$

Proof. Specializing (2.7) [or equivalently (2.12)] to $\mathbf{n} = \mathbf{1}$, we obtain a sum over ordered partitions $\pi = (\pi_1, \dots, \pi_\ell)$ of V into ℓ nonempty blocks. Passing from ordered to unordered partitions, we get a factor $\ell!$ that cancels the $1/\ell!$ in (2.7)/(2.12). \square

Proposition 3.4 is very powerful because we are free to choose q_1 as we please; then $Z_G(q_2, \mathbf{v})$ can be written in terms of $Z_{G[B]}(q_1, \mathbf{v})$. The two most important special cases are $q_1 = 0$ and $q_1 = 1$.¹⁶

¹⁵Readers who dislike dividing by indeterminates can reinterpret this formula by writing $r = q_2/q_1$ and hence $q_2 = r q_1$, as in (2.7).

¹⁶The case $q_1 = -1$ is also of interest, as it is related to acyclic orientations. See Lass [35] for some interesting material that may be relevant in the present context.

Case $q_1 = 0$. Taking the limit $q_1 \rightarrow 0$ in (3.28), or equivalently just setting $q_1 = 0$ in (3.29), we obtain

$$Z_G(q, \mathbf{v}) = \sum_{\pi \in \Pi(V)} q^{|\pi|} \prod_{B \in \pi} C_{G[B]}(\mathbf{v}). \quad (3.31)$$

[This formula can alternatively be derived from (2.3) by the same reasoning as used in proving Proposition 3.4.] Of course, (3.31) also has an obvious direct combinatorial proof, based on the definition (3.1): first we classify subsets $A \subseteq E$ according to the partition π of the vertex set V that is induced by the connected components of the subgraph (V, A) ; then we sum over ways of connecting up each component.

Case $q_1 = 1$. This special case is slightly less obvious, and it is worth stating it explicitly:

Corollary 3.5. *Let $G = (V, E)$ be a finite loopless graph. Then*

$$Z_G(q, \mathbf{v}) = \sum_{\pi \in \Pi(V)} q^{|\pi|} \prod_{B \in \pi} \prod_{e \in E(G[B])} (1 + v_e) \quad (3.32)$$

where the sum runs over (unordered) partitions π of the vertex set V , the outermost product runs over blocks B of π , and the innermost product runs over edges in the induced subgraph $G[B]$. In particular,

$$C_G(\mathbf{v}) = \sum_{\pi \in \Pi(V)} (-1)^{|\pi|-1} (|\pi| - 1)! \prod_{B \in \pi} \prod_{e \in E(G[B])} (1 + v_e) \quad (3.33)$$

$$Z_G(-1, \mathbf{v}) = \sum_{\pi \in \Pi(V)} (-1)^{|\pi|} |\pi|! \prod_{B \in \pi} \prod_{e \in E(G[B])} (1 + v_e) \quad (3.34)$$

where $|\pi|$ denotes the number of blocks in π .

Specializing (3.32)–(3.34) to $v_e = -1$ for all edges e , we have:

Corollary 3.6. *Let $G = (V, E)$ be a finite loopless graph with $|V| = n$. Then:*

$$P_G(q) = \sum_{k=1}^n q^k Q_G(k) \quad (3.35)$$

$$P'_G(0) = C_G(-1) = \sum_{k=1}^n (-1)^{k-1} (k-1)! Q_G(k) \quad (3.36)$$

$$P_G(-1) = \sum_{k=1}^n (-1)^k k! Q_G(k) \quad (3.37)$$

where $Q_G(k)$ denotes the number of partitions of V into k nonempty independent subsets.

Of course, (3.35) is well known and has a trivial direct proof by counting proper q -colorings; and (3.36)/(3.37) are immediate corollaries of (3.35).

Let us conclude by giving two direct combinatorial proofs of the fundamental Proposition 3.4: one using the subgraph representation (3.1), and the other using the coloring representation (3.5).

Second Proof of Proposition 3.4. Use the partition formula (3.31) [which itself had a simple direct combinatorial proof] to expand $Z_G(q_2, \mathbf{v})$ on the left-hand side of (3.28) and to expand each $Z_{G[B]}(q_1, \mathbf{v})$ on the right-hand side. We can then interpret the double sum on the right-hand side as a sum over partitions π of V , with weight q_1 for each block, together with a sum over ways of grouping these blocks into nonempty groups, with a weight $(q_2/q_1)^k$ if we have k groups. Using the identity

$$\sum_{k=1}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} (q_2/q_1)^k = (q_2/q_1)^n \quad (3.38)$$

where $n = |\pi|$, we get exactly the expression on the left-hand side. \square

It will not escape the reader's notice that the argument using (3.38) is identical to the one given earlier using (3.21).

Third Proof of Proposition 3.4. Since both sides of (3.28) are polynomials in q_1 and q_2 , it suffices to prove the identity for all pairs of positive integers q_1, q_2 such that $r = q_2/q_1$ is also an integer. We shall do this using Theorem 3.1. Consider a (not-necessarily-proper) coloring $\sigma: V \rightarrow [q_2]$. Fix a bijection $[q_2] \simeq [q_1] \times [r]$ and write $\sigma = (\tau, \psi)$ where $\tau: V \rightarrow [q_1]$ and $\psi: V \rightarrow [r]$. We then trivially have

$$Z_G(q_2, \mathbf{v}) = \sum_{\sigma: V \rightarrow [q_2]} \prod_{e=ij \in E} \left[1 + v_e \delta(\sigma(i), \sigma(j)) \right] \quad (3.39a)$$

$$= \sum_{\psi: V \rightarrow [r]} \sum_{\tau: V \rightarrow [q_1]} \prod_{e=ij \in E} \left[1 + v_e \delta(\tau(i), \tau(j)) \delta(\psi(i), \psi(j)) \right]. \quad (3.39b)$$

The map ψ induces a partition π_ψ of V into its nonempty color classes, and the remaining sum over τ gives (by Theorem 3.1 again) $\prod_{B \in \pi_\psi} Z_{G[B]}(q_1, \mathbf{v})$. On the other hand, each partition π of V arises in this way from $r^{|\pi|}$ different colorings ψ , since we have r choices to color the lexicographically first block of π , $r - 1$ choices for the lexicographically second block, and so forth. This proves (3.28). \square

Important note: The formulae in this subsection — notably (3.33) together with the $q = 1$ case of (3.31) — are strongly reminiscent of Möbius inversion on the lattice of partitions $\Pi(V)$. This is not an accident. In fact, the formula (3.30) is a special case of a “ q_1 - q_2 generalization” of Möbius inversion that we present in Appendix A. We are grateful to an anonymous referee for drawing our attention to the connection of these formulae with Möbius inversion and for challenging us to find the underlying general principle.

3.3.2. *Convolution formulae.* In a similar way we can obtain convolution formulae for the multivariate Tutte polynomial by specializing (2.14)–(2.15):

Proposition 3.7. *Let $G = (V, E)$ be a finite loopless graph. Then*

$$Z_G(q_1 + q_2, \mathbf{v}) = \sum_{W \subseteq V} Z_{G[W]}(q_1, \mathbf{v}) Z_{G[V \setminus W]}(q_2, \mathbf{v}). \quad (3.40)$$

Moreover, for each $i \in V$ we have

$$\widehat{Z}_G(q_1 + q_2, \mathbf{v}) = \sum_{\substack{W \subseteq V \\ W \ni i}} \widehat{Z}_{G[W]}(q_1, \mathbf{v}) Z_{G[V \setminus W]}(q_2, \mathbf{v}) \quad (3.41)$$

and in particular

$$Z_G(q, \mathbf{v}) = \sum_{\substack{W \subseteq V \\ W \ni i}} q C_{G[W]}(\mathbf{v}) Z_{G[V \setminus W]}(q, \mathbf{v}) . \quad (3.42)$$

We also have

$$|V| Z_G(q_2, \mathbf{v}) = \sum_{\emptyset \neq W \subseteq V} \left[(q_1 + q_2)|W| - q_1|V| \right] \widehat{Z}_{G[W]}(q_1, \mathbf{v}) Z_{G[V \setminus W]}(q_2, \mathbf{v}) . \quad (3.43)$$

Proof. (3.40) and (3.41) are just (2.14) and (2.15), respectively, specialized to $\mathbf{n} = \mathbf{1}$. Further specializing the latter to $q_1 = 0$, we obtain (3.42). Finally, (3.43) is obtained by subtracting $(q_1 + q_2) \times (3.41)$ from (3.40) and summing over $i \in V$. \square

For completeness, let us give a simple direct proof of (3.41). [(3.40) is similar but easier, and is left as an exercise for the reader.]

Alternate Proof of Proposition 3.7. From the definition (3.2) we have

$$\widehat{Z}_G(q_1 + q_2, \mathbf{v}) = \sum_{A \subseteq E} (q_1 + q_2)^{k(A)-1} \prod_{e \in A} v_e . \quad (3.44)$$

The factor $(q_1 + q_2)^{k(A)-1}$ can be handled by choosing, for each connected component of (V, A) other than the component containing the distinguished vertex i , to color it either “1” or “2”, with a corresponding factor q_1 or q_2 , and summing over all such choices. Now define $W \subseteq V$ to be the vertex set corresponding to the union of the components colored “1” together with the component containing i . By construction, W is compatible with A in the sense that A has no edges connecting W to $V \setminus W$. Moreover, every $W \ni i$ compatible with A is obtained exactly once by some choice of colors “1” and “2”. On the other hand, the right-hand side of (3.41) is given precisely by such a sum over compatible pairs (W, A) [the $\widehat{}$ on $\widehat{Z}_{G[W]}(q_1, \mathbf{v})$ accounts for the fact that the component containing i carries no color and hence no factor q_1]. \square

We do not know whether the Abel-type extensions (2.19)/(2.20), when applied to the multivariate Tutte polynomial, are of any interest.

3.4. Specialization to the complete graphs. Let us now take G to be the complete graph K_n , with all the edge weights v_e set equal to the same value v . We write $Z_n(q, v)$, $\widehat{Z}_n(q, v)$ and $C_n(v)$ for the corresponding polynomials.

3.4.1. *Partition formulae.* Specializing (3.28), we obtain the general identity

$$Z_n(q_2, v) = \sum_{\pi \in \Pi_n} (q_2/q_1)^{|\pi|} \prod_{i=1}^{|\pi|} Z_{|\pi_i|}(q_1, v) , \quad (3.45)$$

where the sum runs over (unordered) partitions π of $[n]$, say $\pi = \{\pi_1, \dots, \pi_\ell\}$, and $|\pi|$ denotes the number of blocks in π (i.e., $|\pi| = \ell$). Special cases of (3.45) arise at particular values of q_1 :

Case $q_1 = 0$. Specializing (3.31), we obtain the trivial identity

$$Z_n(q, v) = \sum_{\pi \in \Pi_n} q^{|\pi|} \prod_{i=1}^{|\pi|} C_{|\pi_i|}(v). \quad (3.46)$$

Case $q_1 = 1$. Specializing (3.32)–(3.34), we obtain the less trivial relations

$$Z_n(q, v) = \sum_{\pi \in \Pi_n} q^{|\pi|} (1+v)^{n(n-1)/2 - \|\pi\|} \quad (3.47)$$

$$C_n(v) = \sum_{\pi \in \Pi_n} (-1)^{|\pi|-1} (|\pi|-1)! (1+v)^{n(n-1)/2 - \|\pi\|} \quad (3.48)$$

$$Z_n(-1, v) = \sum_{\pi \in \Pi_n} (-1)^{|\pi|} |\pi|! (1+v)^{n(n-1)/2 - \|\pi\|} \quad (3.49)$$

where

$$\|\pi\| = \sum_{1 \leq i < j \leq \ell} |\pi_i| |\pi_j| = \frac{1}{2} \left(n^2 - \sum_{i=1}^{\ell} |\pi_i|^2 \right) \quad (3.50)$$

denotes the number of “cross-edges” in π , i.e., the number of edges in the complete ℓ -partite graph with vertex classes π_1, \dots, π_ℓ . Equivalently,

$$\frac{n(n-1)}{2} - \|\pi\| = \sum_{i=1}^{\ell} \frac{|\pi_i|(|\pi_i|-1)}{2} \quad (3.51)$$

is the number of “internal edges” in π . The formulae (3.47)–(3.49) will play an important role in our study of the large- n asymptotics of $C_n(v)$ and $Z_n(q, v)$ at complex v [55].

3.4.2. Convolution formulae. Specializing (3.40)–(3.43) and exploiting the symmetries of the complete graph, we obtain

$$Z_n(q_1 + q_2, v) = \sum_{k=0}^n \binom{n}{k} Z_k(q_1, v) Z_{n-k}(q_2, v) \quad (3.52)$$

$$\widehat{Z}_n(q_1 + q_2, v) = \sum_{k=1}^n \binom{n-1}{k-1} \widehat{Z}_k(q_1, v) Z_{n-k}(q_2, v) \quad \text{for } n \geq 1 \quad (3.53)$$

$$Z_n(q, v) = \sum_{k=1}^n \binom{n-1}{k-1} q C_k(v) Z_{n-k}(q, v) \quad \text{for } n \geq 1 \quad (3.54)$$

$$Z_n(q_2, v) = \sum_{k=1}^n \left[\binom{n-1}{k-1} (q_1 + q_2) - \binom{n}{k} q_1 \right] \widehat{Z}_k(q_1, v) Z_{n-k}(q_2, v) \quad \text{for } n \geq 1 \quad (3.55)$$

On the other hand, these are just the one-dimensional identities (1.5), (1.6), (1.9) and (1.10) applied to the family $a_n(q) = Z_n(q, v)$, which is indeed of the required form (1.2) by virtue of the generating-function formula (3.9).

The formula (3.54) can be used to compute the $Z_n(q, v)$ inductively given the $C_k(v)$. On the other hand, if we specialize (3.54) to $q = 1$ and use (3.3), we obtain the identity

$$(1 + v)^{n(n-1)/2} = \sum_{k=1}^n \binom{n-1}{k-1} C_k(v) (1 + v)^{(n-k)(n-k-1)/2} \quad \text{for } n \geq 1 \quad (3.56)$$

or equivalently

$$C_n(v) = (1 + v)^{n(n-1)/2} - \sum_{k=1}^{n-1} \binom{n-1}{k-1} C_k(v) (1 + v)^{(n-k)(n-k-1)/2} \quad \text{for } n \geq 1, \quad (3.57)$$

which can be used to compute the $C_n(v)$ inductively *ab initio*. We call (3.56)/(3.57) the “linear” identity for $C_n(v)$. It goes back at least to Leroux [37, eq. (3.3)] and is probably much older; it can be proven by an easy direct argument based on considering the size k of the connected component containing a fixed vertex. [The same argument proves (3.54).]

Alternatively, if we specialize (3.55) to $q_1 = 1$ and use (3.3), we obtain

$$Z_n(q, v) = \sum_{k=1}^n \left[\binom{n-1}{k-1} (1 + q) - \binom{n}{k} \right] (1 + v)^{k(k-1)/2} Z_{n-k}(q, v) \quad \text{for } n \geq 1, \quad (3.58)$$

which allows us to go directly from $Z_n(1, v) = (1 + v)^{n(n-1)/2}$ to $Z_n(q, v)$ without passing through $C_n(v)$. We can also rewrite (3.58) as

$$\begin{aligned} \widehat{Z}_n(q, v) &= (1 + v)^{n(n-1)/2} \\ &\quad + \sum_{k=1}^{n-1} \left[\binom{n-1}{k} q - \binom{n-1}{k-1} \right] (1 + v)^{(n-k)(n-k-1)/2} \widehat{Z}_k(q, v) \end{aligned} \quad \text{for } n \geq 1, \quad (3.59)$$

which manifestly generalizes (3.57) and reduces to it when $q = 0$.

3.5. A generalization. Let $\mathbf{a} = \{a_n\}_{n=0}^\infty$ be an arbitrary sequence of coefficients belonging to the ring R , satisfying $a_0 = 1$, and define the family of polynomials

$$Z_n(q; \mathbf{a}) = \sum_{\pi \in \Pi_n} q^{|\pi|} \prod_{i=1}^{|\pi|} a_{|\pi_i|} \quad (3.60)$$

with the convention $Z_0 = 1$. Note that $Z_n(1; \mathbf{a}) = a_n$. This definition generalizes (3.47)/(3.51), and if we specialize to $a_n = (1 + v)^{n(n-1)/2}$ we obtain $Z_n(q, v)$.

A simple counting argument shows that

$$Z_n(q; \mathbf{a}) = \sum_{b=0}^n \frac{q^b}{b!} \sum_{\substack{n_1, \dots, n_b \geq 1 \\ \sum n_i = n}} \binom{n}{n_1, \dots, n_b} \prod_{i=1}^b a_{n_i}. \quad (3.61)$$

Using this to compute the exponential generating function of the $\{Z_n(q; \mathbf{a})\}$, we find, after a short calculation using the binomial series, that

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} Z_n(q; \mathbf{a}) = \left(\sum_{n=0}^{\infty} \frac{x^n}{n!} a_n \right)^q. \quad (3.62)$$

In other words, the family $\{Z_n(q; \mathbf{a})\}$ is of the form $\{a_n(q)\}$ defined in (1.2), with $a_n(1) = a_n$. But this should hardly be surprising, as (3.61) is simply the one-dimensional case of (2.12) with $q_2 = q$ and $q_1 = 1$. We have thus come full circle.

Likewise, the recursion

$$Z_n(q; \mathbf{a}) = q \sum_{k=1}^n \binom{n-1}{k-1} a_k Z_{n-k}(q-1; \mathbf{a}) \quad (3.63)$$

can be proven from (3.60) by an easy direct argument based on considering the size k of the block of π containing some fixed element of $[n]$. But this identity is nothing other than (1.11) specialized to $q_1 = 1$ and $q_2 = q - 1$.

3.6. A related problem. Suppose we try a definition analogous to (3.60), but with ordinary powers $q^{|\pi|}$ in place of falling factorials $q^{\underline{|\pi|}}$. That is, let $\mathbf{c} = \{c_n\}_{n=1}^{\infty}$ be an arbitrary sequence of coefficients belonging to the ring R , and define the family of polynomials

$$Y_n(q; \mathbf{c}) = \sum_{\pi \in \Pi_n} q^{|\pi|} \prod_{i=1}^{|\pi|} c_{|\pi_i|} \quad (3.64)$$

with the convention $Y_0 = 1$. Of course, the factor $q^{|\pi|}$ is superfluous, because we can simply multiply each c_n by q , but it is convenient to keep it explicit. This definition generalizes (3.46), and if we specialize to $c_n = C_n(v)$ we obtain $Z_n(q, v)$.

A simple counting argument shows that

$$Y_n(q; \mathbf{c}) = \sum_{b=0}^n \frac{q^b}{b!} \sum_{\substack{n_1, \dots, n_b \geq 1 \\ \sum n_i = n}} \binom{n}{n_1, \dots, n_b} \prod_{i=1}^b c_{n_i}. \quad (3.65)$$

Using this to compute the exponential generating function of the $\{Y_n(q; \mathbf{c})\}$, we find, after a short calculation using the exponential series, that

$$\sum_{n=0}^{\infty} \frac{x^n}{n!} Y_n(q; \mathbf{c}) = \exp \left(q \sum_{n=1}^{\infty} \frac{x^n}{n!} c_n \right). \quad (3.66)$$

In other words, the family $\{Y_n(q; \mathbf{c})\}$ is again of the form $\{a_n(q)\}$ defined in (1.2). But this should hardly be surprising, as (3.65) is simply the one-dimensional case of (2.3) [i.e., (1.3).] We have again come full circle.

Likewise, the recursion

$$Y_n(q; \mathbf{c}) = q \sum_{k=1}^n \binom{n-1}{k-1} c_k Y_{n-k}(q; \mathbf{c}), \quad (3.67)$$

which can be proven from (3.64) by considering the size k of the block of π containing some fixed element of $[n]$, is nothing other than (1.9).

4. NONLINEAR IDENTITY FOR THE MULTIVARIATE TUTTE POLYNOMIAL

4.1. General identity. In the preceding section we proved the identities (3.41) and (3.42) for the multivariate Tutte polynomial $Z_G(q, \mathbf{v})$, which are based on choosing a single distinguished vertex i . We shall now prove a different identity that is based on choosing a *pair* of distinguished vertices i, j . Unlike the identities discussed in the preceding section, this one relies on the graphical structure of $Z_G(q, \mathbf{v})$ and does *not* appear to generalize to arbitrary families $\{a_n(q)\}$ of the type (1.2).

So let $G = (V, E)$ be a finite undirected graph with $|V| \geq 2$, and let $i, j \in V$ with $i \neq j$. We then have the following identity:

Theorem 4.1. *Let $G = (V, E)$ be a finite graph, and let $i, j \in V$ with $i \neq j$. Then*

$$Z_G(q, \mathbf{v}) = \sum_{\substack{W \subseteq V \\ W \ni i, W \not\ni j}} \left[q - 1 + \prod_{e \in E(W, j)} (1 + v_e) \right] C_{G[W]}(\mathbf{v}) Z_{G[V \setminus W]}(q, \mathbf{v}) \quad (4.1)$$

where $E(W, j)$ denotes the set of all edges with one endpoint in W and the other endpoint at j . In particular,

$$C_G(\mathbf{v}) = \sum_{\substack{W \subseteq V \\ W \ni i, W \not\ni j}} \left[\prod_{e \in E(W, j)} (1 + v_e) - 1 \right] C_{G[W]}(\mathbf{v}) C_{G[V \setminus W]}(\mathbf{v}). \quad (4.2)$$

We do not know whether there exists a generalization of (4.1) involving q_1 and q_2 (rather than just $q_1 = 0$).

The proof of Theorem 4.1 is based on looking at the connected component of i in the induced subgraph where j is deleted:

Proof. Start from the definition $Z_G(q, \mathbf{v}) = \sum_{A \subseteq E} q^{k(A)} \prod_{e \in A} v_e$. Let $G' = (V, A)$ and $G'' = G' \setminus j$, and let W be the vertex set of the connected component of G'' containing i . Let us now sum over all $A \subseteq E$ that give rise in this way to a specified set $W \subseteq V$, and let us split this sum into two parts according as A does or does not contain at least one edge from j to W . If A does not contain such an edge, then W is a connected component of G' , giving rise to a factor q , and we get $q C_{G[W]}(\mathbf{v}) Z_{G[V \setminus W]}(q, \mathbf{v})$. On the other hand, if A contains at least one such edge, then W forms part of the connected component of j ($\in V \setminus W$) in G' ; hence there is no factor q , but there is a factor v_e for each edge e in the (nonempty) subset $A \cap E(W, j)$. Summing over all such nonempty subsets of $E(W, j)$, we get

$$\left[\prod_{e \in E(W, j)} (1 + v_e) - 1 \right] C_{G[W]}(\mathbf{v}) Z_{G[V \setminus W]}(q, \mathbf{v}). \quad (4.3)$$

Putting everything together gives (4.1). Specialization to $q = 0$ yields (4.2). \square

There is also a variant of (4.1) in which the right-hand side is completely expanded out. For notational simplicity we shall assume that G has no loop at the distinguished vertex j . (In the general case one must multiply by a factor $1 + v_e$ for each loop e at j .)

Proposition 4.2. *Let $G = (V, E)$ be a finite graph, let $j \in V$, and suppose that G has no loop at j . Then*

$$\widehat{Z}_G(q, \mathbf{v}) = \sum_{\pi \in \Pi(V \setminus \{j\})} \prod_{B \in \pi} \left[q - 1 + \prod_{e \in E(B, j)} (1 + v_e) \right] C_{G[B]}(\mathbf{v}) \quad (4.4)$$

where the sum runs over (unordered) partitions π of $V \setminus \{j\}$, and the product runs over blocks B of π . In particular,

$$C_G(\mathbf{v}) = \sum_{\pi \in \Pi(V \setminus \{j\})} \prod_{B \in \pi} \left[\prod_{e \in E(B, j)} (1 + v_e) - 1 \right] C_{G[B]}(\mathbf{v}) \quad (4.5)$$

Proof. In (3.2), consider a term $A \subseteq E$, and let $G' = (V, A)$ and $G'' = G' \setminus j$. Let π be the partition of $V \setminus \{j\}$ into vertex sets of connected components of G'' . We can recover G' from G'' by adjoining, for each block B of π , zero or more edges from the set $E(B, j)$. If we adjoin zero edges, we get an extra factor q because B becomes the vertex set of a connected component of G' that is distinct from the component containing j ; if we adjoin one or more edges, we get no such factor. This proves (4.4). Specialization to $q = 0$ yields (4.5). \square

When $v_e = v$ for all edges e , the formulae (4.4) and (4.5) can be found in Gessel [17, Theorems 13 and 10]. The special case of (4.5) in which $v_e = -1$ for all e was also found by Borgs [8] and used recently by him [9, Lemma 3.2] to bound the complex zeros of chromatic polynomials (a variant of the proof in [56]). Indeed, we were led to formulate (4.4)/(4.5) by meditating on Borgs' special case, oblivious to the fact that they had already been essentially found by Gessel!

Finally, let us use (4.2) to prove an interesting inequality concerning the polynomials $C_G(\mathbf{v})$. Let

$$c(A) = |A| - |V| + k(A) \quad (4.6)$$

be the cyclomatic number of the subgraph (V, A) , and let us define the generalized connected sum

$$C_G(\mathbf{v}, \lambda) = \sum_{\substack{A \subseteq E \\ k(A) = 1}} \lambda^{c(A)} \prod_{e \in A} v_e \quad (4.7a)$$

$$= \lambda^{-(|V|-1)} C_G(\lambda \mathbf{v}). \quad (4.7b)$$

Of course, (4.7b) shows that $C_G(\mathbf{v}, \lambda)$ contains no more information than $C_G(\mathbf{v})$; it is simply a convenient way of scaling all the variables v_e simultaneously while removing a factor $\lambda^{|V|-1}$. In particular, $C_G(\mathbf{v}, \lambda)$ interpolates between the spanning-tree sum ($\lambda = 0$) and the connected-spanning-subgraph sum ($\lambda = 1$). We then have the following result [56, Remark 2 in Section 4.1] [53, Proposition 2.5]:

Proposition 4.3. *Let $G = (V, E)$ be a finite graph ($V \neq \emptyset$) equipped with real edge weights $\mathbf{v} = (v_e)_{e \in E}$ satisfying $-1 \leq v_e \leq 0$ for all $e \in E$. Then*

$$(-1)^{\ell+|V|-1} \frac{\partial^\ell}{\partial \lambda^\ell} C_G(\mathbf{v}, \lambda) \geq 0 \quad (4.8)$$

on $0 \leq \lambda \leq 1$, for all integers $\ell \geq 0$.

This inequality was proven in [53, 56] using a “partitionability” method going back to Penrose [41].¹⁷ Here we prove it using the recursion (4.2):

Proof of Proposition 4.3. By induction on $|V|$. When $|V| = 1$, G necessarily consists of zero or more loops attached to the sole vertex, so

$$C_G(\mathbf{v}, \lambda) = \prod_{e \in E} (1 + \lambda v_e). \quad (4.9)$$

Using $v_e \leq 0$ and $1 + \lambda v_e \geq 0$, it is easy to see that (4.8) holds.

Now assume that $|V| \geq 2$. Replacing \mathbf{v} by $\lambda \mathbf{v}$ in (4.2) and using (4.7b), we obtain

$$C_G(\mathbf{v}, \lambda) = \sum_{\substack{W \subseteq V \\ W \ni i, W \not\ni j}} \lambda^{-1} \left[\prod_{e \in E(W, j)} (1 + v_e) - 1 \right] C_{G[W]}(\mathbf{v}, \lambda) C_{G[V \setminus W]}(\mathbf{v}, \lambda) \quad (4.10a)$$

$$= \sum_{\substack{W \subseteq V \\ W \ni i, W \not\ni j}} \left(\sum_{\emptyset \neq B \subseteq E(W, j)} \lambda^{|B|-1} \prod_{e \in B} v_e \right) C_{G[W]}(\mathbf{v}, \lambda) C_{G[V \setminus W]}(\mathbf{v}, \lambda). \quad (4.10b)$$

Now apply $\partial^\ell / \partial \lambda^\ell$ to both sides: on the right-hand side we will have terms in which ℓ_1 derivatives act on $\sum \lambda^{|B|-1} \prod v_e$, ℓ_2 act on $C_{G[W]}(\mathbf{v}, \lambda)$, and ℓ_3 act on $C_{G[V \setminus W]}(\mathbf{v}, \lambda)$, where $\ell_1 + \ell_2 + \ell_3 = \ell$. The induction hypothesis is applicable because $1 \leq |W| \leq |V| - 1$. So it suffices to check that

$$(-1)^{\ell_1-1} \frac{\partial^{\ell_1}}{\partial \lambda^{\ell_1}} \left(\sum_{\emptyset \neq B \subseteq E(W, j)} \lambda^{|B|-1} \prod_{e \in B} v_e \right) \geq 0. \quad (4.11)$$

To prove this, let us order the elements of $E(W, j)$, i.e., consider $E(W, j) \simeq [m]$ where $m = |E(W, j)|$. Then, by singling out the smallest element in each set B , we can write

$$\sum_{\emptyset \neq B \subseteq [m]} \lambda^{|B|-1} \prod_{e \in B} v_e = \sum_{i=1}^m v_i \prod_{j=i+1}^m (1 + \lambda v_j). \quad (4.12)$$

We have $v_i \leq 0$; each $\partial / \partial \lambda$ brings down some factor $v_j \leq 0$; and the undifferentiated factors satisfy $1 + \lambda v_j \geq 0$. So (4.11) holds. \square

It is worth remarking that (4.11) is nothing other than Proposition 4.3 specialized to the graph $K_2^{(m)}$ consisting of two vertices connected by m parallel edges. Moreover, the identity (4.12) is nothing other than (one version of) the partitionability identity for $K_2^{(m)}$.

¹⁷For the partitionability method (which applies to matroids as well as graphs), see also [7, 19] and the other references mentioned in [53, Section 2.2].

4.2. Specialization to the complete graphs. Specializing (4.1) to the complete graph K_n with equal weights v , we obtain

$$Z_n(q, v) = \sum_{k=1}^{n-1} \binom{n-2}{k-1} [q + (1+v)^k - 1] C_k(v) Z_{n-k}(q, v) \quad \text{for } n \geq 2 \quad (4.13)$$

(see [17, eq. (5)] for an equivalent formula). Specializing this to $q = 0$, we obtain

$$C_n(v) = \sum_{k=1}^{n-1} \binom{n-2}{k-1} [(1+v)^k - 1] C_k(v) C_{n-k}(v) \quad \text{for } n \geq 2, \quad (4.14)$$

a result apparently first proven by Leroux [37, eq. (3.5)] (see also [17, eq. (2)]).¹⁸ And specializing this latter formula to $v = 1$, we obtain the identity of Riordan, Nijenhuis, Wilf and Kreweras [31, 40, 47] for counting connected graphs. We call (4.14) the “nonlinear” identity for $C_n(v)$.

On the other hand, specializing (4.13) to $q = 1$, dividing both sides by $(1+v)^{n-1}$ and relabelling $n \rightarrow n+1$, we recover the “linear” identity (3.56).

The formula (4.14) has also been derived in a different context. Let T be a tree with vertex set $[n]$, rooted at the vertex 1. An *inversion* of T is an ordered pair (j, k) of vertices such that $j > k > 1$ and k is a descendant of j (i.e., the path from 1 to k passes through j). We define the *inversion enumerator for trees* [20, 31, 38] (see also [10, 17, 19]) to be the polynomial

$$I_n(y) = \sum_{\text{trees } T \text{ on } [n]} y^{\text{inv}(T)} \quad (4.15)$$

where $\text{inv}(T)$ denotes the number of inversions in T . This polynomial turns out to be related to $C_n(v)$ by the beautiful formula [3, 20, 31, 38]

$$C_n(v) = v^{n-1} I_n(1+v). \quad (4.16)$$

Now, Mallows, Riordan and Kreweras [31, 38] show that $I_n(y)$ satisfies the recursion

$$I_n(y) = \sum_{k=1}^{n-1} \binom{n-2}{k-1} \left(\sum_{j=0}^{k-1} y^j \right) I_k(y) I_{n-k}(y) \quad \text{for } n \geq 2. \quad (4.17)$$

But using (4.16), it is easily seen that (4.17) is equivalent to (4.14).

Let us remark, finally, that even without using (4.15), it follows immediately from the recursion (4.17) [together with the initial condition $I_1(y) = 1$] that the polynomials $I_n(y) \equiv C_n(y-1)/(y-1)^{n-1}$ have nonnegative (indeed, strictly positive) integer coefficients. On the other hand, the nonnegativity of the derivatives of $I_n(y)$ at $y = 0$ is also a special case of Proposition 4.3.

APPENDIX A. A GENERALIZATION OF MÖBIUS INVERSION ON THE PARTITION LATTICE

In this appendix we present an apparently new generalization of Möbius inversion on the lattice of partitions of a finite set, which is inspired by the formulae in Section 3.3.1 and in particular by (3.30), and more generally by (2.13). We refer to [59, Chapter 3] for basic facts about posets and Möbius inversion.

¹⁸See also [5, p. 306, Exercise 4.2.2] for a proof of (4.14) in species language.

To begin with, let P be a finite poset. Then the *zeta function* ζ on P is the function $\zeta: P \times P \rightarrow \mathbb{Z}$ defined by

$$\zeta(x, y) = \begin{cases} 1 & \text{if } x \leq y \\ 0 & \text{if } x \not\leq y \end{cases} \quad (\text{A.1})$$

(We think of ζ as a matrix whose rows and columns are indexed by P .) The *Möbius function* μ on P is the two-sided matrix inverse of ζ , i.e., it satisfies $\zeta\mu = \mu\zeta = I$. It can be computed by the recursion

$$\mu(x, y) = \begin{cases} 1 & \text{if } x = y \\ -\sum_{x \leq z < y} \mu(x, z) & \text{if } x < y \\ 0 & \text{if } x \not\leq y \end{cases} \quad (\text{A.2})$$

Now let S be a finite set, and let P be the lattice $\Pi(S)$ of partitions of S , ordered by refinement. It is well known [59, p. 128] that the Möbius function of $\Pi(S)$ is given by

$$\mu(\sigma, \pi) = \begin{cases} \prod_{i=1}^k (-1)^{\lambda_i-1} (\lambda_i - 1)! & \text{if } \sigma \leq \pi = \{B_1, \dots, B_k\} \text{ and } B_i \text{ is} \\ & \text{partitioned into } \lambda_i \text{ blocks in } \sigma \\ 0 & \text{if } \sigma \not\leq \pi \end{cases} \quad (\text{A.3})$$

We generalize this as follows: let q_1 and q_2 be indeterminates, and define

$$\mu_{q_1, q_2}(\sigma, \pi) = \begin{cases} \prod_{i=1}^k \prod_{j=1}^{\lambda_i-1} (q_2 - jq_1) & \text{if } \sigma \leq \pi = \{B_1, \dots, B_k\} \text{ and } B_i \text{ is} \\ & \text{partitioned into } \lambda_i \text{ blocks in } \sigma \\ 0 & \text{if } \sigma \not\leq \pi \end{cases} \quad (\text{A.4})$$

Let us observe for future reference that

$$\prod_{j=1}^{m-1} (q_2 - jq_1) = q_1^m q_2^{-1} (q_2/q_1)^m = q_1^{m-1} (q_2/q_1 - 1)^{m-1}. \quad (\text{A.5})$$

Then we have in particular

$$\mu_{0,1} = \zeta \quad (\text{A.6a})$$

$$\mu_{1,0} = \mu \quad (\text{A.6b})$$

$$\mu_{q,q} = I \quad \text{for all } q \quad (\text{A.6c})$$

We shall prove the following generalization of Möbius inversion:

Theorem A.1. *Let S be a finite set, and define the matrices μ_{q_1, q_2} on $\Pi(S)$ by (A.4). Then*

$$\mu_{q_1, q_2} \mu_{q_2, q_3} = \mu_{q_1, q_3} \quad (\text{matrix multiplication}) \quad (\text{A.7})$$

We interpret (A.7) as an identity in the polynomial ring $\mathbb{Z}[q_1, q_2, q_3]$, but of course it also holds when q_1, q_2, q_3 are specialized to specific integer values (or more generally to specific values in a commutative ring R). In particular, Möbius inversion corresponds to the special cases $(q_1, q_2, q_3) = (0, 1, 0)$ and $(1, 0, 1)$.

The proof of Theorem A.1 will be based on the following lemma, which we think is of some interest in its own right. It is probably not new, but we have been unable to find any reference.

Lemma A.2. *Let m be a positive integer, and let r and s be indeterminates. Then*

$$\sum_{\omega \in \Pi_m} r^{|\omega|} \left(\prod_{B \in \omega} s^{|B|} \right) = (rs)^m \quad (\text{A.8})$$

as an identity in the polynomial ring $\mathbb{Z}[r, s]$.

Let us remark that if one divides (A.8) by s^m and takes $s \rightarrow \infty$ (or equivalently just extracts the coefficient of s^m), one obtains the well-known formula (3.21)/(3.38); while if one sets $r = q/\epsilon$, $s = \epsilon q$ and takes $\epsilon \rightarrow 0$, then one obtains the formula

$$\sum_{\pi \in \Pi_m} \mu(\hat{0}, \pi) q^{|\pi|} = q^m \quad (\text{A.9})$$

due to Rota [50, Section 9] [59, pp. 128, 162, 187] (here $\hat{0}$ denotes the partition in which every element is a singleton).

Proof of Lemma A.2. It suffices to prove (A.8) for positive integers r, s . Then the right-hand side counts the proper colorings of the complete graph K_m with rs colors. But so does the left-hand side, if we use the color set $[r] \times [s]$ and define ω to be the partition of $[m]$ in which two vertices are placed in the same block if and only if they receive a color with the same first index. \square

Second Proof of Lemma A.2 (suggested independently by Christian Krattenthaler and Richard Stanley). We use the isomorphism of weighted species

$$\text{Partitions} = \text{Sets} \circ \text{NonemptySets}, \quad (\text{A.10})$$

where a Set of cardinality b is given a weight r^b , a NonemptySet of cardinality k is given a weight s^k , and a partition ω is weighted as on the left-hand side of (A.8).¹⁹ The corresponding exponential generating functions are

$$F(x) = \sum_{b=0}^{\infty} \frac{x^b}{b!} r^b = (1+x)^r \quad (\text{A.11})$$

$$G(x) = \sum_{k=1}^{\infty} \frac{x^k}{k!} s^k = (1+x)^s - 1 \quad (\text{A.12})$$

and hence

$$(F \circ G)(x) = (1+x)^{rs} = \sum_{m=0}^{\infty} \frac{x^m}{m!} (rs)^m, \quad (\text{A.13})$$

which proves (A.8) for all $m \geq 0$. \square

¹⁹See [5, pp. 44, 86] for this isomorphism with a different weighting. For the reader unfamiliar with the theory of combinatorial species [5], it suffices to observe that a partition of a finite set S is simply a set of nonempty subsets of S that disjointly cover S ; one can then invoke [60, Theorem 5.1.4] to complete the proof.

Proof of Theorem A.1. Let $\sigma \leq \pi = \{B_1, \dots, B_k\}$ where B_i is partitioned into λ_i blocks in σ , and let us compute

$$(\mu_{q_1, q_2} \mu_{q_2, q_3})(\sigma, \pi) = \sum_{\sigma \leq \tau \leq \pi} \mu_{q_1, q_2}(\sigma, \tau) \mu_{q_2, q_3}(\tau, \pi). \quad (\text{A.14})$$

The partition τ is specified by saying, for each $i \in [k]$, how the corresponding λ_i blocks of σ get grouped in τ . The sum (A.14) will then factorize over i , with the i th factor given by

$$\begin{aligned} & \sum_{\substack{\omega \in \Pi_{\lambda_i} \\ \omega = \{\hat{B}_1, \dots, \hat{B}_{\mu_i}\}}} \left(\prod_{\alpha=1}^{\mu_i} \prod_{j=1}^{|\hat{B}_\alpha|-1} (q_2 - jq_1) \right) \prod_{k=1}^{\mu_i-1} (q_3 - kq_2) \\ &= \sum_{\substack{\omega \in \Pi_{\lambda_i} \\ \omega = \{\hat{B}_1, \dots, \hat{B}_{\mu_i}\}}} \left(\prod_{\alpha=1}^{\mu_i} q_1^{|\hat{B}_\alpha|} q_2^{-1} (q_2/q_1)^{|\hat{B}_\alpha|} \right) q_2^{\mu_i} q_3^{-1} (q_3/q_2)^{\mu_i} \\ &= q_1^{\lambda_i} q_3^{-1} \sum_{\substack{\omega \in \Pi_{\lambda_i} \\ \omega = \{\hat{B}_1, \dots, \hat{B}_{\mu_i}\}}} \left(\prod_{\alpha=1}^{\mu_i} (q_2/q_1)^{|\hat{B}_\alpha|} \right) (q_3/q_2)^{\mu_i} \\ &= q_1^{\lambda_i} q_3^{-1} (q_3/q_1)^{\lambda_i} \end{aligned} \quad (\text{A.15})$$

by Lemma A.2. \square

Second Proof of Theorem A.1 (suggested by Richard Stanley). We prove (A.7) for $S = [n]$ simultaneously for all $n \geq 1$, using the interpretation of convolution of multiplicative functions on $\mathbf{\Pi} = (\Pi_1, \Pi_2, \dots)$ in terms of composition of exponential generating functions [60, pp. 7–8]. We first observe that μ_{q_1, q_2} is a multiplicative function on $\mathbf{\Pi}$, with underlying numerical function

$$f_{q_1, q_2}(m) = \prod_{j=1}^{m-1} (q_2 - jq_1) = q_1^m q_2^{-1} (q_2/q_1)^m \quad (\text{A.16})$$

and corresponding exponential generating function

$$F_{q_1, q_2}(x) = \sum_{m=1}^{\infty} f_{q_1, q_2}(m) \frac{x^m}{m!} = q_2^{-1} [(1 + q_1 x)^{q_2/q_1} - 1]. \quad (\text{A.17})$$

It is easy to check that

$$F_{q_2, q_3}(F_{q_1, q_2}(x)) = F_{q_1, q_3}(x). \quad (\text{A.18})$$

By [60, Theorem 5.1.11], the identities (A.7) for all $n \geq 1$ are an immediate consequence.²⁰ \square

²⁰Using [60, Theorem 5.1.11] with the coefficient field K taken to be (say) \mathbb{R} , this argument proves (A.7) for $q_1, q_2, q_3 \in \mathbb{R} \setminus \{0\}$, which is more than enough to imply it as a polynomial identity. Alternatively, we can use [60, Theorem 5.1.11] with K taken to be (say) the field $\mathbb{R}(q_1, q_2, q_3)$ of rational functions in the indeterminates q_1, q_2, q_3 , which yields (A.7) directly as a polynomial identity.

Remark. If we define the diagonal matrix D_r by

$$D_r(\sigma, \pi) = \begin{cases} r^{|\sigma|} & \text{if } \sigma = \pi \\ 0 & \text{if } \sigma \neq \pi \end{cases} \quad (\text{A.19})$$

then an easy calculation shows that

$$D_r \mu_{q_1, q_2} D_r^{-1} = \mu_{rq_1, rq_2}. \quad (\text{A.20})$$

This is why many things involving μ_{q_1, q_2} depend only on the ratio q_2/q_1 [cf. (3.28)].

We can apply Theorem A.1 as follows: Let R be a commutative ring, and fix a function $f: \Pi(S) \rightarrow R$ and an element $q_0 \in R$ (it could be an indeterminate if desired). Now introduce a new indeterminate q , and define $F_q: \Pi(S) \rightarrow R[q]$ by

$$F_q = f \mu_{q_0, q} \quad (\text{A.21})$$

or in more detail

$$F_q(\pi) = \sum_{\sigma \in \Pi(S)} f(\sigma) \mu_{q_0, q}(\sigma, \pi). \quad (\text{A.22})$$

By construction we have imposed the “initial condition”

$$F_{q_0} = f \quad (\text{A.23})$$

(here F_{q_0} means of course that the indeterminate q is replaced by the value $q_0 \in R$). Most importantly, Theorem A.1 implies that the q -dependence of F_q is “coherent” in the sense that

$$F_{q_1} \mu_{q_1, q_2} = F_{q_2}. \quad (\text{A.24})$$

The formula (3.30) is now the special case of (A.24) in which we take $S = V$,

$$F_q(\pi) = \prod_{B \in \pi} \widehat{Z}_{G[B]}(q, \mathbf{v}), \quad (\text{A.25})$$

and we evaluate (A.24) at $\pi = \hat{1}$ (the partition in which all of V belongs to a single block). More generally, in place of $\widehat{Z}_{G[B]}(q, \mathbf{v})$ we can use $\widehat{a}_{\mathbf{1}_B}(q)$ where $\{\widehat{a}_{\mathbf{n}}(q)\}$ is *any* family arising as in Section 2; we then obtain (2.13) specialized to $\mathbf{n} = \mathbf{1}$.²¹

The formulae in this appendix seem to be special for the partition lattice. Still, it may not be totally absurd to ask: Might there exist “ q_1 – q_2 generalizations” of Möbius inversion for some other posets? Maybe even for some natural *class* of posets?

ACKNOWLEDGMENTS

We are grateful to Pierre Leroux for an extremely careful reading of the manuscript and for many suggestions, both substantive and bibliographic. In particular, it is thanks to him that our attention was drawn to [5, Section 3.1] and notably to [5, p. 189, Exercise 3.1.22], from which we learned about the Abel-type extensions (1.16)–(1.19).

We finished the first version of this paper, by unhappy circumstance, on the same day that we learned that Pierre had been struck with a grave illness, to which he succumbed two weeks later. We dedicate this work to his memory.

²¹Note also that, in these formulae, we use $\widehat{a}_{\mathbf{n}}(q)$ [or $c_{\mathbf{n}}$] only for $\mathbf{0} \leq \mathbf{n} \leq \mathbf{1}$.

Finally, we are grateful to an anonymous referee for pointing out the connection of the formulae of Section 3.3.1 with Möbius inversion — which led us to discover the results reported in the Appendix — and to Christian Krattenthaler and Richard Stanley for comments on these results.

This research was supported in part by U.S. National Science Foundation grant PHY-0424082, by U.K. Engineering and Physical Sciences Research Council grant GR/S26323/01, and by a Leverhulme Visiting Professorship.

REFERENCES

- [1] G.E. Andrews, R. Askey and R. Roy, *Special Functions* (Cambridge University Press, Cambridge–New York, 1999), Section 6.2.
- [2] W. Magnus, F. Oberhettinger, and F.G. Tricomi, *Higher Transcendental Functions*, vol. 2 (McGraw-Hill, New York, 1953).
- [3] J.S. Beissinger, On external activity and inversions in trees, *J. Combin. Theory B* **33**, 87–92 (1982).
- [4] G.D. Birkhoff, A determinant formula for the number of ways of coloring a map, *Ann. Math.* **14**, 42–46 (1912).
- [5] F. Bergeron, G. Labelle and P. Leroux, *Combinatorial Species and Tree-Like Structures* (Cambridge University Press, Cambridge–New York, 1998).
- [6] N. Biggs, *Algebraic Graph Theory*, 2nd ed. (Cambridge University Press, Cambridge–New York, 1993), p. 103, item 13e.
- [7] A. Björner, The homology and shellability of matroids and geometric lattices, in N. White (editor), *Matroid Applications* (Encyclopedia of Mathematics and its Applications #40), Chapter 7, pp. 226–283 (Cambridge University Press, Cambridge, 1992).
- [8] C. Borgs, Polymer systems, Pirogov–Sinai theory and conductance bounds. Lecture notes, Microsoft Research, October 2000, unpublished, cited in [9, p. 68].
- [9] C. Borgs, Absence of zeros for the chromatic polynomial of bounded degree graphs, *Combin. Probab. Comput.* **15**, 63–74 (2006).
- [10] C. Chauve, P. Duchon and S. Dulucq, Une généralisation des résultats de Kreweras sur le polynôme énumérateur des inversions dans les arborescences, *Ann. Sci. Math. Québec* **24**, 129–144 (2000).
- [11] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions* (D. Reidel, Dordrecht–Boston, 1974), Section 3.3.
- [12] J.P. Fillmore and S.G. Williamson, A linear algebra setting for the Rota–Mullin theory of polynomials of binomial type, *Linear and Multilinear Algebra* **1**, 67–80 (1973).
- [13] D. Foata, *La série génératrice exponentielle dans les problèmes d'énumération* (Presses de l'Université de Montréal, Montréal, 1974).
- [14] D. Foata and M.-P. Schützenberger, *Théorie géométrique des polynômes eulériens*, Lecture Notes in Mathematics #138 (Springer-Verlag, Berlin–New York, 1970), Chapitre III.
- [15] C.M. Fortuin and P.W. Kasteleyn, On the random-cluster model. I. Introduction and relation to other models, *Physica* **57**, 536–564 (1972).
- [16] A.M. Garsia, An exposé of the Mullin–Rota theory of polynomials of binomial type, *Linear and Multilinear Algebra* **1**, 47–65 (1973).
- [17] I.M. Gessel, Enumerative applications of a decomposition for graphs and digraphs, *Discrete Math.* **139**, 257–271 (1995).
- [18] I.M. Gessel, Applications of the classical umbral calculus, *Algebra Univers.* **49**, 397–434 (2003).
- [19] I.M. Gessel and B.E. Sagan, The Tutte polynomial of a graph, depth-first search, and simplicial complex partitions, *Electron. J. Combin.* **3**, no. 2, #R9 (1996).
- [20] I. Gessel and D.-L. Wang, Depth-first search as a combinatorial correspondence, *J. Combin. Theory A* **26**, 308–313 (1979).
- [21] H.W. Gould, Some generalizations of Vandermonde's convolution, *Amer. Math. Monthly* **63**, 84–91 (1956).

- [22] H.W. Gould, Final analysis of Vandermonde's convolution, *Amer. Math. Monthly* **64**, 409–415 (1957).
- [23] H.W. Gould, Coefficient identities for powers of Taylor and Dirichlet series, *Amer. Math. Monthly* **81**, 3–14 (1974).
- [24] H.W. Gould and J. Kaucký, Evaluation of a class of binomial coefficient summations, *J. Comb. Theory* **1**, 233–247 (1966); erratum, *J. Comb. Theory A* **12**, 309–310 (1972).
- [25] R.L. Graham, D.E. Knuth and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Science*, 2nd ed. (Addison-Wesley, Reading, Mass., 1994).
- [26] G. Grimmett, *The Random-Cluster Model* (Springer-Verlag, New York, 2006).
- [27] F. Harary and E.M. Palmer, *Graphical Enumeration* (Academic Press, New York, 1973).
- [28] W.P. Johnson, The Pfaff/Cauchy derivative identities and Hurwitz type extensions, *Ramanujan J.* **13**, 167–201 (2007).
- [29] P.W. Kasteleyn and C.M. Fortuin, Phase transitions in lattice systems with random local properties, *J. Phys. Soc. Japan* **26** (Suppl.), 11–14 (1969).
- [30] D.E. Knuth, Convolution polynomials, *Mathematica J.* **2**, 67–78 (1992); [arXiv:math.CA/9207221](https://arxiv.org/abs/math.CA/9207221).
- [31] G. Kreweras, Une famille de polynômes ayant plusieurs propriétés énumératives, *Period. Math. Hungar.* **11**, 309–320 (1980).
- [32] G. Labelle, Sur l'inversion et l'itération continue des séries formelles, *Europ. J. Combin.* **1**, 113–138 (1980).
- [33] G. Labelle, Une nouvelle démonstration combinatoire des formules d'inversion de Lagrange, *Adv. Math.* **42**, 217–247 (1981).
- [34] G. Labelle and P. Leroux, An extension of the exponential formula in enumerative combinatorics, *Electron. J. Combin.* **3**, no. 2, #R12 (1996).
- [35] B. Lass, Orientations acycliques et le polynôme chromatique, *European J. Combin.* **22**, 1101–1123 (2001).
- [36] N.N. Lebedev, *Special Functions and Their Applications* (Prentice-Hall, Englewood Cliffs, N.J., 1965).
- [37] P. Leroux, Methoden der Anzahlbestimmung für einige Klassen von Graphen [Methods of enumeration for some classes of graphs], *Bayreuth. Math. Schr.* **26**, 1–36 (1988).
- [38] C.L. Mallows and J. Riordan, The inversion enumerator for labeled trees, *Bull. Amer. Math. Soc.* **74**, 92–94 (1968).
- [39] R. Mullin and G.-C. Rota, On the foundations of combinatorial theory. III. Theory of binomial enumeration, in *Graph Theory and its Applications*, edited by Bernard Harris (Academic Press, New York, 1970), pp. 167–213.
- [40] A. Nijenhuis and H.S. Wilf, The enumeration of connected graphs and linked diagrams, *J. Combin. Theory A* **27**, 356–359 (1979).
- [41] O. Penrose, Convergence of fugacity expansions for classical systems, in T.A. Bak (editor), *Statistical Mechanics: Foundations and Applications*, pp. 101–109 (Benjamin, New York–Amsterdam, 1967).
- [42] J. Pitman, Random mappings, forests, and subsets associated with Abel–Cayley–Hurwitz multinomial expansions, *Sém. Lothar. Combin.* **46**, article B46h (2001).
- [43] J. Pitman, Forest volume decompositions and Abel–Cayley–Hurwitz multinomial expansions, *J. Combin. Theory A* **98**, 175–191 (2002).
- [44] R.B. Potts, Some generalized order-disorder transformations, *Proc. Cambridge Philos. Soc.* **48**, 106–109 (1952).
- [45] D.L. Reiner, Multivariate sequences of binomial type, *Stud. Appl. Math.* **57**, 119–133 (1977).
- [46] J. Riordan, *Combinatorial Identities* (Wiley, New York, 1968), especially Sections 1.5 and 1.6.
- [47] J. Riordan, unpublished, cited in [27, p. 8] and [40, pp. 356–357].
- [48] S.M. Roman and G.-C. Rota, The umbral calculus, *Adv. Math.* **27**, 95–188 (1978).
- [49] S.M. Roman, *The Umbral Calculus* (Academic Press, New York, 1984).
- [50] G.-C. Rota, On the foundations of combinatorial theory. I. Theory of Möbius functions, *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* **2**, 340–368 (1964).

- [51] G.-C. Rota, D. Kahaner and A. Odlyzko, On the foundations of combinatorial theory. VIII. Finite operator calculus, *J. Math. Anal. Appl.* **42**, 684–760 (1973). [Reprinted in G.-C. Rota, *Finite Operator Calculus* (Academic Press, New York–London, 1975), Chapter 2.]
- [52] G.-C. Rota and B.D. Taylor, The classical umbral calculus, *SIAM J. Math. Anal.* **25**, 694–711 (1994).
- [53] A.D. Scott and A.D. Sokal, The repulsive lattice gas, the independent-set polynomial, and the Lovász local lemma, *J. Stat. Phys.* **118**, 1151–1261 (2005); [arXiv:cond-mat/0309352](#).
- [54] A.D. Scott and A.D. Sokal, On dependency graphs and the lattice gas, *Combin. Probab. Comput.* **15**, 253–279 (2006).
- [55] A.D. Scott and A.D. Sokal, Asymptotic behavior at complex parameter for the Tutte polynomials (alias Potts-model partition functions) of the complete graphs K_n , in preparation.
- [56] A.D. Sokal, Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions, *Combin. Probab. Comput.* **10**, 41–77 (2001); [arXiv:cond-mat/9904146](#).
- [57] A.D. Sokal, The multivariate Tutte polynomial (alias Potts model) for graphs and matroids, in *Surveys in Combinatorics, 2005*, edited by Bridget S. Webb (Cambridge University Press, Cambridge–New York, 2005), pp. 173–226; [arXiv:math.CO/0503607](#).
- [58] A.J. Stam, Two identities in the theory of polynomials of binomial type, *J. Math. Anal. Appl.* **122**, 439–443 (1987).
- [59] R.P. Stanley, *Enumerative Combinatorics*, vol. 1 (Wadsworth & Brooks/Cole, Monterey, California, 1986). Reprinted by Cambridge University Press, 1997.
- [60] R.P. Stanley, *Enumerative Combinatorics*, vol. 2 (Cambridge University Press, Cambridge–New York, 1999).
- [61] V. Strehl, Identities of Rothe–Abel–Schläfli–Hurwitz-type, *Discrete Math.* **99**, 321–340 (1992).
- [62] J.R. Swenson, The chromatic polynomial of a complete bipartite graph, *Amer. Math. Monthly* **80**, 797–798 (1973).
- [63] W.T. Tutte, On dichromatic polynomials, *J. Combin. Theory* **2**, 301–320 (1967).
- [64] G.E. Uhlenbeck and G.W. Ford, The theory of linear graphs with applications to the theory of the virial development of the properties of gases, in J. de Boer and G.E. Uhlenbeck (editors), *Studies in Statistical Mechanics*, vol. I, pp. 119–211 (North-Holland, Amsterdam, 1962).
- [65] D.J.A. Welsh and C. Merino, The Potts model and the Tutte polynomial, *J. Math. Phys.* **41**, 1127–1152 (2000).
- [66] H. Whitney, A logical expansion in mathematics, *Bull. Amer. Math. Soc.* **38**, 572–579 (1932).
- [67] H.S. Wilf, *generatingfunctionology*, 2nd ed. (Academic Press, San Diego–London, 1994), Chapter 3.
- [68] F.Y. Wu, The Potts model, *Rev. Mod. Phys.* **54**, 235–268 (1982); erratum **55**, 315 (1983).
- [69] F.Y. Wu, Potts model of magnetism (invited), *J. Appl. Phys.* **55**, 2421–2425 (1984).
- [70] J. Zeng, Multinomial convolution polynomials, *Discrete Math.* **160**, 219–228 (1996).