

# On some polynomials enumerating Fully Packed Loop configurations



Philippe Nadeau (Univ. of Vienna)

Joint work with Tiago Fonseca (LPTHE, Univ. Paris 6)

SLC 64, Lyon, March 29th 2010.

# Introduction


Fully Packed Loop (FPL) configurations are combinatorial structures that arose in statistical mechanics, in simple bijection with known combinatorial objects.



$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 1 \\ 1 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

# Introduction

Fully Packed Loop (FPL) configurations are combinatorial structures that arose in statistical mechanics, in simple bijection with known combinatorial objects.




$$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 & 1 \\ 1 & 0 & 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}$$

One can define certain polynomials  $A_\pi(X)$  (indexed by noncrossing matchings  $\pi$ ) which count FPLs when specialized to nonnegative integers.

We will here formulate conjectures for the polynomials  $A_\pi(X)$ , hinting at “combinatorial reciprocity”.

# Introduction

Start with the square grid  $G_n$  with  $n^2$  vertices and  $4n$  external edges (here  $n = 7$ ).



A **FPL** configuration of size  $n$  is a subgraph of the grid  $G_n$


- (1) such that around each vertex of  $G_n$ , 2 edges out of 4 are selected; (“**Fully Packed**”)
- (2) containing every other external edge. (“**Boundary condition**”)

$A_n$  = number of FPLs of size  $n$

# Introduction


A **link pattern**  $\pi$  of size  $n$  is a set of  $n$  noncrossing chords between  $2n$  points on a disk.

To each FPL  $F$  is associated a link pattern  $\pi(F)$ .



# Introduction


We will write link patterns linearly, as noncrossing matchings :



Given a noncrossing matching  $\pi$ , we note  $A_\pi$  the number of FPLs  $F$  such that  $\pi(F) = \pi$ .


# Introduction

We will write link patterns linearly, as noncrossing matchings :




Given a noncrossing matching  $\pi$ , we note  $A_\pi$  the number of FPLs  $F$  such that  $\pi(F) = \pi$ .

Matchings with **nested arches** : for  $p \geq 0$ ,



# Introduction : the Razumov-Stroganov ex-conjecture


**Definition** : The operators  $e_i, i \in \llbracket 1, 2n \rrbracket$ , act on matchings by  $\{i, j\}, \{i + 1, k\} \in \pi \rightarrow \{i, i + 1\}, \{j, k\} \in e_i(\pi)$ .





# Introduction : the Razumov-Stroganov ex-conjecture

**Definition** : The operators  $e_i, i \in \llbracket 1, 2n \rrbracket$ , act on matchings by  $\{i, j\}, \{i + 1, k\} \in \pi \rightarrow \{i, i + 1\}, \{j, k\} \in e_i(\pi)$ .




## Markov chain $\mathcal{M}$

- States =  $LP_n$  ;
- Transition probabilities :  $P(\pi \rightarrow \pi') = \frac{k}{2n}$  where  $k$  is the number of  $i \in \{1, \dots, 2n\}$  such that  $e_i(\pi) = \pi'$ .

# Introduction : the Razumov-Stroganov ex-conjecture

**Definition** : The operators  $e_i, i \in \llbracket 1, 2n \rrbracket$ , act on matchings by  $\{i, j\}, \{i + 1, k\} \in \pi \rightarrow \{i, i + 1\}, \{j, k\} \in e_i(\pi)$ .



## Markov chain $\mathcal{M}$

- States =  $LP_n$  ;
- Transition probabilities :  $P(\pi \rightarrow \pi') = \frac{k}{2n}$  where  $k$  is the number of  $i \in \{1, \dots, 2n\}$  such that  $e_i(\pi) = \pi'$ .

**Stationary distribution** : Let  $P$  be the matrix defined by

$P_{\pi\pi'} = P(\pi \rightarrow \pi')$  where  $\pi, \pi' \in LP_n$ .

Then there is a **unique probability distribution**  $(\psi)_\pi$  on  $LP_n$  such that  $P\psi = \psi$ .

# Introduction : the Razumov-Stroganov ex-conjecture

**RS conjecture ([Cantini and Sportiello '10]) :**

$$\forall \pi \in LP_n, \quad \psi_\pi = \frac{A_\pi}{A_n}$$

The proof consists in showing that the numbers  $A_\pi/A_n$  verify the stationary equations of  $\mathcal{M}$ , which can be written explicitly :

$$\forall \pi, \quad 2nA_\pi = \sum_{(i, \pi'), e_i(\pi') = \pi} A_{\pi'}$$

# Introduction : the Razumov-Stroganov ex-conjecture

**RS conjecture ([Cantini and Sportiello '10]) :**

$$\forall \pi \in LP_n, \quad \psi_\pi = \frac{A_\pi}{A_n}$$

The proof consists in showing that the numbers  $A_\pi/A_n$  verify the stationary equations of  $\mathcal{M}$ , which can be written explicitly :

$$\forall \pi, \quad 2nA_\pi = \sum_{(i, \pi'), e_i(\pi')=\pi} A_{\pi'}$$

The numbers  $\psi_\pi$  were studied in detail by Di Francesco and Zinn-Justin  $\rightarrow$  integral expressions (up to a change of basis), multivariate versions, computations in special cases.

# Introduction : the Razumov-Stroganov ex-conjecture

**RS conjecture ([Cantini and Sportiello '10]) :**

$$\forall \pi \in LP_n, \quad \psi_\pi = \frac{A_\pi}{A_n}$$

The proof consists in showing that the numbers  $A_\pi/A_n$  verify the stationary equations of  $\mathcal{M}$ , which can be written explicitly :

$$\forall \pi, \quad 2nA_\pi = \sum_{(i, \pi'), e_i(\pi')=\pi} A_{\pi'}$$

The numbers  $\psi_\pi$  were studied in detail by Di Francesco and Zinn-Justin  $\rightarrow$  integral expressions (up to a change of basis), multivariate versions, computations in special cases.

**Consequence** : to prove results about the numbers  $A_\pi$  one can either use their combinatorial definitions or use the expressions from Di-Francesco and Zinn-Justin.

# Introduction

**Theorem**[Caselli, Krattenthaler, Lass, N. '05]



For a fixed  $\pi$ , the quantity  $A_{(\pi)_p}$  is polynomial in  $p$ ; let  $A_\pi(X)$  be the polynomial such that  $A_\pi(p) = A_{(\pi)_p}$  for  $p \in \mathbb{N}$ . Then  $A_\pi(X)$  is a polynomial of degree  $d(\pi)$ , with leading coefficient  $\frac{1}{H_\pi}$ , such that  $d(\pi)! \cdot A_\pi(X)$  has integer coefficients.

# Introduction

**Theorem**[Caselli, Krattenthaler, Lass, N. '05]

For a fixed  $\pi$ , the quantity  $A_{(\pi)_p}$  is polynomial in  $p$ ; let  $A_\pi(X)$  be the polynomial such that  $A_\pi(p) = A_{(\pi)_p}$  for  $p \in \mathbb{N}$ . Then  $A_\pi(X)$  is a polynomial of degree  $d(\pi)$ , with leading coefficient  $\frac{1}{H_\pi}$ , such that  $d(\pi)! \cdot A_\pi(X)$  has integer coefficients.

$d(\pi)$  is the number of boxes in the Young diagram  $Y(\pi)$  :




# Introduction


**Theorem**[Caselli, Krattenthaler, Lass, N. '05]

For a fixed  $\pi$ , the quantity  $A_{(\pi)_p}$  is polynomial in  $p$ ; let  $A_\pi(X)$  be the polynomial such that  $A_\pi(p) = A_{(\pi)_p}$  for  $p \in \mathbb{N}$ . Then  $A_\pi(X)$  is a polynomial of degree  $d(\pi)$ , with leading coefficient  $\frac{1}{H_\pi}$ , such that  $d(\pi)! \cdot A_\pi(X)$  has integer coefficients.

$d(\pi)$  is the number of boxes in the Young diagram  $Y(\pi)$  :



$H_\pi$  is the product of the hook lengths of  $Y(\pi)$ .





# Introduction

We have the following properties :

$$A_{\pi}(X) = A_{\pi^*}(X).$$
$$A_{(\pi)}(X) = A_{\pi}(X + 1).$$

Here  $\pi^*$  is the matching reflected vertically ; this is proved by checking the evaluations  $X = p \in \mathbb{N}$ .

# Introduction

We have the following properties :

$$A_{\pi}(X) = A_{\pi^*}(X).$$
$$A_{(\pi)}(X) = A_{\pi}(X + 1).$$

Here  $\pi^*$  is the matching reflected vertically ; this is proved by checking the evaluations  $X = p \in \mathbb{N}$ .

We formulate several conjectures about the polynomials  $A_{\pi}(X)$ , and emphasize the combinatorics related to them.

# Some simple combinatorics

# Combinatorial constructions (1)

Let  $\pi$  be a matching of size  $|\pi| = n$ , and consider an integer  $i$  in  $\llbracket 1, n-1 \rrbracket$ .

Let  $\widehat{x} = 2n + 1 - x$ , and  $(I)$  be the interval  $\llbracket i + 1, \widehat{i + 1} \rrbracket$ , while  $(O)$  is defined as  $\llbracket 1, 2n \rrbracket - (I)$ .


**Definition**  $m_i(\pi) :=$  half the number of arcs in  $\pi$  linking the regions  $(O)$  and  $(I)$ .

# Combinatorial constructions (1)

Let  $\pi$  be a matching of size  $|\pi| = n$ , and consider an integer  $i$  in  $\llbracket 1, n-1 \rrbracket$ .

Let  $\hat{x} = 2n + 1 - x$ , and  $(I)$  be the interval  $\llbracket i + 1, \widehat{i + 1} \rrbracket$ , while  $(O)$  is defined as  $\llbracket 1, 2n \rrbracket - (I)$ .


**Definition**  $m_i(\pi) :=$  half the number of arcs in  $\pi$  linking the regions  $(O)$  and  $(I)$ .



$$m_4(\pi_0) = \frac{4}{2} = 2.$$

# Combinatorial constructions (1)


This is better visualized by folding  $\pi$  on itself :



We obtain  $m_i(\pi_0) = 0, 1, 2, 2, 2, 1, 1$  for  $i = 1 \dots 7$ .

# Combinatorial constructions (1)

This is better visualized by folding  $\pi$  on itself :




We obtain  $m_i(\pi_0) = 0, 1, 2, 2, 2, 1, 1$  for  $i = 1 \dots 7$ .

One has the easy properties :

- $m_i(\pi) = m_i(\pi^*)$ , where  $\pi^*$  is the reflected matching ;
- $m_i(\pi) = m_{i+1}(\pi')$  where  $\pi' = (\pi)$ .

# Combinatorial constructions (1)

This is better visualized by folding  $\pi$  on itself :



We obtain  $m_i(\pi_0) = 0, 1, 2, 2, 2, 1, 1$  for  $i = 1 \dots 7$ .

One has the easy properties :


- $m_i(\pi) = m_i(\pi^*)$ , where  $\pi^*$  is the reflected matching ;
- $m_i(\pi) = m_{i+1}(\pi')$  where  $\pi' = (\pi)$ .

We now define other integers  $m_i^{bis}(\pi)$ , defined also for  $i = 1, 2, \dots, n - 1$ .



## Combinatorial constructions (2)

We use here the Young diagram  $Y(\pi)$  attached to  $\pi$ .



## Combinatorial constructions (2)

We use here the Young diagram  $Y(\pi)$  attached to  $\pi$ .


1) Label the cells by putting  $n - 1$  in the top left corner, and letting labels decrease by 1; decompose  $Y(\pi)$  in rims.

7	6	5	4	3
6	5	4	3	
5	4	3	2	
4	3			

## Combinatorial constructions (2)

We use here the Young diagram  $Y(\pi)$  attached to  $\pi$ .

- 1) Label the cells by putting  $n - 1$  in the top left corner, and letting labels decrease by 1; decompose  $Y(\pi)$  in rims.
- 2) For each rim  $R$ , construct the multiset union of  $\{k\}$ ,  $\llbracket k + 1, e_1 \rrbracket$  and  $\llbracket k + 1, e_2 \rrbracket$ , where  $k$  is the minimum label in  $R$ , and  $e_1, e_2$  are the labels at both extremities.




## Combinatorial constructions (2)

We use here the Young diagram  $Y(\pi)$  attached to  $\pi$ .

- 1) Label the cells by putting  $n - 1$  in the top left corner, and letting labels decrease by 1; decompose  $Y(\pi)$  in rims.
- 2) For each rim  $R$ , construct the multiset union of  $\{k\}$ ,  $\llbracket k + 1, e_1 \rrbracket$  and  $\llbracket k + 1, e_2 \rrbracket$ , where  $k$  is the minimum label in  $R$ , and  $e_1, e_2$  are the labels at both extremities.
- 3) Now do the union  $U$  of all these multisets :

$$m_i^{bis}(\pi) := \text{multiplicity of } i \text{ in } U.$$



$$U = \{2, 3^2, 4^2, 5^2, 6, 7\}$$

$$\rightarrow m_i^{bis} = 0, 1, 2, 2, 2, 1, 1$$

for  $i = 1 \dots 7$ .

## Combinatorial constructions (2)

In fact we have :

**Proposition :** For all  $\pi, i$ , we have  $m_i(\pi) = m_i^{bis}(\pi)$

One can show this by induction on the number of rims in the rim decomposition of  $Y(\pi)$ .

## Combinatorial constructions (2)

In fact we have :

**Proposition :** For all  $\pi, i$ , we have  $m_i(\pi) = m_i^{bis}(\pi)$

One can show this by induction on the number of rims in the rim decomposition of  $Y(\pi)$ .

**Proposition :** For any  $\pi$ , we have

- $\sum_i m_i(\pi) \leq d(\pi)$  ;
- $\sum_i m_i(\pi) \equiv d(\pi) \pmod{2}$ .

*Proof :* Use the  $m_i^{bis}(\pi)$  construction.

By convention, we set  $m_i(\pi) = 0$  if  $i \in \llbracket 1, n - 1 \rrbracket$ .

# The conjectures

# The conjectures

We will formulate conjectures about the polynomials  $A_\pi(X)$ , which concern :

- the real roots ;
- the values at negative integers ;
- the coefficients.



# The conjectures

We will formulate conjectures about the polynomials  $A_\pi(X)$ , which concern :

- the real roots ;
- the values at negative integers ;
- the coefficients.

The  $A_\pi(X)$  have been explicitly computed for  $|\pi| \leq 8$ , and the conjectures hold for these polynomials.

(There are  $C_8 = 1430$  matchings  $\pi$  with 8 arches. The maximal degree  $d(\pi)$  of the corresponding polynomials  $A_\pi(X)$  is 28.)

# The root conjecture

**Root conjecture :** All real roots of  $A_\pi(X)$  are negative integers. The multiplicity of  $-i$  is exactly  $m_i(\pi)$ .

Equivalently, we have the factorization :

$$A_\pi(X) = \left( \prod_i (X + i)^{m_i(\pi)} \right) \cdot Q_\pi(X)$$

where  $Q_\pi(X)$  has no real roots.

# The root conjecture

**Root conjecture :** All real roots of  $A_\pi(X)$  are negative integers. The multiplicity of  $-i$  is exactly  $m_i(\pi)$ .

Equivalently, we have the factorization :

$$A_\pi(X) = \left( \prod_i (X + i)^{m_i(\pi)} \right) \cdot Q_\pi(X)$$

where  $Q_\pi(X)$  has no real roots.

**(Remark :** A consequence of the conjecture is that it gives the *sign variations* of the real function  $x \mapsto A_\pi(x)$ .)

**Proposition :** If  $(1, 2n)$  is not an arc in  $\pi$ , then  $A_\pi(-1) = 0$ .

This is a very special case of the conjecture.

# The root conjecture

We need to check that the root conjecture is compatible with what we know about  $A_\pi(X)$  and  $m_i(\pi)$ .

1)  $A_\pi(X)$  has degree  $d(\pi)$ , so  $\sum_i m_i(\pi)$  cannot be larger than  $d(\pi)$ . Furthermore  $Q_\pi(X)$  has even degree, so  $d(\pi) - \sum_i m_i(\pi)$  is even.

$\Rightarrow$  We already checked both facts.

2) We have  $A_\pi(X) = A_{\pi^*}(X)$ .

$\Rightarrow$  indeed  $m_i(\pi) = m_i(\pi^*)$  for all  $i$ .

3)  $A_{(\pi)}(X) = A_\pi(X + 1)$

$\Rightarrow$  we have  $m_i((\pi)) = m_{i+1}(\pi)$  as expected.

# The ghost value conjecture

1) **Definition** For any  $\pi$  we define  $G_\pi := A_\pi(-|\pi|)$ .

By the root conjecture, its sign is  $(-1)^{d(\pi)}$ .


2) **Composition of matchings**



$\alpha$



$\beta$



$\alpha \circ \beta$

# The ghost value conjecture

1) **Definition** For any  $\pi$  we define  $G_\pi := A_\pi(-|\pi|)$ .

By the root conjecture, its sign is  $(-1)^{d(\pi)}$ .


2) **Composition of matchings**



$\alpha$



$\beta$



$\alpha \circ \beta$

**Ghost value conjecture :** Let  $i \in \llbracket 1, n - 1 \rrbracket$  such that  $m_i(\pi) = 0$ , and write  $\pi = \alpha \circ \beta$  where  $|\alpha| = i$ ,  $|\beta| = n - i$ .  
Then

$$A_\pi(-i) = G_\alpha A_\beta.$$

# The ghost value conjecture

The values  $G_\pi$  play thus a special role, and we have conjectures for them :

**Conjecture** : For all  $n \geq 1$ , we have

$$\sum_{\pi:|\pi|=n} |G_\pi| = A_n$$

Here  $A_n$  is the total number of FPLs of size  $n$ .

So, like the  $A_\pi$ , the values  $G_\pi$  seem to be associated to a partition of FPLs indexed by non-crossing matchings ( It is easily checked that  $A_\pi \neq G_\pi$  in general).

# The positivity conjecture

**Positivity conjecture** : The polynomial  $A_\pi(X)$  has nonnegative coefficients.

This is already known for :

- the constant term  $A_\pi(0) = A_\pi$ , and
- the leading coefficient in degree  $d(\pi)$  which is  $1/H_\pi$ .



# The positivity conjecture

**Positivity conjecture** : The polynomial  $A_\pi(X)$  has nonnegative coefficients.

This is already known for :

- the constant term  $A_\pi(0) = A_\pi$ , and
- the leading coefficient in degree  $d(\pi)$  which is  $1/H_\pi$ .

**Theorem** The positivity conjecture holds for the coefficient of  $X^{d(\pi)-1}$  in  $A_\pi(X)$ .

Two proofs can be given : either using the formula for  $\psi_\pi$ , or an expression for the polynomials  $A_\pi(X)$  based on “FPL configurations in a triangle” ([N. '10]).

**Remark** : As a byproduct of these proofs, we obtained certain summation formulas involving hook products  $H_\pi$ .

## A last word of support

We sum up the various sources of supporting evidence for the conjectures :

- Computation of the  $A_\pi(X)$  for small  $|\pi|$  ;
- Compatibility of the conjectures with known facts ;
- Coherence of the conjectures among themselves.
- Proof of the conjectures for special values ;
- Proof of the conjectures for certain families of matchings  $\pi$ , for which  $A_\pi(X)$  is known explicitly [Di-Francesco and al '04, Caselli and Krattenthaler '05].

## A last word of support

We sum up the various sources of supporting evidence for the conjectures :

- Computation of the  $A_\pi(X)$  for small  $|\pi|$  ;
- Compatibility of the conjectures with known facts ;
- Coherence of the conjectures among themselves.
- Proof of the conjectures for special values ;
- Proof of the conjectures for certain families of matchings  $\pi$ , for which  $A_\pi(X)$  is known explicitly [Di-Francesco and al '04, Caselli and Krattenthaler '05].

I wish this were enough to turn them into theorems ; it's not...  
But anyway, they are true.

## A last word of support

We sum up the various sources of supporting evidence for the conjectures :

- Computation of the  $A_\pi(X)$  for small  $|\pi|$  ;
- Compatibility of the conjectures with known facts ;
- Coherence of the conjectures among themselves.
- Proof of the conjectures for special values ;
- Proof of the conjectures for certain families of matchings  $\pi$ , for which  $A_\pi(X)$  is known explicitly [Di-Francesco and al '04, Caselli and Krattenthaler '05].

I wish this were enough to turn them into theorems ; it's not...  
But anyway, they are true.

I guess.

# Conclusion

- The conjectures lead us to believe that there is *combinatorial reciprocity* result underlying the  $A_\pi(X)$ , à la Ehrhart polynomial :  
 $\Rightarrow$  there “should be” nice objects enumerated by the values  $A_\pi(-i)$ .

Especially interesting is to conjecture/prove :

What do the numbers  $G_\pi$  count ?


# Conclusion

- The conjectures lead us to believe that there is *combinatorial reciprocity* result underlying the  $A_\pi(X)$ , à la Ehrhart polynomial :  
 $\Rightarrow$  there “should be” nice objects enumerated by the values  $A_\pi(-i)$ .

Especially interesting is to conjecture/prove :

What do the numbers  $G_\pi$  count ?

- **The  $\tau$  case** : There exists a refinement of the probabilities  $\psi_\pi$  to polynomials in  $\tau$ , with no known equivalent for the  $A_\pi$ , which specialize to our previous setting for  $\tau = 1$ .  
Our conjectures all have “ $\tau$  versions” dealing with bivariate polynomials  $\psi_\pi(X, \tau)$ .



$$A_{\pi_0}(X) = \frac{(2+X)(3+X)^2(4+X)^2(5+X)^2(6+X)(7+X)}{145152000} \\ \times (9X^6 + 284X^5 + 4355X^4 + 39660X^3 + 225436X^2 \\ + 757456X + 123120)$$