

Some Surprising Results on a One-Dimensional Elliptic Boundary Value Blow-Up Problem

Y. J. Cheng

Abstract. In this paper we consider the one-dimensional elliptic boundary blow-up problem

$$\left. \begin{aligned} \Delta_p u &= f(u) \quad (a < t < b) \\ u(a) &= u(b) = +\infty \end{aligned} \right\}$$

where $\Delta_p u = (|u'(t)|^{p-2}u'(t))'$ is the usual p -Laplace operator. We show that the structure of the solutions can be very rich even for a simple function f which gives a leading that a similar results might hold also in higher dimensional spaces

Keywords: *Boundary blow-up, multiplicity, concave and convex nonlinearity*

AMS subject classification: 34B, 35J

1. Introduction and formulation of main results

In the last few years there is a great of interests in the investigation of boundary blow-up solutions for elliptic equations [2, 3, 7], which comes originally from differential geometry [6] and electrohydrodynamics [5]. Very recently some existence results of two (one positive and one sign-changing) solutions have been established in [1, 8]. The purpose of the present paper is to show through one-dimensional examples that the structure of the solutions can be very rich even for a simple right-hand side. More precisely, we consider the problem

$$\left. \begin{aligned} \Delta_p u &= \lambda f(u) \quad (a < t < b) \\ u(a) &= u(b) = +\infty \end{aligned} \right\} \quad (1)$$

where

$$\Delta_p u = (|u'(t)|^{p-2}u'(t))'$$

is the p -Laplace operator as usual, $\lambda > 0$ is a parameter, and f is a given continuous function. By a solution $u = u(t)$ of problem (1) we mean that u satisfies the equation in (1), i.e. $(|u'(t)|^{p-2}u'(t))' = \lambda f(u(t))$ for all $t \in (a, b)$, and $\lim_{t \rightarrow a+} u(t) = \lim_{t \rightarrow b-} u(t) = +\infty$. By a sign-changing solution $u(t)$ of (1) we mean that there exist $t_1, t_2 \in (a, b)$ such that $u(t_1) > 0$ and $u(t_2) < 0$.

The results for problem (1) in this paper are summarized in the followings three theorems.

Yuanji Cheng: Univ. of Malmö, School of Techn. and Management., 205 06 Malmö, Sweden

Theorem 1. For given positive constants q, ε, s and α, r such that $q > p-1 > r > s$ let

$$f(u) = u^q + \varepsilon u^s \quad (u \geq 0) \quad \text{and} \quad f(u) = \alpha |u|^r \quad (u \leq 0).$$

Then the following statements hold:

(i) There exists a constant $\lambda_1 > 0$ such that problem (1) has at least one sign-changing solution if $\lambda > \lambda_1$, and no such kind of solutions can exist if $\lambda < \lambda_1$.

(ii) There exist a constant $\varepsilon_0 > 0$ such that the solution is unique if $\varepsilon \geq \varepsilon_0$.

(iii) If $\varepsilon \in (0, \varepsilon_0)$, then there are constants $\lambda_2 < \lambda_3 \leq \lambda_4$ such that problem (1) has at least two sign-changing solutions if $\lambda \in (\lambda_1, \lambda_4)$, has at least three sign-changing solutions if $\lambda \in (\lambda_2, \lambda_3)$, and has a unique sign-changing solution if $\lambda > \lambda_4$.

Figure 1

Theorem 2. For given positive constants q, ε, s and α, r, δ, τ such that $q, \tau > p-1 > r > s$ let

$$f(u) = u^q + \varepsilon u^s \quad (u \geq 0) \quad \text{and} \quad f(u) = \alpha |u|^r + \delta |u|^\tau \quad (u \leq 0).$$

Then:

(i) There are constants $\Lambda > 0$ and $\Lambda_- > 0$ such that problem (1) has at least one sign-changing solution if $\lambda \leq \Lambda$, has no sign-changing solutions if $\lambda > \Lambda$, and the solution is unique if $\lambda < \Lambda_-$.

Figure 2

(ii) For small ε and δ satisfying $\varepsilon \ll \delta$ there are $\lambda_1 < \lambda_2$ such that if $\lambda \in (\lambda_1, \lambda_2)$, then problem (1) has at least four sign-changing solutions.

Next we consider for simplicity the semilinear ($p = 2$) problem

$$\left. \begin{aligned} u'' &= \lambda f(u) \quad (a < t < b) \\ u(a) &= u(b) = +\infty. \end{aligned} \right\} \tag{2}$$

For this simple problem we have the following, a somehow surprising result.

Theorem 3. For given positive constants α, q and β such that $q > 3$ and $\beta < 1$ let

$$f(u) = \alpha u^q \quad (u \geq 0) \quad \text{and} \quad f(u) = (1 + \beta \sin |u|)|u| \quad (u \leq 0)$$

$(\lambda(2) = (\frac{\pi}{b-a})^2)$. Then the following results hold:

(i) There are $\lambda_1, \lambda_2 > 0$ such that problem (2) has at least one sign-changing solution if $\lambda \geq \lambda_1$ and no sign-changing solutions if $\lambda < \lambda_1$. The sign-changing solution is unique, when $\lambda > \lambda_2$.

(ii) For any integer $n \geq 1$ there exists $\delta > 0$ such that problem (2) has at least n distinct sign-changing solutions when $\lambda \in (\lambda(2) - \delta, \lambda(2) + \delta)$.

(iii) For $\lambda = \lambda(2)$ problem (2) has infinitely many sign-changing solutions.

Figure 3

Remark. Problem (2) has a unique positive solution for all $\lambda > 0$. The problems treated in Theorems 1 and 2 have also a unique positive solution if $\lambda < \lambda_0$ where λ_0 is defined by

$$\int_0^\infty \frac{du}{\sqrt{\frac{1}{q+1} u^{1+q} + \frac{\varepsilon}{s+1} u^{1+s}}} = \frac{b-a}{2} \sqrt[p]{p' \lambda_0}$$

and has no positive solution if $\lambda \geq \lambda_0$.

2. Some basic analysis

It is easy to see that the equation in (1) has an first integral

$$\frac{1}{p'}|u'(t)|^p = \lambda F(u) + C \quad \text{with } F(u) = \int_0^u f(x) dx \quad (3)$$

where $\frac{1}{p} + \frac{1}{p'} = 1$. Let $t_0 \in (a, b)$ be a minimum point of $u(t)$, which exists by the boundary condition. Then $u'(t_0) = 0$ and $C = -\lambda F(u_0)$, $u_0 = \min u(t)$, and

$$|u'(t)|^p = \lambda p'(F(u) - F(u_0)) \quad (a < t < b).$$

If $f = f(u)$ is non-negative, then we see that $u = u(t)$ is convex and the minimum point $t = t_0$ is unique. Consequently, $u'(t) \leq 0$ for $t \in (a, t_0)$ and $u'(t) \geq 0$ for $t \in (t_0, b)$. Moreover,

$$u'(t) = \sqrt[p']{\lambda p'(F(u) - F(u_0))} \operatorname{sign}(t - t_0) \quad (a < t < b).$$

Direct integration yields

$$(b - t_0) \sqrt[p']{p'\lambda} = \int_{u_0}^{+\infty} \frac{du}{\sqrt[p']{F(u) - F(u_0)}} = (t_0 - a) \sqrt[p']{p'\lambda}$$

which implies $t_0 = \frac{a+b}{2}$ and thus $u = u(t)$ must be symmetric.

To establish the existence and the structure of solutions of problem (1) it suffices to study the nonlinear integral equation

$$\int_{u_0}^{+\infty} \frac{du}{\sqrt[p']{F(u) - F(u_0)}} = \frac{b - a}{2} \sqrt[p']{p'\lambda}. \quad (4)$$

Obviously, a necessary condition for the existence of solutions for problem (1) is

$$\int^{+\infty} \frac{du}{\sqrt[p']{F(u)}} < +\infty \quad (5)$$

and so throughout this paper we shall assume that this condition holds.

Rewriting the integral in (4) gives that it is equivalent to

$$\int_0^{+\infty} \frac{du}{\sqrt[p']{F(u + u_0) - F(u_0)}} = \frac{b - a}{2} \sqrt[p']{p'\lambda}. \quad (6)$$

It follows from here that problem (1) has at most one (positive) solution under the condition that $f(u)$ is non-decreasing on \mathbb{R} (or on \mathbb{R}_+). On the other hand, if $f(u) > 0$ for $u > 0$, then it has at least one positive solution for all $\lambda < \lambda_0$, where $\lambda_0 \in (0, +\infty]$ is defined by

$$\int_0^{+\infty} \frac{du}{\sqrt[p']{F(u)}} = \frac{b - a}{2} \sqrt[p']{p'\lambda_0}. \quad (7)$$

By summarizing, let λ_0 be as above. Then we have the following

Theorem 4. *If $f(u) > 0$ ($u > 0$), then problem (1) has at least one positive solution for all $\lambda < \lambda_0$. Moreover, if f is also non-decreasing on \mathbb{R}_+ , then the solution is unique.*

Example. Consider the problem

$$\left. \begin{aligned} \Delta_p u &= \lambda(u^q + \varepsilon u^s) \quad (a < t < b) \\ u(a) &= u(b) = +\infty \end{aligned} \right\} \tag{8}$$

where $0 < r < q$, $\varepsilon > 0$ and $q > p - 1$. Then condition (5) is satisfied, and further $\lambda_0 = +\infty$ if $s \geq p - 1$ and $\lambda_0 < +\infty$ if $s \in (0, p - 1)$. Hence problem (8) has a unique positive solution for all $\lambda > 0$ if $q > r \geq p - 1$, and for all $\lambda < \lambda_0$ if $0 < s < p - 1$, where λ_0 satisfies

$$\int_0^{+\infty} \frac{du}{\sqrt[p]{\frac{1}{q+1} u^{1+q} + \frac{\varepsilon}{s+1} u^{1+s}}} = \frac{b-a}{2} \sqrt[p']{p'\lambda_0}.$$

3. Proofs

To investigate solutions of problem (1) which change its sign, we define

$$f_+(u) = f(u) \quad (u \geq 0) \quad \text{and} \quad f_-(u) = f(-u) \quad (u \leq 0).$$

Then we have

$$F(u) = \begin{cases} F_+(u) = \int_0^u f_+(x) dx & \text{for } u \geq 0 \\ -F_-(-u) = -\int_0^{-u} f_-(x) dx & \text{for } u < 0. \end{cases}$$

Now equation (6) becomes

$$\int_0^{+\infty} \frac{du}{\sqrt[p]{F_+(u) + F_-(v_0)}} + \int_0^{v_0} \frac{du}{\sqrt[p]{F_-(v_0) - F_-(u)}} = \frac{b-a}{2} \sqrt[p']{p'\lambda} \tag{9}$$

where $v_0 = -u_0 > 0$. First observe that the first integral in (9) is strictly decreasing in v_0 if $f_-(u)$ is non-negative. For the second integral in (9), using

$$\int_0^{v_0} \frac{du}{\sqrt[p]{F_-(v_0) - F_-(u)}} = \int_0^1 \frac{v_0 ds}{\sqrt[p]{F_-(v_0) - F_-(sv_0)}}$$

and

$$F_-(v_0) - F_-(sv_0) = \int_0^1 f_-((s+t-st)v_0)(1-s)v_0 dt,$$

we see that it is decreasing or increasing in v_0 if $f_-(u)u^{1-p}$ is increasing or decreasing, respectively. Therefore, if $f_-(u)$ is non-negative and $f_-(u)u^{1-p}$ is increasing, then (9) has at most one solution, which gives the uniqueness of sign-changing solutions of problem (1). On the other hand, if $f_-(u)u^{1-p}$ is decreasing, then the first and the second integrals in (9) will compete to each other and thus the existence of a multiple solution is possible. In particular, if

$$\int_0^\infty \frac{du}{\sqrt[p]{F_+(u)}} = +\infty \quad \text{and} \quad \lim_{u \rightarrow +\infty} f_-(u)u^{1-p} = 0,$$

then the left side in (9) goes to infinity, when v_0 goes either to zero or to infinity. Hence there is $\lambda_- > 0$ such that problem (1) has at least two sign-changing solutions for $\lambda > \lambda_-$, and no sign-changing solution if $\lambda < \lambda_-$. A typical example is

$$f_+(u) = \sum \alpha_i u^{q_i} \quad \text{and} \quad f_-(u) = \sum \beta_j u^{r_j}$$

where $\alpha_i, \beta_j > 0$ and $q_i > p - 1 > r_j$. When f_+ and f_- are simply given by αu^q and βu^r ($q > p - 1 > r$), respectively, then we have a complete characterization of signchanging solutions, namely, two solutions if $\lambda > \lambda_-$, a unique solution if $\lambda = \lambda_-$, and no solutions if $\lambda < \lambda_-$. In other situations the multiplicity of sign-changing solutions can be very complicated and three representative examples are as in Theorems 1 - 3.

Proof of Theorem 1. To this end we study the function in the left-hand side of (9). In this case we have

$$F_+(u) = \frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} \quad \text{and} \quad F_-(u) = \frac{\alpha}{r+1} u^{r+1}$$

and

$$\int_0^1 \frac{v_0 ds}{\sqrt[p]{F_-(v_0) - F_-(sv_0)}} = c_2 v_0^{1-\frac{1+r}{p}} \quad \text{with} \quad c_2 = \int_0^1 \frac{ds}{\sqrt[p]{\alpha(1-s^{r+1})/(1+r)}}$$

and the function in (9) is

$$\int_0^\infty \frac{du}{\sqrt[p]{\frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} + \frac{\alpha}{1+r} v_0^{r+1}}} + c_2 v_0^{1-\frac{r+1}{p}} =: F_\varepsilon(v_0). \tag{10}$$

The first conclusion of Theorem 1 follows from that $F_\varepsilon(v_0) > 0$ is continuous on $[0, +\infty)$ and goes to infinity, as $v_0 \rightarrow \infty$. To get a complete picture for the existence of sign-changing solutions we let

$$k_1 = (1+r)\left(\frac{1}{s+1} - \frac{1}{p}\right), \quad k_2 = \frac{(1+r)(q-s)}{1+s}, \quad k_3 = 1 - \frac{r+1}{p}$$

and

$$g(v_0) = \int_0^{+\infty} \left(\frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} + \frac{\alpha}{1+r} v_0^{r+1}\right)^{-\frac{1}{p}} du.$$

Then $F_\varepsilon(v_0) = g(v_0) + c_2 v_0^{k_3}$. We first study the property of $g(v_0)$ in a neighbourhood of the origin.

By the change of variable $u = x v_0^{\frac{1+r}{1+s}}$ we get for $v_0 > 0$

$$g(v_0) = \int_0^{+\infty} \frac{v_0^{k_1} dx}{\sqrt[p]{\frac{1}{q+1} x^{q+1} v_0^{k_2} + \frac{\varepsilon}{1+s} x^{1+s} + \frac{\alpha}{1+r}}}$$

$$g(0) = \int_0^{+\infty} \frac{v_0^{k_1} dx}{\sqrt[p]{\frac{1}{q+1} x^{q+1} v_0^{k_2} + \frac{\varepsilon}{1+s} x^{1+s}}}$$

Consequently,

$$\frac{g(v_0) - g(0)}{v_0^{k_1}} = - \int_0^{+\infty} G(v_0, x) dx$$

where

$$G(v_0, x) = \frac{1}{\sqrt[p]{\frac{1}{q+1} x^{q+1} v_0^{k_2} + \frac{\varepsilon}{1+s} x^{1+s}}} - \frac{1}{\sqrt[p]{\frac{1}{q+1} x^{q+1} v_0^{k_2} + \frac{\varepsilon}{1+s} x^{1+s} + \frac{\alpha}{1+r}}}$$

Writing the difference as an integral we see

$$G(v_0, x) = \frac{\alpha}{p(r+1)} \int_0^1 \left(\frac{1}{q+1} x^{q+1} v_0^{k_2} + \frac{\varepsilon}{1+s} x^{1+s} + \frac{\alpha}{1+r} \theta \right)^{-\frac{p+1}{p}} d\theta$$

which implies that $G(v_0, x)$ is decreasing in v_0 . Since

$$G(0, x) = \frac{\alpha}{p(r+1)} \int_0^1 \left(\frac{\varepsilon}{1+s} x^{1+s} + \frac{\alpha}{1+r} \theta \right)^{-\frac{p+1}{p}} d\theta$$

is integrable over $(0, +\infty)$, due to $s \in (0, p-1)$, we deduce by the dominate convergence theorem that

$$\lim_{v_0 \rightarrow 0} \frac{g(v_0) - g(0)}{v_0^{k_1}} = - \int_0^{+\infty} G(0, x) dx < 0.$$

Thus we obtain that near the origin F_ε is increasing if $k_1 > 1 - \frac{r+1}{p}$, which is equivalent to $r > s$, and is decreasing when $r < s$. Similarly, we get that

$$\lim_{v_0 \rightarrow \infty} g'(v_0) v_0^{1+(r+1)(\frac{1}{p} - \frac{1}{q+1})} = -\frac{\alpha}{p} \int_0^1 \left(\frac{1}{q+1} x^{q+1} + \frac{\alpha}{1+r} \right)^{-\frac{p+1}{p}} dx$$

which implies that F_ε is increasing for large $v_0 > 0$, because $q > p - 1$. Since $\frac{dF_\varepsilon}{dv_0}$ is increasing in ε and tends to $(1 - \frac{r+1}{p})c_2 v_0^{-\frac{r+1}{p}}$ uniformly in v_0 as $\varepsilon \rightarrow \infty$, we deduce in the case $r > s$ that there is an $\varepsilon_0 \geq 0$ such that $\min_{v_0} \frac{dF_\varepsilon}{dv_0} > 0$ for $\varepsilon > \varepsilon_0$. Moreover, when $\varepsilon = 0$,

$$F_0(v_0) = c_1 v_0^{1-\frac{q+1}{p}} + c_2 v_0^{1-\frac{r+1}{p}}$$

for $1 - \frac{q+1}{p} < 0 < 1 - \frac{r+1}{p}$ and thus $\frac{dF_0}{dv_0} < 0$ for small $v_0 > 0$. Further, $F_\varepsilon(v_0) \rightarrow F_0(v_0)$ on compacta in $C^1(0, +\infty)$ as $\varepsilon \rightarrow 0$ and we assert that $\varepsilon_0 > 0$ and $\min_{v_0} \frac{dF_\varepsilon}{dv_0} < 0$ for $\varepsilon \in (0, \varepsilon_0)$. Whence, we obtain that the sign-changing solution is unique when $\varepsilon \geq \varepsilon_0$. If $\varepsilon \in (0, \varepsilon_0)$, using the fact that F_ε is increasing for both small and large v_0 , we deduce that there are $0 < v_1 < v_2 \leq v_3 < +\infty$ such that F_ε is increasing on $(0, v_1)$ and $(v_3, +\infty)$ and is decreasing on (v_1, v_2) . Thus let

$$\lambda_1 = \min F_\varepsilon(v_0), \quad \lambda_2 = F_\varepsilon(v_1), \quad \lambda_3 = F_\varepsilon(v_2), \quad \lambda_4 = \max_{F'_\varepsilon(v_0)=0} F_\varepsilon(v_0).$$

Then problem (1) has no sign-changing solutions if $\lambda < \lambda_1$ and has at least two sign-changing solutions $\lambda \in (\lambda_1, \lambda_4)$. In particular, it has at least three sign-changing solutions if $\lambda \in (\lambda_2, \lambda_3)$ and has a unique sign-changing solution as $\lambda > \lambda_4$. The proof is complete ■

Proof of Theorem 2. The idea is the same as in the proof of Theorem 1, so we will be sketch in many places. In view of

$$F_+(u) = \frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} \quad \text{and} \quad F_-(u) = \frac{\alpha}{r+1} u^{r+1} + \frac{\delta}{\tau+1} u^{\tau+1}$$

it follows that equation (9) takes the form

$$\begin{aligned} & \int_0^{+\infty} \frac{du}{\sqrt[p]{\frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} + F_-(v_0)}} \\ & + \int_0^1 \frac{v_0 dx}{\sqrt[p]{r(x) v_0^{s+1} + \tau(x) v_0^{\tau+1}}} = \frac{b-a}{2} \sqrt[p]{\alpha p' \lambda} \end{aligned} \tag{11}$$

where

$$r(x) = \frac{\alpha}{r+1} (1 - x^{r+1}) \quad \text{and} \quad \tau(x) = \frac{\delta}{\tau+1} (1 - x^{\tau+1}).$$

Likely as in the proof of Theorem 1 we define

$$L(v_0) = \ell_1(v_0) + \ell_2(v_0)$$

where

$$\begin{aligned} \ell_1(v_0) &= \int_0^{+\infty} \frac{du}{\sqrt[p]{\frac{1}{q+1} u^{q+1} + \frac{\varepsilon}{1+s} u^{1+s} + F_-(v_0)}} \\ \ell_2(v_0) &= \int_0^1 \frac{v_0 dx}{\sqrt[p]{r(x) v_0^{s+1} + \tau(x) v_0^{\tau+1}}}. \end{aligned}$$

The key points to prove that the function $L(v_0)$ have a graph as shown in Figure 2 are

$$L(v_0) = F_0(v_0) = c_1 v_0^{\frac{p-q-1}{p}} + c_2 v_0^{\frac{p-r-1}{p}} \quad \text{when } \varepsilon = \delta = 0$$

which has the property $F'_0(v_0)(v_0 - v) > 0$ for some $v > 0$ and L converges to F_0 on compacta of $(0, +\infty)$ in C^1 as $\varepsilon, \delta \rightarrow 0$ and $L \rightarrow 0$ as $v_0 \rightarrow +\infty$.

To complete the proof we need to show two more things: Monotonicity of L near the origin and $L(0) \leq L(v_2)$ where v_2 is the second local maximum of L . But

$$L(0) = c_1 \varepsilon^{\frac{q-p-1}{p(q-s)}} \quad \text{and} \quad L(\delta^{\frac{1}{r-\tau}}) \geq \ell_2(\delta^{\frac{1}{r-\tau}}) = c_2 \delta^{\frac{r+1-p}{p(\tau-r)}}$$

for some constants $c_1 > 0$ and $c_2 > 0$ from which it follows that if ε and δ satisfy $c_1 \varepsilon^{\frac{q-p-1}{p(q-s)}} \leq c_2 \delta^{\frac{r+1-p}{p(\tau-r)}}$, then we are done.

To get the monotonicity of L in the nearby of the origin, we exploit the technique in the proof of Theorem 1 and can show that

$$\lim_{v_0 \rightarrow 0} \frac{\ell_1(v_0) - \ell_1(0)}{v_0^{k_1}} = -\frac{\alpha}{p(r+1)} \int_0^{+\infty} dx \int_0^1 \left(\frac{\varepsilon}{1+s} x^{s+1} + \frac{\alpha}{r+1} \theta \right)^{-\frac{p+1}{p}} d\theta < 0$$

where we have used that $r > \tau$ and $F_-(v_0) = \frac{\alpha}{r+1} v_0^{r+1} (1 + o(1))$ as $v_0 \rightarrow 0$,

$$\lim_{v_0 \rightarrow 0} \frac{\ell_2(v_0) - \ell_2(0)}{v_0^{k_3}} = \int_0^1 \frac{dx}{\sqrt[r]{r(x)}} > 0 \quad \text{and} \quad \lim_{v_0 \rightarrow 0} \frac{L(v_0) - L(0)}{v_0^{k_1}} < 0.$$

Finally, the uniqueness follows from the results in [4] that the function ℓ_2 has the following property: there is $v > 0$ such that $\ell'_2(v_0) > 0$ for $v_0 \in (0, v)$ and $\ell'_2(v_0) < 0$ for $v_0 \in (v, +\infty)$. Then $L'(v_0) < 0$ for $v_0 \in [v, +\infty)$ since clearly $\ell'_1(v_0) < 0$ for $v_0 > 0$. The proof is complete \blacksquare

Proof of Theorem 3. By virtue of $p = 2$ and

$$F_+(u) = \frac{\alpha}{q+1} u^{q+1} \quad \text{and} \quad F_-(u) = \frac{1}{2} u^2 + \beta(\sin u - u \cos u)$$

we have that

$$\int_0^{+\infty} \frac{du}{\sqrt{F_+(u) + F_-(v_0)}} = c(q) \left(\frac{1}{2} v_0^2 + \beta(\sin v_0 - v_0 \cos v_0) \right)^{\frac{1}{q+1} - \frac{1}{2}} =: g_1(v_0)$$

$$\int_0^{v_0} \frac{du}{\sqrt{F_-(v_0) - F_-(u)}} = \int_0^1 \frac{v_0 dx}{\sqrt{h(x, v_0)}} =: g_2(v_0)$$

where

$$\begin{aligned}
 h(x, v_0) &= e(x) v_0^2 + \beta \varepsilon(x, v_0) \\
 \varepsilon(x, v_0) &= \sin v_0 - \sin(xv_0) - v_0 (\cos v_0 - x \cos(xv_0)) \\
 c(q) &= \int_0^{+\infty} \left(1 + \frac{\alpha}{q+1} u^{q+1}\right)^{-\frac{1}{2}} du \\
 e(x) &= \frac{1-x^2}{2}
 \end{aligned}$$

and the function in the left-hand side of (9) takes the form

$$g_1(v_0) + g_2(v_0) := H(v_0).$$

Clearly, H is well defined and continuous on $(0, +\infty)$. Since $q > 3$, we see that

$$g_1(v_0) \rightarrow +\infty \quad \text{as } v_0 \rightarrow 0+$$

and

$$\left. \begin{aligned}
 g_1(v_0) &\rightarrow 0 \\
 g_2(v_0) &\rightarrow \int_0^1 \frac{dx}{\sqrt{e(x)}} = \frac{\pi}{\sqrt{2}}
 \end{aligned} \right\} \quad \text{as } v_0 \rightarrow +\infty$$

and therefore $H(0+) = +\infty$ and $H(+\infty) = \frac{\pi}{\sqrt{2}}$.

To show the results (ii) and (iii) we need to prove that H oscillates around the line $H = \frac{\pi}{\sqrt{2}}$, and first show that g_1 oscillates around $H = \frac{\pi}{\sqrt{2}}$.

Choosing $v_0 = n\pi$ ($n \in \mathbb{N}$ is even) we get

$$\begin{aligned}
 h(x, n\pi) &= (n\pi)^2 e(x) - \beta \left(\sin(n\pi x) + n\pi(1 - x \cos(n\pi x)) \right) \\
 &\leq (n\pi)^2 e(x) - \beta \left(n\pi(1 - x) - \sin(n\pi(1 - x)) \right) \\
 &\leq (n\pi)^2 e(x)
 \end{aligned}$$

thereafter

$$g_2(n\pi) > \int_0^1 \frac{n\pi dx}{\sqrt{e(x)(n\pi)^2}} = \frac{\pi}{\sqrt{2}}.$$

Analogously, for odd integer n , $g_2(n\pi) < \frac{\pi}{\sqrt{2}}$.

To show the function H is also oscillating, it suffices to show that, for odd n , $H(n\pi) < \frac{\pi}{\sqrt{2}}$ since $g_1 > 0$ and $g_2(n\pi) > \frac{\pi}{\sqrt{2}}$ for even n . For this purpose, first we have

$$\begin{aligned}
 \frac{\pi}{\sqrt{2}} - g_2(n\pi) &= \int_0^1 \left(\frac{n\pi}{\sqrt{e(x)(n\pi)^2}} - \frac{n\pi}{\sqrt{e(x)(n\pi)^2 + \beta \varepsilon(x, n\pi)}} \right) dx \\
 &= \frac{1}{2} \int_0^1 dx \int_0^1 \frac{n\pi \beta \varepsilon(x, n\pi)}{(e(x)(n\pi)^2 + \theta \beta \varepsilon(x, n\pi))^{\frac{3}{2}}} d\theta \\
 &\geq \frac{1}{2} \int_0^1 \frac{n\pi \beta \varepsilon(x, n\pi)}{(e(x)(n\pi)^2 + \beta \varepsilon(x, n\pi))^{\frac{3}{2}}} dx,
 \end{aligned}$$

due to $\varepsilon(x, n\pi) > 0$ for all $x \in (0, 1)$. Using $f_-(u) \geq (1 - \beta)u$, we deduce that

$$\begin{aligned} e(x)(n\pi)^2 + \beta \varepsilon(x, n\pi) &= n\pi(1 - x) \int_0^1 f_-((x + t - tx)n\pi) dt \\ &\geq n\pi(1 - x) \int_0^1 (1 - \beta)(x + t - tx)n\pi dt \\ &\geq \frac{1}{2}(1 - \beta)(1 - x)(n\pi)^2 \end{aligned}$$

for all $x \in (0, 1)$. Thereafter

$$\frac{\pi}{\sqrt{2}} - g_2(n\pi) \geq \frac{2^{\frac{3}{2}}\beta}{(1 - \beta)^{\frac{3}{2}}}(n\pi)^{-2} \int_0^1 \frac{\varepsilon(x, n\pi)}{(1 - x)^{\frac{3}{2}}} dx,$$

and by a change of variable $z = \frac{1}{\sqrt{1-x}}$ (then $x = 1 - \frac{1}{z^2}$)

$$\begin{aligned} \int_0^1 \frac{\varepsilon(x, n\pi)}{(1 - x)^{\frac{3}{2}}} dx &= 2 \int_1^{+\infty} \varepsilon\left(1 - \frac{1}{z^2}, n\pi\right) dz \\ &= 2 \int_1^{+\infty} \left(-\sin\left(\frac{n\pi}{z^2}\right) + n\pi\left(1 - \left(1 - \frac{1}{z^2}\right)\cos\frac{n\pi}{z^2}\right)\right) dz \quad cr \\ &= 2 \int_1^{+\infty} \left(n\pi\left(1 - \cos\frac{n\pi}{z^2}\right) + \left(-\sin\frac{n\pi}{z^2} + \frac{n\pi}{z^2}\cos\frac{n\pi}{z^2}\right)\right) dz \end{aligned}$$

where we have used n being an odd integer. Change variables once more, we obtain

$$\int_1^{+\infty} \left(1 - \cos\frac{n\pi}{z^2}\right) dz = \int_{\frac{1}{\sqrt{n\pi}}}^{+\infty} \left(1 - \cos\frac{1}{w^2}\right) \sqrt{n\pi} dw \geq C\sqrt{n\pi}$$

where $C = \int_1^{+\infty} (1 - \cos \frac{1}{z^2}) dz$. On the other hand,

$$\int_1^{+\infty} \left(-\sin\frac{n\pi}{z^2} + \frac{n\pi}{z^2}\cos\frac{n\pi}{z^2}\right) dz \leq \int_1^{+\infty} \frac{2n\pi}{z^2} dz = 2n\pi.$$

Whence

$$\int_0^1 \frac{\varepsilon(x, n\pi)}{(1 - x)^{\frac{3}{2}}} dx \geq 2(Cn\pi\sqrt{n\pi} - 2n\pi)$$

and consequently for large n the estimate

$$\frac{\pi}{\sqrt{2}} - g_2(n\pi) \geq \frac{2^{\frac{3}{2}}\beta}{(1-\beta)^{\frac{3}{2}}} C(n\pi)^{-\frac{1}{2}}$$

holds. Combing the above estimates, we obtain by noting $g_1(n\pi) \leq C_1(n\pi)^{\frac{2}{q+1}-1}$ that

$$\frac{\pi}{\sqrt{2}} - H(n\pi) = \frac{\pi}{\sqrt{2}} - g_2(n\pi) - g_1(n\pi) \geq C_2(n\pi)^{-\frac{1}{2}} - C_1(n\pi)^{\frac{2}{q+1}-1} > 0$$

since $-\frac{1}{2} > \frac{2}{q+1} - 1$ (this is the source for the condition on q).

Assertions (ii) and (iii) follow from the oscillatory property of H . Using

$$\lim_{v_0 \rightarrow \infty} H(v_0) = \frac{\pi}{\sqrt{2}}, \quad \lim_{v_0 \rightarrow 0^+} H(v_0) = +\infty, \quad H(n\pi) < \frac{\pi}{\sqrt{2}}$$

we deduce from the continuity of H that the minimum of $H(v_0)$ on $(0, +\infty)$ achieves and thus the equation $H(v_0) = \mu$ is solvable if and only if $\mu \geq \min H(v_0)$ which complete the proof of assertion (i) ■

4. Final remarks

In this note we have only carried out some basic calculations to exhibit the rich structure for boundary blow-up problems, even it is very elementary (just calculus), but it is certainly not easy to give a complete bifurcation picture for all involved parameters, for instance $q, r, s, \tau, \varepsilon, \delta, \alpha$ in Theorem 2. From our one-dimensional examples we can see that there is a big difference between Dirichlet boundary value problems and boundary blow-up problems. If the boundary is Dirichlet, then there are infinitely many sign solutions in the superlinear case, but it can have only finite number of sign solutions for boundary blow-up problems (note, the function in (2) is not superlinear at $-\infty$). As our examples are of one-dimensional character, one may say that it would not be representative, so it will be interesting to study those equations in two-dimensional or higher dimensional domains.

References

- [1] Aftalion, A. and W. Reichel: *Existence of two boundary blow-up solutions for semilinear elliptic equations*. J. Diff. Equ. 141 (1997), 400 – 421.
- [2] Bandle, C. and M. Essén: *On the solutions of quasilinear elliptic problems with boundary blow-up*. In: Partial Differential Equations of Elliptic Type (Symposia Mathematica: Vol. 35; eds.: A. Alvino et al.). Univ. Press, Cambridge (1994), pp. 93 – 111.
- [3] Bandle, C. and M. Marcus: *Asymptotic behaviour of solutions and their derivatives, for semilinear elliptic problems with boundary blow-up*. Ann. Inst. Henri Poincaré 12 (1995), 155 – 171.

- [4] Cheng, Y. J.: *On an open problem of Ambrosetti, Brezis and Cerami*. Diff. Int. Equ. (to appear).
- [5] Keller, J. B.: *Electrohydrodynamics. Part I: The equilibrium of a charged gas in a container*. J. Rat. Mech. Anal. 5 (1956), 715 – 724.
- [6] Loewner, C. and L. Nirenberg: *Partial differential equations invariant under conformal or projective transformations*. In: Contribution to Analysis (ed.: L. Ahlfors). Acad. Press, New York (1974), pp. 245 – 272.
- [7] Matero, J.: *Nonlinear Elliptic Problems with Boundary Blow-Up*. Diss. in Math. 6. Uppsala: Univ. 1997.
- [8] Mckenna, P. J., Reichel, W. and W. Walter: *Symmetry and multiplicity for nonlinear elliptic differential equations with boundary blow-up*. Nonlin. Anal. Theory, Methods and Appl. 28 (1997), 1213 – 1225.

Received 26.10.1998; in revised form 21.04.1999