

LAMB SHIFT IN CLASSICAL ELECTRODYNAMICAL MODEL OF ATOM

V. SIMULIK and I. KRIVSKY

*Institute of Electron Physics, Ukrainian National Academy of Sciences
21 Universitetska Street, 88000 Uzhgorod, Ukraine*

Abstract. The Lamb shift in hydrogen spectrum is obtained here in the framework of some weak generalization of the classical Maxwell electrodynamics. It is shown that weakly generalized classical Maxwell electrodynamics can describe the intra-atomic phenomena with the same success as relativistic quantum mechanics can do. Group-theoretical grounds for the description of fermionic states by bosonic system are presented briefly. The advantages of generalized electrodynamics in intra-atomic region in comparison with standard Maxwell electrodynamics are demonstrated on testing example of hydrogen atom. We are able to obtain some results which are impossible in the framework of standard Maxwell electrodynamics. The Sommerfeld–Dirac formula for the fine structure of the hydrogen atom spectrum is obtained on the basis of such Maxwell equations without appealing to the Dirac equation. The Bohr postulates are proved to be the consequences of the equations under consideration. The relationship of the new model with the Dirac theory is investigated. Possible directions of unification of such electrodynamics with gravity are mentioned.

1. Introduction

Lamb shift is a specific effect of quantum electrodynamical consideration. It is explained in this theory as a consequence of polarization of vacuum. Below we are able to explain this effect in the framework of classical electrodynamics. More precisely speaking this effect is obtained in the framework of weakly generalized classical electrodynamics, which contains electric and magnetic gradient-type sources.

Before such nonstandard consideration of the Lamb shift the group-theoretical grounds of a new model of atom and the main assertions of this model must

be presented. Let us consider firstly the reasons why the classical Maxwell electrodynamics needs some weak generalization and after that the foundations of the new model of atom, which follows from such generalized Maxwell equations.

There is no doubt that the classical Maxwell electrodynamics of macroworld (without any generalization) is sufficient for the description of electrodynamic phenomena in macro region. On the other hand it is well known that for micro phenomena (inneratomic region) the classical Maxwell electrodynamics (as well as the classical mechanics) cannot work and must be replaced by quantum theory. Trying to extend the limits of classical electrodynamics application to the intra-atomic region we came to the conclusion that it is possible by means of generalization of the standard Maxwell electrodynamics in the direction of extension of its symmetry. We also use the relationships between the Dirac and Maxwell equations for these purposes. Furthermore, the relationships between relativistic quantum mechanics and classical microscopical electrodynamics of media are investigated. Such relationships are considered here not only from the mathematical point of view — they are used for construction of fundamentals of a nonquantum-mechanical model of microworld.

Our nonquantum-mechanical model of microworld is a model of atom on the basis of weakly generalized Maxwell's equations, i. e. in the framework of weakly extended classical microscopical electrodynamics of media. This model is free from probability interpretation and can explain many intra-atomic phenomena by means of classical physics. Despite the fact that we construct the classical model, for the purposes of such construction we use essentially the analogy with the Dirac equation and the results which were achieved on the basis of this equation. Note also that electrodynamics is considered here in terms of field strengths without appealing to the vector potentials as the primary (input) variables of the theory.

The first step in our consideration is the unitary relationship (and wide range analogy) between the Dirac equation and weakly generalized Maxwell equations [39, 40].

Our second step is the symmetry principle. On the basis of this principle we introduced in [41, 42] the most symmetrical form of the generalized Maxwell equations which can describe both bosons and fermions because they have (see [41, 42]) both spin 1 and spin 1/2 symmetries. On the other hand, namely these equations are unitarily connected with the Dirac equation. So, we have one more important argument to suggest these equations in order to describe intra-atomic phenomena, i. e. to be the equations of specific intra-atomic classical electrodynamics.

In our third step we refer to Sallhofer, who suggested in [34–36] the possi-

bility of introduction of interaction with external field as the interaction with specific media (a new way of introduction of the interaction into the field equations). Nevertheless, our model of atom (and of electron) [39,40] is essentially different from the Sallhofer's one.

On the basis of these three main ideas we are able to postulate the weakly generalized Maxwell equations as the equations for intra-atomic classical electrodynamics which may work in atomic, nuclear and particle physics on the same level of success as the Dirac equation can do. Below we illustrate it considering hydrogen atom within the classical model.

The interest to the problem of relationship between the Dirac and Maxwell equations dates back to the time of creation of quantum mechanics [2, 6, 7, 10, 11, 24, 25, 27–30]. But the authors of these papers considered the most simple example of free and massless Dirac equation. The interest to this relationship has grown in recent years due to the results [34–36], where the investigations of the case $m_0 \neq 0$ and the interaction potential $\Phi \neq 0$ were started. Another approach was developed in [3, 4, 8, 9, 16, 17, 32, 33], where the quadratic relations between the fermionic and bosonic amplitudes were found and used. In our above mentioned papers [39,42], in publications [19–22, 37, 38] and herein we consider the linear relations between the fermionic and bosonic amplitudes. In [20–22, 37, 38] we have found the relationship between the symmetry properties of the Dirac and Maxwell equations, the complete set of 8 transformations linking these equations, the relationship between the conservation laws for the electromagnetic and spinor fields, and the relationship between the Lagrangians for these fields. Here we summarize our previous results and give some new details (Lamb shift consideration) of the intra-atomic electrodynamics and its application to the hydrogen atom. The possibilities of unification with gravitation are briefly discussed.

2. Weakly Generalized Maxwell Equations with Maximal Symmetry Properties

Consider the Maxwell equations with specific (gradient-type) form of electric and magnetic sources and the symmetry properties of such equations. The corresponding equations for the system of electromagnetic and scalar fields $(\vec{E}, \vec{H}, E^0, H^0)$ have the form:

$$\begin{aligned} \partial_0 \vec{E} &= \text{rot } \vec{H} - \text{grad } E^0, & \partial_0 \vec{H} &= -\text{rot } \vec{E} - \text{grad } H^0, \\ \text{div } \vec{E} &= -\partial_0 E^0, & \text{div } \vec{H} &= -\partial_0 H^0. \end{aligned} \quad (1)$$

The Eqs (1) are nothing more than the weakly generalized Maxwell equations with gradient-like electric and magnetic sources $j_\mu^e = -\partial_\mu E^0$, $j_\mu^{\text{mag}} = -\partial_\mu H^0$,

i. e.

$$\vec{j}_e = \text{grad } E^0, \quad \vec{j}_{\text{mag}} = \text{grad } H^0, \quad \rho_e = -\partial_0 E^0, \quad \rho_{\text{mag}} = -\partial_0 H^0. \quad (2)$$

In terms of complex 4-component object $\mathcal{E} \equiv (\mathcal{E}^\mu) = E - iH$

$$\mathcal{E} \equiv \begin{bmatrix} \vec{\mathcal{E}} \\ \mathcal{E}^0 \end{bmatrix} = [E^1 - iH^1, E^2 - iH^2, E^3 - iH^3, E^0 - iH^0]^\top, \quad (3)$$

where $\vec{\mathcal{E}} = \vec{E} - i\vec{H}$ is the well-known complex form for the electromagnetic field used by Majorana in far 1930 (see, e. g. [28]) Eqs (1) may be rewritten in the manifestly covariant form

$$\partial_\mu \mathcal{E}_\nu - \partial_\nu \mathcal{E}_\mu + i\varepsilon_{\mu\nu\rho\sigma} \partial^\rho \mathcal{E}^\sigma = 0, \quad \partial_\mu \mathcal{E}^\mu = 0. \quad (4)$$

It is useful also to consider the following form of Eqs (1), (4):

$$(i\partial_0 - \vec{S} \cdot \vec{p}) \vec{\mathcal{E}} - i \text{grad } \mathcal{E}^0 = 0, \quad \partial_\mu \mathcal{E}^\mu = 0, \quad (5)$$

where $\vec{S} \equiv (S^j)$ are the generators of the irreducible representation $D(1)$ of the group $SU(2)$, i. e. the quantum-mechanical spin 1 operators:

$$S^1 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{bmatrix}, \quad S^2 = \begin{bmatrix} 0 & 0 & i \\ 0 & 0 & 0 \\ -i & 0 & 0 \end{bmatrix}, \quad S^3 = \begin{bmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad (6)$$

$$\vec{S}^2 = 1(1+1)I.$$

Equations (1-5) are directly connected with the free massless Dirac equation

$$i\gamma^\mu \partial_\mu \Psi(x) = 0. \quad (7)$$

The substitution of

$$\psi = \begin{bmatrix} E^3 + iH^0 \\ E^1 + iE^2 \\ E^0 + iH^3 \\ -H^2 + iH^1 \end{bmatrix} = U\mathcal{E}, \quad U = \begin{bmatrix} 0 & 0 & C_+ & C_- \\ C_+ & iC_+ & 0 & 0 \\ 0 & 0 & C_- & C_+ \\ C_- & iC_- & 0 & 0 \end{bmatrix}, \quad (8)$$

$$C_\mp \equiv \frac{1}{2}(C \mp 1), \quad C\Psi \equiv \Psi^*, \quad C\mathcal{E} \equiv \mathcal{E}^*$$

into the Dirac equation (7) with γ matrices in standard Pauli-Dirac representation guarantees its transformation into the generalized Maxwell equations (1-5). The complete set of 8 transformations like (8), which relate generalized Maxwell equations (4) and massless Dirac equation (7), was found in [37, 38]. Unitary relationship between the generalized Maxwell equations (4) and massless Dirac equation (7) can be found in our papers [20, 22].

The unitarity of the operator (8) can be verified easily by taking into account that the equations

$$(AC)^\dagger = CA^\dagger, \quad aC = Ca^*, \quad (aC)^* = Ca \quad (9)$$

hold for an arbitrary matrix A and a complex number a . We note that in the real algebra (i. e. the algebra over the field of real numbers) and in the Hilbert space of quantum mechanical amplitudes this operator has all properties of unitarity. Equations (4) (or their another representations (1-5)) are the maximally symmetrical form of the generalized Maxwell equations. We consider here representation (4) as an example. The following theorem is valid.

Theorem 1. *The generalized Maxwell equations (4) are invariant with respect to the three different transformations, which are generated by three different representations P^V , P^{TS} , P^S of the Poincaré group $P(1,3)$ given by the formulae*

$$\begin{aligned} \mathcal{E}(x) &\rightarrow \mathcal{E}^V(x) = \Lambda \mathcal{E}(\Lambda^{-1}(x - a)) \\ \mathcal{E}(x) &\rightarrow \mathcal{E}^{TS}(x) = F(\Lambda) \mathcal{E}(\Lambda^{-1}(x - a)) \\ \mathcal{E}(x) &\rightarrow \mathcal{E}^S(x) = S(\Lambda) \mathcal{E}(\Lambda^{-1}(x - a)) \end{aligned} \quad (10)$$

where Λ is a vector, i. e. $(\frac{1}{2}, \frac{1}{2})$, $F(\Lambda)$ is a tensor-scalar $(0,1) \otimes (0,0)$ and $S(\Lambda)$ is a spinor representation $(0, \frac{1}{2}) \otimes (\frac{1}{2}, 0)$ of $SL(2, \mathbb{C})$ group. This means that the equations (44) have both spin 1 and spin 1/2 symmetries.

Proof: Let us write the infinitesimal transformations, following from (10), in the form

$$\mathcal{E}^{V,TS,S}(x) = \left(1 - a^\rho \partial_\rho - \frac{1}{2} \omega^{\rho\sigma} j_{\rho\sigma}^{V,TS,S}\right) \mathcal{E}(x). \quad (11)$$

Then the generators of the transformations (11) have the form

$$\partial_\rho = \frac{\partial}{\partial x^\rho}, \quad j_{\rho\sigma}^{V,TS,S} = x_\rho \partial_\sigma - x_\sigma \partial_\rho + s_{\rho\sigma}^{V,TS,S}, \quad (12)$$

where

$$\begin{aligned} (s_{\rho\sigma}^V)^\mu_\nu &= \delta_\rho^\mu g_{\sigma\nu} - \delta_\sigma^\mu g_{\rho\nu}, \quad s_{\rho\sigma}^V \in \left(\frac{1}{2}, \frac{1}{2}\right), \\ s_{\rho\sigma}^{TS} &= \begin{bmatrix} s_{\rho\sigma}^T & 0 \\ 0 & 0 \end{bmatrix} \in (1,0) \oplus (0,0), \quad s_{\rho\sigma}^T = -s_{\sigma\rho}^T, \\ s_{mn}^T &= -i\varepsilon^{mnj} S^j, \quad s_{0j}^T = S^j, \end{aligned} \quad (13)$$

with S^j given by the formula (6) and

$$s_{\rho\sigma}^S = \frac{1}{4}[\tilde{\gamma}_\rho, \tilde{\gamma}_\sigma], \quad \tilde{\gamma} = U^\dagger \gamma U. \quad (14)$$

The unitary operator U is given by the formula (8), $\tilde{\gamma}$ matrices here may be easily found according to (14). The explicit form of $\tilde{\gamma}$ matrices is as follows:

$$\begin{aligned} \tilde{\gamma}^0 &= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix} C, & \tilde{\gamma}^1 &= \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix} C, \\ \tilde{\gamma}^2 &= \begin{bmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & 1 \\ -i & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix} C, & \tilde{\gamma}^3 &= \begin{bmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix} C. \end{aligned} \quad (15)$$

We call the representation (15) the bosonic representation of the γ matrices. In this representation the imaginary unit i is represented by the 4×4 matrix operator.

$$ViV^\dagger = i\Gamma, \quad \Gamma \equiv \begin{bmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix} = \Gamma^\dagger = \Gamma^{-1}, \quad \Gamma^2 = 1. \quad (16)$$

Due to the unitarity of the operator U in (8), the $\tilde{\gamma}^\mu$ matrices not only obey the Clifford-Dirac algebra

$$\tilde{\gamma}^\mu \tilde{\gamma}^\nu + \tilde{\gamma}^\nu \tilde{\gamma}^\mu = 2g^{\mu\nu} \quad (17)$$

but also have the same Hermitian properties as the Pauli-Dirac γ^μ matrices:

$$\tilde{\gamma}^{0\dagger} = \tilde{\gamma}^0, \quad \tilde{\gamma}^{k\dagger} = -\tilde{\gamma}^k. \quad (18)$$

Thus, the formulae (15) give indeed an exotic representation of γ^μ matrices. Now the proof of the theorem is reduced to the verification that all the generators (12) obey the commutation relations of the $P(1,3)$ group and commute with the operator of the generalized Maxwell equations (4-5), which can be rewritten in the Dirac form

$$\tilde{\gamma}^\mu \partial_\mu \mathcal{E}(x) = 0 \quad (19)$$

(for some details see [41,42]). \square

This result about the generalized Maxwell equations (4) means the following. From group theoretical point of view these equations can describe both bosons and fermions. This means that one has direct group-theoretical grounds to apply these equations for the description of electron, as it is presented below in Section 5.

A distinctive feature of the equation (4) for the system $\mathcal{E} = (\vec{\mathcal{E}}, \mathcal{E}^0)$ (i. e. for the system of interacting irreducible $(0, 1)$ and $(0, 0)$ fields) is the following. It is a manifestly covariant equation with minimal number of components, i. e. the equation without redundant components for this system.

Note that each of the three representations (10) of the $P(1, 3)$ group is a local one, because each matrix part of transformations (10) (matrices Λ , $F(\Lambda)$ and $S(\Lambda)$) does not depend on the coordinates $x \in \mathbb{R}^4$, and, consequently, the generators (12) belong to the Lie class of operators. Each of the transformations in (10) may be understood as connected with special relativity transformations in the space-time $\mathbb{R}^4 = \{x\}$, i. e. with transformations in the manifold of inertial frame of references.

We emphasize that the equation (19) has the form of massless Dirac equation for fermionic fields. In such consideration of equation (19) the $\tilde{\gamma}^\mu$ matrices may be chosen in arbitrary representation (e. g., in each of representations of Pauli–Dirac, Majorana, Weyl, etc). However, only in exotic representation (15) equation (19) is the Maxwell equation for the system of interacting electromagnetic $\vec{\mathcal{E}} = \vec{E} - i\vec{H}$ and scalar $\mathcal{E}^0 = E^0 - iH^0$ fields (therefore, we have called the representation (15) the bosonic one). Thus, if one considers the equation (19) as bosonic one, the representation of γ^μ matrices and their explicit form *must be fixed* in the form (15). In the case of bosonic interpretation of Eq. (7) one must fix the explicit form of γ^μ in standard Pauli–Dirac representation and fix the form of Ψ as in (8).

It follows from the Eqs (4) that the field $\mathcal{E} = (\vec{\mathcal{E}}, \mathcal{E}^0)$ is massless, i. e. $\partial^\nu \partial_\nu \mathcal{E}^\mu = 0$. Therefore it is interesting to note that neither P^V , nor P^{TS} symmetries cannot be extended to the local conformal $C(1, 3)$ symmetry. Only the spinor C^S representation of $C(1, 3)$ group, obtained from the local P^S representation, is the symmetry group for the generalized Maxwell equations (4). This fact is understandable: the electromagnetic field $\vec{\mathcal{E}} = \vec{E} - i\vec{H}$ obeying Eqs (4) is not free, it interacts with the scalar field \mathcal{E}^0 .

Consider the particular case of standard (non-generalized) Maxwell equations, namely, the case of equations (4) without magnetic charge and current densities, i. e. the case when $H^0 = 0$ but $E^0 \neq 0$. The symmetry properties of such standard equations are strongly restricted in comparison with the generalized Eqs (4): they are invariant only with respect to tensor-scalar (spin 1 and 0) representations of Poincaré group defined by the corresponding representation

$(0, 1) \otimes (0, 0)$ of the proper orthochronous Lorentz group $SL(2, C)$. Other symmetries mentioned in the theorem are lost for this case too. The proof of this assertion follows from the fact that the vector $\left(\frac{1}{2}, \frac{1}{2}\right)$ and the spinor $\left(0, \frac{1}{2}\right) \oplus \left(\frac{1}{2}, 0\right)$ transformations of $\mathcal{E} = (\vec{\mathcal{E}}, \mathcal{E}^0)$ mix the \mathcal{E}^0 and $\vec{\mathcal{E}}$ components of the field \mathcal{E} , and only the tensor-scalar $(0, 1) \oplus (0, 0)$ transformations do not mix them.

For the free Maxwell equation in vacuum without sources (the case $E^0 = H^0 = 0$) the losing of above mentioned symmetries is evident from the same reasons. Moreover, it is well known that such equations are invariant only with respect to tensor (spin 1) representations of Poincaré and conformal groups and with respect to dual transformation: $\vec{E} \rightarrow \vec{H}, \vec{H} \rightarrow -\vec{E}$. We have obtained the extended 32-dimensional Lie algebra [18] (and the corresponding group) of invariance of free Maxwell equations, which is isomorphic to $C(1, 3) \oplus C(1, 3) \oplus$ dual algebra. We were successful to prove it appealing not to Lie class of symmetry operators but to a more general, namely, to the simplest Lie-Backlund class of operators. The corresponding generalization of symmetries of Eqs (4) presented in the above theorem leads to a wide 246-dimensional Lie algebra in the class of first order Lie-Backlund operators. Thus, the Maxwell equations (4) with electric and magnetic gradient-like sources have the maximally possible symmetry properties among the standard and generalized equations of classical electrodynamics.

The general solution of Eqs (4) was found in [21, 22] directly by the application of Fourier method. In terms of helicity amplitudes $c^\mu(\vec{k})$ this solution has the form

$$\mathcal{E}(x) = \int d^3k \sqrt{\frac{2\omega}{(2\pi)^3}} \left\{ \begin{aligned} & [c^1 e_1 + c^3 (e_3 + e_4)] e^{-ikx} \\ & + [c^{*2} e_1 + c^{*4} (e_3 + e_4)] e^{ikx} \end{aligned} \right\}, \quad (20)$$

where the 4-component basis vectors e_α are taken in the form

$$e_1 = \begin{bmatrix} \vec{e}_1 \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} \vec{e}_2 \\ 0 \end{bmatrix}, \quad e_3 = \begin{bmatrix} \vec{e}_3 \\ 0 \end{bmatrix}, \quad e_4 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}. \quad (21)$$

Here the 3-component basis vectors which, without any loss of generality, can be taken as

$$\vec{e}_1 = \frac{1}{\omega \sqrt{2(k^1 k^1 + k^2 k^2)}} \begin{bmatrix} \omega k^2 - i k^1 k^3 \\ -\omega k^1 - i k^2 k^3 \\ i(k^1 k^1 + k^2 k^2) \end{bmatrix}, \quad \vec{e}_2 = \vec{e}_1^*, \quad \vec{e}_3 = \frac{\vec{k}}{\omega}, \quad (22)$$

are the eigenvectors for the quantum-mechanical helicity operator for the spin $s = 1$.

Note that if the quantities E^0, H^0 in Eqs (1) are some given functions for which the representation

$$E^0 - iH^0 = \int d^3k \sqrt{\frac{2\omega}{(2\pi)^3}} (c^3 e^{-ikx} + c^4 e^{ikx}) \quad (23)$$

is valid, then Eqs (1) are the Maxwell equations with the given sources, $j_\mu^e = -\partial_\mu E^0$, $j_\mu^{\text{mag}} = -\partial_\mu H^0$ (we call these 4 currents the gradient-like sources). In this case the general solution of the Maxwell equations (1-4) with the given sources, as follows from (20), has the form

$$\begin{aligned} \vec{E}(x) &= \int d^3k \sqrt{\frac{\omega}{2(2\pi)^3}} (c^1 \vec{e}_1 + c^2 \vec{e}_2 + \alpha \vec{e}_3) e^{-ikx} + \text{c.c.} \\ \vec{H}(x) &= i \int d^3k \sqrt{\frac{\omega}{2(2\pi)^3}} (c^1 \vec{e}_1 - c^2 \vec{e}_2 + \beta \vec{e}_3) e^{-ikx} + \text{c.c.} \end{aligned} \quad (24)$$

where the amplitudes of longitudinal waves $\vec{e}_3 \exp(-ikx)$ are $\alpha = c^3 + c^4$, $\beta = c^3 - c^4$ and c^3, c^4 are determined by the functions E^0, H^0 according to the Eq. (23).

The longitudinal electromagnetic waves were the object of long time investigation by Hovorostenko [13]. Here we are able (i) to add to his results the exact solution of the Maxwell equations with gradient-like sources which contains such a waves and (ii) to make the assertion about location of these waves in the same space-time area where the gradient-like sources are located (the reason: the amplitudes c^3, c^4 which define this waves and the gradient-like sources are the same).

Now, knowing the operator U from (8), it is easy to obtain the relationship between the amplitudes $a^r(\vec{k}), b^r(\vec{k})$ determining the well known fermionic solution of the massless Dirac equation, and the amplitudes $c^\alpha(\vec{k})$, determining the bosonic solution (20). Corresponding formulae relating fermionic and bosonic amplitudes were found in [21, 22]. Therefore, the fermionic states may be constructed over bosonic states. This assertion finishes the consideration of group-theoretical grounds of our model.

3. New Classical Electrodynamical Hydrogen Atom Model

The generalized Maxwell equations (1) may be extended on the case of specific inneratomic medium. Namely these equations are put into the ground of our model of the atom.

Consider the weakly generalized Maxwell equations with gradient-type sources in a medium:

$$\begin{aligned} \operatorname{rot} \vec{H} - \partial_0 \varepsilon \vec{E} &= \vec{j}_e, & \operatorname{rot} \vec{E} + \partial_0 \mu \vec{H} &= \vec{j}_{\text{mag}}, \\ \operatorname{div} \varepsilon \vec{E} &= \rho_e, & \operatorname{div} \mu \vec{H} &= \rho_{\text{mag}}, \end{aligned} \quad (25)$$

where \vec{E} and \vec{H} are the electromagnetic field strengths, ε and μ are the electric and magnetic permeabilities of the medium being the same as in the electro-dynamical hydrogen atom model of Sallhofer [34, 36, 23]:

$$\varepsilon(\vec{x}) = 1 - \frac{\Phi(\vec{x}) + m_0}{\omega}, \quad \mu(\vec{x}) = 1 - \frac{\Phi(\vec{x}) - m_0}{\omega} \quad (26)$$

where $\Phi(\vec{x}) = -Ze^2/r$ (we use the units: $\hbar = c = 1$). The current and charge densities in equations (25) have the form

$$\begin{aligned} \vec{j}_e &= \operatorname{grad} E^0, & \vec{j}_{\text{mag}} &= -\operatorname{grad} H^0, \\ \rho_e &= -\varepsilon \mu \partial_0 E^0 + \vec{E} \operatorname{grad} \varepsilon, & \rho_{\text{mag}} &= \varepsilon \mu \partial_0 H^0 + \vec{H} \operatorname{grad} \mu, \end{aligned} \quad (27)$$

where E^0, H^0 is the pair of functions (two real scalar fields) generating the densities of gradient-like sources.

One can easily see that equations (25) are not standard electro-dynamical equations known from the Maxwell theory. These equations have the additional terms which can be considered as the magnetic current and charge densities — in one possible interpretation, or equations (25) can be considered as the equations for compound system of electromagnetic (\vec{E}, \vec{H}) and scalar E^0, H^0 fields in another possible interpretation.

Contrary to [39, 40], here the equations (25) are solved directly by means of separation of variables method. It is useful to rewrite these equations in the mathematically equivalent form where the sources are maximally simple:

$$\begin{aligned} \operatorname{rot} \vec{H} - \varepsilon \partial_0 \vec{E} &= \vec{j}_e, & \operatorname{rot} \vec{E} + \mu \partial_0 \vec{H} &= \vec{j}_{\text{mag}}, \\ \operatorname{div} \varepsilon \vec{E} &= \tilde{\rho}_e, & \operatorname{div} \mu \vec{H} &= \tilde{\rho}_{\text{mag}}, \end{aligned} \quad (28)$$

where

$$\vec{j}_e = \operatorname{grad} E^0, \quad \vec{j}_{\text{mag}} = -\operatorname{grad} H^0, \quad \tilde{\rho}_e = -\mu \partial_0 E^0, \quad \tilde{\rho}_{\text{mag}} = -\varepsilon \partial_0 H^0. \quad (29)$$

Consider the stationary solutions of equations (28). Assuming the harmonic time dependence for the functions E^0, H^0

$$\begin{aligned} E^0(t, \vec{x}) &= E_A^0(\vec{x}) \cos \omega t + E_B^0(\vec{x}) \sin \omega t, \\ H^0(t, \vec{x}) &= H_A^0(\vec{x}) \cos \omega t + H_B^0(\vec{x}) \sin \omega t, \end{aligned} \quad (30)$$

we are looking for the solutions of equations (28) in the form

$$\begin{aligned}\vec{E}(t, \vec{x}) &= \vec{E}_A(\vec{x}) \cos \omega t + \vec{E}_B(\vec{x}) \sin \omega t, \\ \vec{H}(t, \vec{x}) &= \vec{H}_A(\vec{x}) \cos \omega t + \vec{H}_B(\vec{x}) \sin \omega t.\end{aligned}\quad (31)$$

For the 16 time-independent amplitudes we obtain the following two nonlinked subsystems

$$\begin{aligned}\text{rot } \vec{H}_A - \omega \varepsilon \vec{E}_B &= \text{grad } E_A^0, & \text{rot } \vec{E}_B - \omega \mu \vec{H}_A &= -\text{grad } H_B^0, \\ \text{div } \varepsilon \vec{E}_B &= \omega \mu E_A^0, & \text{div } \mu \vec{H}_A &= -\omega \varepsilon H_B^0,\end{aligned}\quad (32)$$

and

$$\begin{aligned}\text{rot } \vec{H}_B + \omega \varepsilon \vec{E}_A &= \text{grad } E_B^0, & \text{rot } \vec{E}_A + \omega \mu \vec{H}_B &= -\text{grad } H_A^0, \\ \text{div } \varepsilon \vec{E}_A &= -\omega \mu E_B^0, & \text{div } \mu \vec{H}_B &= \omega \varepsilon H_A^0.\end{aligned}\quad (33)$$

Below we consider only the first subsystem (32). It is quite enough because the subsystems (32) and (33) are connected with transformations

$$\begin{aligned}E &\rightarrow H, & H &\rightarrow -E, & \varepsilon E &\rightarrow \mu H, & \mu H &\rightarrow -\varepsilon E, \\ \varepsilon &\rightarrow \mu, & \mu &\rightarrow \varepsilon,\end{aligned}\quad (34)$$

which are the generalizations of duality transformation of free electromagnetic field. Due to this fact the solutions of subsystem (33) can be easily obtained from the solutions of subsystem (32).

Furthermore, it is useful to separate equations (32) into the following subsystems:

$$\begin{aligned}\omega \varepsilon E_B^3 - \partial_1 H_A^2 + \partial_2 H_A^1 + \partial_3 E_A^0 &= 0 \\ \omega \varepsilon H_B^0 - \partial_1 H_A^1 + \partial_2 H_A^2 + \partial_3 H_A^3 &= 0 \\ -\omega \mu E_A^0 - \partial_1 E_B^1 + \partial_2 E_B^2 + \partial_3 E_B^3 &= 0 \\ \omega \mu H_A^3 - \partial_1 E_B^2 + \partial_2 E_B^1 - \partial_3 H_B^0 &= 0\end{aligned}\quad (35)$$

and

$$\begin{aligned}\omega \varepsilon E_B^1 - \partial_1 H_A^3 + \partial_3 H_A^2 + \partial_1 E_A^0 &= 0 \\ \omega \varepsilon E_B^2 - \partial_3 H_A^1 + \partial_1 H_A^3 + \partial_2 E_A^0 &= 0 \\ \omega \mu H_A^1 - \partial_2 E_B^3 + \partial_3 E_B^2 - \partial_1 H_B^0 &= 0 \\ \omega \mu H_A^2 - \partial_3 E_B^1 + \partial_1 E_B^3 - \partial_2 H_B^0 &= 0.\end{aligned}\quad (36)$$

Assuming the spherical symmetry case, when $\Phi(\vec{x}) = \Phi(r)$, $r \equiv |\vec{x}|$, we are making the transition into the spherical coordinate system and looking for the

solutions in the spherical coordinates in the form

$$(E, H)(\vec{r}) = R_{(E,H)}(r) f_{(E,H)}(\theta, \phi), \quad (37)$$

where $E \equiv (\vec{E}, E^0)$, $H \equiv (\vec{H}, H^0)$. We choose for the subsystem (35) the d'Alembert Ansatz in the form

$$\begin{aligned} \bar{E}_A^0 &= \bar{C}_{E_4} R_{H_4} P_{l_{H_4}}^{\bar{m}_4} e^{-i\bar{m}_4\phi} \\ \bar{E}_B^k &= \bar{C}_{E_k} R_{E_k} P_{l_{E_k}}^{\bar{m}_k} e^{-i\bar{m}_k\phi} \\ \bar{H}_B^0 &= \bar{C}_{H_4} R_{E_4} P_{l_{E_4}}^{\bar{m}_4} e^{-i\bar{m}_4\phi} \\ \bar{H}_A^k &= \bar{C}_{H_k} R_{H_k} P_{l_{H_k}}^{\bar{m}_k} e^{-i\bar{m}_k\phi} \end{aligned} \quad k = 1, 2, 3. \quad (38)$$

We use the following representation for $\partial_1, \partial_2, \partial_3$ operators in spherical coordinates

$$\begin{aligned} \partial_1 CRP_l^m e^{\mp im\phi} &= \frac{e^{\mp im\phi} C}{2l+1} \cos\phi (R_{,l+1} P_{l-1}^{m+1} - R_{,-l} P_{l+1}^{m+1}) \\ &\quad + e^{\mp i(m-1)\phi} C \frac{m}{\sin\theta} P_l^m \frac{R}{r}, \\ \partial_2 CRP_l^m e^{\mp im\phi} &= \frac{e^{\mp im\phi} C}{2l+1} \sin\phi (R_{,l+1} P_{l-1}^{m+1} - R_{,-l} P_{l+1}^{m+1}) \\ &\quad \mp e^{\mp i(m-1)\phi} C \frac{im}{\sin\theta} P_l^m \frac{R}{r}, \\ \partial_3 CRP_l^m e^{\mp im\phi} &= \frac{e^{\mp im\phi} C}{2l+1} (R_{,l+1} (l+m) P_{l-1}^m + R_{,-l} (l-m+1) P_{l+1}^m). \end{aligned} \quad (39)$$

Substitutions (38) and (39) together with the assumptions

$$\begin{aligned} R_{E_\alpha} &= R_E, & l_{E_\alpha} &= l_E, & R_{H_\alpha} &= R_H, & l_{H_\alpha} &= l_H, \\ \bar{m}_1 &= \bar{m}_2 = \bar{m}_3 - 1 = \bar{m}_4 - 1 = m, \\ \bar{C}_{H_1} &= i\bar{C}_{H_2}, & \bar{C}_{E_2} &= -i\bar{C}_{E_1}, & \bar{C}_{H_4} &= -i\bar{C}_{E_3}, & \bar{C}_{H_3} &= -i\bar{C}_{E_4}, \\ \bar{C}_{H_2}^I &= \bar{C}_{E_4}^I (l_H^I + m + 1), & \bar{C}_{E_3}^I &= -\bar{C}_{E_4}^I \equiv \bar{C}^I, \\ \bar{C}_{E_1}^I &= \bar{C}_{E_3}^I (l_E^I - m), & l_H^I &= l_E^I - 1 \equiv l^I, \\ \bar{C}_{H_2}^{II} &= -\bar{C}_{E_4}^{II} (l_H^{II} - m), & \bar{C}_{E_3}^{II} &= -\bar{C}_{E_4}^{II} \equiv \bar{C}^{II}, \\ \bar{C}_{E_1}^{II} &= -\bar{C}_{E_3}^{II} (l_E^{II} + m + 1), & l_H^{II} &= l_E^{II} + 1 \equiv l^{II} \end{aligned} \quad (40)$$

into the subsystem (35) guarantee the separation of variables in these equations and lead to a pair of equations for the two radial functions R_E, R_H (for the

subsystem (36) the procedure is similar):

$$\varepsilon\omega R_E^I - R_{H,-l}^I = 0, \quad \mu\omega R_H^I + R_{E,l+2}^I = 0, \quad (41)$$

$$\varepsilon\omega R_E^{II} - R_{H,l+1}^{II} = 0, \quad \mu\omega R_H^{II} + R_{E,-l+1}^{II} = 0, \quad (42)$$

$$R_{,a} \equiv \left(\frac{d}{dr} + \frac{a}{r} \right) R.$$

For the case $\Phi = -ze^2/r$ the equations (41, 42) coincide exactly with the radial equations for the hydrogen atom of the Dirac theory and, therefore, the procedure of their solution is the same as in textbooks on relativistic quantum mechanics. It leads to the well-known Sommerfeld–Dirac formula for the fine structure of the hydrogen spectrum. We note only that here the discrete picture of energy spectrum in the domain $0 < \omega < m_0c^2$ is guaranteed by the demand for the solutions of the radial equations (41), (42) to decrease at infinity (when $r \rightarrow \infty$). From the equations (41), (42) and this condition the Sommerfeld–Dirac formula

$$\omega = \omega_{nj}^{\text{hyd}} = \frac{m_0c^2}{\hbar \sqrt{1 + \frac{\alpha^2}{(n_r + \sqrt{k^2 - \alpha^2})^2}}} \quad (43)$$

follows, where the notations of the Dirac theory (see, e. g. [1]) are used: $n_r = n - k$, $k = j + 1/2$, $\alpha = e^2/\hbar c$. Let us note once more that the result (43) is obtained here not from the Dirac equation, but from the Maxwell equations (25) with sources (27) in the medium (26).

Substituting (40) into (38) one can easily obtain the angular part of the hydrogen solutions for the $(\vec{E}, \vec{H}, E^0, H^0)$ field and calculate according to (27) the corresponding currents and charges. Let us write down the explicit form for the set of electromagnetic field strengths (\vec{E}, \vec{H}) , which are the hydrogen solutions of equations (25), and also for the currents and charges generating these field strengths (the complete set of solutions is represented in [39, 40]):

$$\vec{E}^I = R_E^I \begin{bmatrix} (-l + m - 1)P_{l+1}^m \cos m\phi \\ (l - m + 1)P_{l+1}^m \sin m\phi \\ -P_{l+1}^{m+1} \cos(m+1)\phi \end{bmatrix}, \quad \vec{H}^I = R_H^I \begin{bmatrix} (l + m + 1)P_l^m \sin m\phi \\ (l + m + 1)P_l^m \cos m\phi \\ -P_l^{m+1} \sin(m+1)\phi \end{bmatrix}$$

$$\vec{j}_e^I = \text{grad } R_H^I P_l^{m+1} \cos(m+1)\phi$$

$$\vec{j}_{\text{mag}}^I = -\text{grad } R_E^I P_{l+1}^{m+1} \sin(m+1)\phi \quad (44)$$

$$\rho_e^I = -(\varepsilon R_E^I)_{,l+2} P_l^{m+1} \cos(m+1)\phi$$

$$\rho_{\text{mag}}^I = -(\mu R_H^I)_{,-l} P_{l+1}^{m+1} \sin(m+1)\phi$$

$$\begin{aligned}
\vec{E}^{II} &= R_E^{II} \begin{bmatrix} (l+m)P_{l-1}^m \cos m\phi \\ (-l-m)P_{l-1}^m \sin m\phi \\ P_{l-1}^{m+1} \cos(m+1)\phi \end{bmatrix}, & \vec{H}^{II} &= R_H^{II} \begin{bmatrix} (-l+m)P_l^m \sin m\phi \\ (-l+m)P_l^m \cos m\phi \\ -P_l^{m+1} \sin(m+1)\phi \end{bmatrix}, \\
\vec{j}_e^{II} &= \text{grad } R_H^{II} P_l^{m+1} \cos(m+1)\phi \\
\vec{j}_{\text{mag}}^{II} &= -\text{grad } R_E^{II} P_{l-1}^{m+1} \sin(m+1)\phi \\
\rho_e^{II} &= -(\varepsilon R_E^{II})_{,-l+1} P_l^{m+1} \cos(m+1)\phi \\
\rho_{\text{mag}}^{II} &= -(\mu R_H^{II})_{,l+1} P_{l-1}^{m+1} \sin(m+1)\phi.
\end{aligned} \tag{45}$$

In one of the possible interpretations the states of the hydrogen atom are described by these field strength functions \vec{E} , \vec{H} generated by the corresponding currents and charge densities.

It is evident from (25) that currents and charges in (44–45) are generated by scalar fields (E^0, H^0). Corresponding to (44–45), the (E^0, H^0) solutions of equations (25) are the following:

$$\begin{aligned}
E^{I0} &= R_H^I P_l^{m+1} \cos(m+1)\phi, & H^{I0} &= R_E^I P_{l+1}^{m+1} \sin(m+1)\phi, \\
E^{II0} &= R_H^{II} P_l^{m+1} \cos(m+1)\phi, & H^{II0} &= R_E^{II} P_{l-1}^{m+1} \sin(m+1)\phi.
\end{aligned} \tag{46}$$

As in quantum theory, the numbers $n = 0, 1, 2, \dots$, $j = k - \frac{1}{2} = l \mp \frac{1}{2}$ ($k = 1, 2, \dots, n$) and $m = -l, -l+1, \dots, l$ mark both the terms (43) and the corresponding exponentially decreasing field functions \vec{E}, \vec{H} (and E^0, H^0) in (44–46), i. e. they mark the different discrete states of the classical electrodynamical field (and the densities of the currents and charges) which by definitions describes the corresponding states of hydrogen atom in the model under consideration.

Note that the radial equations (41), (42) cannot be obtained if one neglects the sources in equations (29), or one (electric or magnetic) of these sources. Moreover, in this case there is no solution effectively concentrated in atomic region.

Now we can show on the basis of this model that the assertions known as *Bohr's postulates are the consequences of equations (25) and of their classical interpretation*, i. e. these assertions can be derived from the model, there is no necessity to postulate them outside the framework of the classical physics as it was in Bohr's theory. To derive the first Bohr's postulate one can calculate the generalized Poynting vector for the hydrogen solutions (44–46), i. e. for the compound system of stationary electromagnetic and scalar fields ($\vec{E}, \vec{H}, E^0, H^0$)

$$\vec{P}_{\text{gen}} = \int d^3x (\vec{E} \times \vec{H} - \vec{E}E^0 - \vec{H}H^0). \tag{47}$$

The straightforward calculations show that not only the vector (47) is identically equal to zero but the Poynting vector itself and the term with scalar fields (E^0, H^0) are also identically equal to zero:

$$\vec{P} = \int d^3x(\vec{E} \times \vec{H}) \equiv 0, \quad \int d^3x(\vec{E}E^0 + \vec{H}H^0) \equiv 0. \quad (48)$$

This means that in stationary states the hydrogen atom does not emit any Poynting radiation neither due to the electromagnetic (\vec{E}, \vec{H}) field, nor to the scalar (E^0, H^0) field. That is the mathematical proof of the first Bohr postulate. The similar calculations of the energy for the same system

$$\mathbf{W} \equiv \frac{1}{2} \int d^3x(\vec{E}^2 + \vec{H}^2 + E_0^2 + H_0^2) \quad (49)$$

give a constant \mathbf{W}_{nl} , depending on n, l (or n, j) and independent of m . In our model this constant is to be identified with the parameter ω in equations (1) which in the stationary states of $(\vec{E}, \vec{H}, E^0, H^0)$ field appears to be equal to the Sommerfeld-Dirac value ω_{nj}^{hyd} (43). By abandoning the $\hbar = c = 1$ system and putting arbitrary A in equations (25) instead of \hbar we obtain final ω_{nj}^{hyd} with A instead of \hbar . Then the numerical value of \hbar can be obtained by comparison of ω_{nj}^{hyd} containing A with the experiment. These facts complete the proof of the second Bohr postulate.

This result means that in this model the Bohr postulates are no longer postulates, but direct consequences of the classical electro-dynamical equation (25). Moreover, this means that together with Dirac or Schrödinger equations we have now the new equation which can be used for finding the solutions of atomic spectroscopy problems. In a contrast to the well-known equations of quantum mechanics our equation is classical.

Being aware that a few interpretations of quantum mechanics (e. g. Copenhagen, statistical, Feynman's, Everett's, transactional, see e. g. [5, 15, 43, 31]) exist, we are far from thinking that the interpretation here can be the only one. But the main point is that now the classical interpretation (without probabilities) is possible.

Today we prefer the following interpretation of hydrogen atom in the approach, when one considers only the motion of electron in the external field of the nucleon. In our model the interacting field of the nucleon and electron is represented by the medium with permeabilities ε, μ given by the formulae (26). The atomic electron is interpreted as the stationary electromagnetic-scalar wave $(\vec{E}, \vec{H}, E^0, H^0)$ in medium (26), i. e. as the stationary electromagnetic wave interacting with massless scalar fields (E^0, H^0) , or with complex massless scalar

field $\mathcal{E}^0 = E^0 - iH^0$ with spin $s = 0$. In other words, the electron can be interpreted as an object having the structure consisting of a photon and a massless meson with zero spin connected, probably, with leptonic charge. The role of the massless scalar field is the following: it generates the densities of electric and magnetic currents and charges (ρ, \vec{j}) , which are the secondary objects in such model. The mass is the secondary parameter too. There is no electron as an input charged massive corpuscle in this model! The mass and the charge of electron appear only outside such atom according to the law of electromagnetic induction and its gravitational analogy. That is why no difficulties of Rutherford–Bohr’s model (about different models of atom see, e. g. [23]) of atom are present here! The Bohr postulates are shown to be the consequences of the model. This interpretation is based on the hypothesis of bosonic nature of matter (on the speculation of the bosonic structure of fermions) according to which all the fermions can be constructed from different bosons (something like new SUSY theory). Of course, before the experiment intended to observe the structure of electron and before the registration of massless spinless meson it is only the hypothesis but based on the mathematics presented here. We note that such massless spinless boson has many similar features with the Higgs boson and the transition here from intra-atomic (with high symmetry properties) to macroelectrodynamics (with loss of many symmetries) looks similarly to the symmetry breakdown mechanism.

The successors of magnetic monopole can try to develop here the monopole interpretation (see [26] for the review and some new ideas about the monopole) — we note that there are few interesting possibilities of interpretation but we want to mark first of all the mathematical facts which are more important than different ways of interpretation.

4. The Unitary Relationship Between the Relativistic Quantum Mechanics and Classical Electrodynamics in Medium

Let us consider the connection between the stationary Maxwell equations

$$\begin{aligned} \operatorname{rot} \vec{H} - \omega \varepsilon \vec{E} &= \operatorname{grad} E^0, & \operatorname{rot} \vec{E} - \omega \mu \vec{H} &= -\operatorname{grad} H^0, \\ \operatorname{div} \vec{E} &= \omega \mu E^0, & \operatorname{div} \vec{H} &= -\omega \varepsilon H^0, \end{aligned} \quad (50)$$

which follow from the system (32) after omitting indices A, B , and the stationary Dirac equation following from the ordinary Dirac equation

$$(i\gamma^\mu \partial_\mu - m_0 + \gamma^0 \Phi) \Psi = 0, \quad \Psi \equiv (\Psi^\alpha), \quad (51)$$

with $m_0 \neq 0$ and the interaction potential $\Phi \neq 0$. Assuming the ordinary time dependence

$$\Psi(x) = \Psi(\vec{x}) e^{-i\omega t} \implies \partial_0 \Psi(x) = -i\omega \Psi(x), \quad (52)$$

for the stationary states and using the standard Pauli–Dirac representation for the γ matrices, one obtains the following system of equations for the components Ψ^α of the spinor Ψ :

$$\begin{aligned} -i\omega\varepsilon\Psi^1 + (\partial_1 - i\partial_2)\Psi^4 + \partial_3\Psi^3 &= 0, \\ -i\omega\varepsilon\Psi^2 + (\partial_1 + i\partial_2)\Psi^3 - \partial_3\Psi^4 &= 0, \\ -i\omega\mu\Psi^3 + (\partial_1 - i\partial_2)\Psi^2 + \partial_3\Psi^1 &= 0, \\ -i\omega\mu\Psi^4 + (\partial_1 + i\partial_2)\Psi^1 - \partial_3\Psi^2 &= 0, \end{aligned} \quad (53)$$

where ε and μ are the same as in (26). After substitution in Eqs (54) instead of Ψ the following column

$$\Psi = [-H^0 + iE^3, -E^2 + iE^1, E^0 + iH^3, -H^2 + iH^1]^T. \quad (54)$$

one obtains Eqs (50). A complete set of 8 transformations with the same role was obtained with the help of the Pauli–Gursey symmetry operators [14] in our papers [37, 38].

The relationship (54) may be written down in terms of unitary operator similarly to that in (8). Further consideration of unitary relationship between the equations (50) and (53) is similar to the procedure in Section 1 and may be omitted. The details were published in [39, 40].

The mathematical facts considered here prove the one-to-one correspondence between the solutions of the stationary Dirac equation and the stationary Maxwell equations with 4-currents of gradient-like type. Hence, one can, using (54), write down the hydrogen solutions of the Maxwell equations (25) (or (28)) starting from the well-known hydrogen solutions of the Dirac equation (51), i. e. without special procedure of finding the solutions of the Maxwell equations, see [39, 40].

5. Lamb Shift

It is very useful to consider the case of Lamb shift in the approach presented here. This specific quantum electrodynamical effect (as modern theory asserts) can be described here in the framework of classical electrodynamics of media. In order to obtain the Lamb shift one must add to $\Phi(\vec{x}) = -Ze^2/r$ in (2) the

quasipotential (known, e. g. from [12], which follows, of course, from quantum electrodynamics)

$$-\frac{Ze^4}{60\pi^2 m_0^2} \delta(r) \quad (55)$$

and solve the equations (1) for such medium similarly to the procedure of Section 2 (of course, by another approximation method). Finally one obtains the Lamb shift correction to the Sommerfeld–Dirac formula (43). Such Lamb shift can be interpreted as a pure classical electrodynamical effect. It can be considered here as a consequence of polarization of medium (26) and not of polarization of such abstract concept as the vacuum in quantum electrodynamics. This brief consideration of the example mentioned in the title of this section demonstrates that our approach can essentially extend the limits of classical theory application in microworld, which was the main purpose of our investigations.

6. A Brief Remark About Gravity

The unified theory of electromagnetic and gravitational phenomena may be constructed within the approach under consideration in the following way. The main primary equations again are written as (1) and gravity is considered as a medium in these equations, i. e. the electric ε and magnetic μ permeabilities of the medium are some functions of the gravitational potential Φ_{grav} :

$$\varepsilon = \varepsilon(\Phi_{\text{grav}}), \quad \mu = \mu(\Phi_{\text{grav}}). \quad (56)$$

Gravity as a medium may generate all the phenomena which in standard Einstein's gravity are generated by Riemannian geometry. For example, the refraction of the light beam near a big mass star is a typical medium effect in such a unified model of electromagnetic and gravitational phenomena. The idea of such consideration consists in the following. The gravitational interaction between massive objects may be represented as the interaction with some medium, similarly as in Eqs (25) the electromagnetic interaction between charged particles.

7. Conclusions

One of the conclusions of our investigation presented here and in [41, 42, 18, 19] is that a field equation itself does not answer the question what kind of particles (Bose or Fermi) are described by this equation. To answer this question one needs to find all representations of the Poincaré group under which the equation is invariant. If more than one such Poincaré representations are found

[41, 42, 18, 19], including the representations with integer and half-integer spins, then the given equation describes both Bose and Fermi particles, and both quantization types (Bose and Fermi) [21, 22] of the field function, obeying this equation, satisfy the microcausality condition. The strict group-theoretical ground of this assertion is the following [41, 42]: both weakly generalized Maxwell equations (1) (with $\varepsilon = \mu = 1$) and Dirac equation (7) (with $m_0 = 0$, $\Phi = 0$) are invariant with respect to three different local representations of Poincaré group, namely the standard spinor, vector and tensor-scalar representations generating by the $(0, \frac{1}{2}) \otimes (\frac{1}{2}, 0)$, $(\frac{1}{2}, \frac{1}{2})$, $(0, 1) \otimes (0, 0)$ representations of the Lorentz $SL(2, \mathbb{C})$ group, respectively.

A few words can be said about the interpretation of the Dirac Ψ function. As follows from the consideration presented here, e. g. from the relationships (8) and (54), the new interpretation of the Dirac Ψ function can be suggested too: Ψ function is the combination of the electromagnetic field strengths (\vec{E}, \vec{H}) and two scalar fields (E^0, H^0) generating the electromagnetic sources, i. e. in this case the probability or Copenhagen interpretation of the function Ψ is not necessary.

The main conclusion is that the limit of classical theory application in the microworld may be essentially extended on the basis of equations and models which are considered here. Lamb shift may be explained as the classical effect in the framework of the classical model of atom.

Acknowledgements

The authors are much grateful to Prof. Hans Sallhofer for numerous discussions of the details and essential support of the main idea, and to Prof. Boris Struminsky for useful discussions of alternatives in gravity.

This work is supported by the National Fund of Fundamental Researches of Ukraine, grant #02.07/253.

References

- [1] Bethe H. and Salpeter E., *Quantum Mechanics of One- and Two-Electron Atoms*, Springer, Berlin 1957.
- [2] Borhgardt A., *Wave Equations for the Photon*, Sov. Phys. JETP **34**(2) (1958) 334–341.
- [3] Campolattaro A., *New Spinor Representation of Maxwell Equations*, Int. J. Theor. Phys. **19**(2) (1980) 99–126.
- [4] Campolattaro A., *Generalized Maxwell Equations and Quantum Mechanics*, Internat. J. Theor. Phys. **29**(2) (1990) 141–155.

-
- [5] Cramer J., *The Transactional Interpretation of Quantum Mechanics*, Rev. Mod. Phys. **58**(3) (1986) 647–687.
- [6] Da Silveira A., *Dirac-like Equation for the Photon*, Z. Naturforsch **A34** (1979) 646–647.
- [7] Darwin C., *The Wave Equations of the Electron*, Proc. Roy. Soc. London **A118** (1928) 654–680.
- [8] Daviau C., *Electromagnetisme, monopoles magnetiques at ondes de matiere dans l'algebre d'espace-temps*, Ann. Fondation. Louis de Broglie **14**(3) (1989) 273–390.
- [9] Daviau C. and Lochak G., *Sur un modele d'equation spinorielle non lineaire*, Ann. Fondation. Louis de Broglie **16**(1) (1991) 43–71.
- [10] Giannetto E., *A Majorana–Oppenheimer Formulation of Quantum Electrodynamics*, Lett. Nuov. Cim. **44**(3) (1985) 140–144.
- [11] Good R., *Particle Aspect of the Electromagnetic Field Equations*, Phys. Rev. **105**(6) (1957) 1914–1919.
- [12] Halzen F. and Martin A., *Quarks and Leptons*, John Wiley & Sons, New York 1984.
- [13] Hvorostenko N., *Longitudinal Electromagnetic Waves*, Sov. Phys. J. (Izvestiya Vuzov) Ser. Physics. #7 (1992) 24–29.
- [14] Ibragimov N., *Invariant Variation Problems and Conservation Laws*, Theor. Math. Phys. **1**(3) (1969) 350–359.
- [15] Jammer M., *The Philosophy of Quantum Mechanics. The Interpretations of Quantum Mechanics in Historical Perspective*, Wiley, New York 1974.
- [16] Keller J., *On the Electron Theory*, Proc. of the Int. Conf. “The Theory of Electron”, Mexico 24–27 September 1995, Adv. Appl. Cliff. Algebras **7** (Special) (1997) 3–26.
- [17] Keller J., *The Geometric Content of the Electron Theory*, Adv. Appl. Cliff. Algebras **9**(2) (1999) 309–395.
- [18] Krivsky I. and Simulik V., *Lagrangian for Electromagnetic Field in the Terms of Field Strengths and Conservation Laws*, Ukrainian Phys. J. **30**(10) (1985) 1457–1459.
- [19] Krivsky I. and Simulik V., *Foundations of Quantum Electrodynamics in Field Strengths Terms*, Naukova Dumka, Kiev 1992.
- [20] Krivsky I. and Simulik V., *The Dirac Equation and Spin 1 Representation. Relationship with the Symmetries of the Maxwell Equation*, Theor. Math. Phys. **90**(3) (1992) 388–406.
- [21] Krivsky I. and Simulik V., *Unitary Connection in Maxwell–Dirac Isomorphism and the Clifford Algebra*, Adv. Appl. Cliff. Alg. **6**(2) (1996) 249–259.
- [22] Krivsky I. and Simulik V., *The Maxwell Equations with Gradient-type Currents and their Relationship with the Dirac Equation*, Ukrainian Phys. J., **44**(5) (1999) 661–665.
- [23] Lakhtakia A., *Models and Modelers of Hydrogen*, World Scientific, London 1996.

- [24] Laporte O. and Uhlenbeck G., *Application of Spinor Analysis to the Maxwell and Dirac Equations*, Phys. Rev. **37** (1931) 1380–1397.
- [25] Ljolje K., *Some Remarks on Variational Formulations of Physical Fields*, Fortschr. Phys. **36**(1) (1988) 9–32.
- [26] Lochak J., *The Symmetry Between Electricity and Magnetism and the Problem of the Existence of the Magnetic Monopole* In: *Advanced Electromagnetism*, World Scientific, London 1995 pp 105–148.
- [27] Lomont J., *Dirac-like Wave Equations for Particles of Zero Rest Mass and their Quantization*, Phys. Rev. **111**(6) (1958) 1710–1716.
- [28] Mignani R., Recami E. and Baldo M., *About a Dirac-like Equation for the Photon According to Ettore Majorana*, Lett. Nuovo Cimento **11**(12) (1974) 572–586.
- [29] Moses H., *A Spinor Representation of Maxwell's Equations*, Nuovo Cimento Suppl. **7**(1) (1958) 1–18.
- [30] Oppenheimer J., *Note on Light Quanta and the Electromagnetic Field*, Phys. Rev. **38** (1931) 725–746.
- [31] Paty M., *Are Quantum Systems Physical Objects with Physical Properties*, Eur. J. Phys. **20** (1999) 373–388.
- [32] Rodrigues W. Jr. and de Oliveira E., *Dirac and Maxwell equations in the Clifford and Spin-Clifford Bundles*, Int. J. Theor. Phys. **29**(4) (1990) 397–412.
- [33] Rodrigues W. Jr. and Vaz J. Jr., *From Electromagnetism to Relativistic Quantum Mechanics*, Found. Phys. **28**(5) (1998) 789–814.
- [34] Sallhofer H., *Elementary Derivation of the Dirac Equation I*, Z. Naturforsch **A33** (1978) 1379–1381.
- [35] Sallhofer H., *Hydrogen in Electrodynamics. VI. The General Solution*, Z. Naturforsch **A45** (1990) 1361–1366.
- [36] Sallhofer H. and Radharose D., *Here Erred Einstein*, World Scientific, London 2001.
- [37] Simulik V., *Relationship Between the Symmetry Properties of the Dirac and Maxwell Equations. Conservation Laws*, Theor. Math. Phys. **87** (1991) 76–85.
- [38] Simulik V., *Some Algebraic Properties of Maxwell-Dirac Isomorphism*, Z. Naturforsch **A49** (1994) 1074–1076.
- [39] Simulik V., *Solutions of the Maxwell Equations Describing the Spectrum of Hydrogen*, Ukrainian Mathematical J. **49**(7) (1997) 1075–1088.
- [40] Simulik V. and Krivsky I., *Clifford Algebra in Classical Electrodynamical Hydrogen Atom Model*, Adv. Appl. Cliff. Algebras **7**(1) (1997) 25–34.
- [41] Simulik V. and Krivsky I., *Fermionic Symmetries of the Maxwell Equations with Gradient-like Sources*, Proc. of the Int. Conf. “Symmetry in Nonlinear Mathematical Physics”, Kiev 7–13 July, 1997 p. 475–482.
- [42] Simulik V. and Krivsky I., *Bosonic Symmetries of the Massless Dirac Equation*, Adv. Appl. Cliff. Algebras **8**(1) (1998) 69–82.
- [43] Sudbery A., *Quantum Mechanics and the Particles of Nature*, Cambridge University Press, Cambridge 1986.